

## $L_p$ -MIXED AFFINE SURFACE AREA

TIAN LI AND WEIDONG WANG

(Communicated by M. A. Hernandez Cifre)

*Abstract.* Lutwak introduced the notion of  $L_p$ -affine surface area by  $L_p$ -mixed volume and obtained some related inequalities. In this article, based on the  $L_p$ -mixed quermassintegrals, we define the concept of the  $L_p$ -mixed affine surface area and extend some of Lutwak's result.

### 1. Introduction

We say that  $K$  is a convex body if  $K$  is a compact, convex subset in  $n$ -dimensional Euclidean space  $\mathbb{R}^n$  with non-empty interior. The set of all convex bodies in  $\mathbb{R}^n$  is written as  $\mathcal{K}^n$ , and its subset  $\mathcal{K}_o^n$  denotes the set of convex bodies containing the origin in their interiors. Similarly,  $\mathcal{K}_c^n$  denotes the set of convex bodies with centroid at the origin. Let  $\mathcal{F}^n$  ( $\mathcal{F}_o^n$ ) denote the subset of  $\mathcal{K}^n$  ( $\mathcal{K}_o^n$ ) that have a positive continuous curvature function. Besides  $\mathcal{S}_o^n$  denotes the set of star bodies (with respect to the origin) and  $\mathcal{S}_c^n$  denotes the set of star bodies whose centroid lie at the origin in  $\mathbb{R}^n$ . Let  $S^{n-1}$  denote the unit sphere in  $\mathbb{R}^n$  and  $V(K)$  denote the  $n$ -dimensional volume of the body  $K$ , for the standard unit ball  $B$  in  $\mathbb{R}^n$ , denote  $\omega_n = V(B)$ .

The notion of classical affine surface area was defined first by Blaschke ([1]). For a smooth convex body  $K$  in  $\mathbb{R}^3$ , the affine surface area,  $\Omega(K)$ , of  $K$  is given by

$$\Omega(K) = \int_{S^{n-1}} f(K, u)^{3/4} dS(u),$$

where  $f(K, \cdot)$  denotes the curvature function of  $K \in \mathcal{K}^n$ , and  $dS(\cdot)$  denotes the infinitesimal of Lebesgue measure  $S(\cdot)$  on the unit sphere  $S^{n-1}$ . Later,  $\Omega(K)$  was naturally considered for sufficiently smooth  $K$  in  $\mathcal{K}^n$  by Leichtweiss ([6]) as

$$\Omega(K) = \int_{S^{n-1}} f(K, u)^{\frac{n}{n+1}} dS(u).$$

In 1989, Leichtweiß ([7]) extended the domain of  $\Omega$ :  $\mathcal{F}^n \rightarrow (0, +\infty)$  from  $\mathcal{F}^n$  to  $\mathcal{K}^n$  as follows: For  $K \in \mathcal{K}^n$ , the affine surface area,  $\Omega(K)$ , of  $K$  is defined by

$$n^{-\frac{1}{n}} \Omega(K)^{\frac{n+1}{n}} = \inf\{nV_1(K, Q^*)V(Q)^{\frac{1}{n}} : Q \in \mathcal{S}_o^n\}.$$

*Mathematics subject classification* (2010): 52A20, 52A40.

*Keywords and phrases:*  $L_p$ -affine surface area,  $L_p$ -mixed affine surface area,  $L_p$ -mixed quermassintegrals.

Research is supported in part by the Natural Science Foundation of China (No. 11371224) and Innovation Foundation of Graduate Student of China Three Gorges University (No. SDYC2016118).

Here  $Q^*$  denotes the polar body of  $Q$  and  $V_1(M, N)$  denotes the mixed volume of convex bodies  $M$  and  $N$ .

Based on the classical affine surface area, Lutwak (see [12]) introduced the classical notion of mixed affine surface area and obtained some isoperimetric inequalities for this notion. During the past three decades, the investigations of the classical affine surface area have received great attention from many articles (see [2, 3, 7, 8, 10, 11, 12, 13, 14, 15, 16, 17, 20, 34, 35, 36]).

In 1996, according to the  $L_p$ -mixed volume, Lutwak ([15]) introduced the notion of  $L_p$ -affine surface area. For  $K \in \mathcal{K}_o^n$ ,  $p \geq 1$ , the  $L_p$ -affine surface area,  $\Omega_p(K)$ , of  $K$  is defined by

$$n^{-\frac{p}{n}}\Omega_p(K)^{\frac{n+p}{n}} = \inf\{nV_p(K, Q^*)V(Q)^{\frac{p}{n}} : Q \in \mathcal{S}_o^n\}.$$

Here  $V_p(M, N)$  denotes the  $L_p$ -mixed volume of  $M, N \in \mathcal{K}_o^n$ . Obviously, for  $p = 1$ ,  $\Omega_p(K)$  is the classical affine surface area  $\Omega(K)$ .

In 2007, Wang and Leng ([24]) introduced the notion of  $i$ th  $L_p$ -mixed affine surface area and obtained results related to it. Regarding the studies of the  $L_p$ -affine surface areas also see ([21, 22, 25, 26, 27, 28, 30, 31, 32, 33]).

Based on the definition of  $L_p$ -affine surface area, Lutwak ([15]) proved the following results:

**THEOREM 1.A<sub>a</sub>.** *If  $p \geq 1$  and  $K \in \mathcal{K}_o^n$ , then*

$$\left[ \frac{\Omega_p(K)^{n+p}}{n^{n+p}V(K)^{n-p}} \right]^{\frac{1}{p}} \leq V(K)V(K^*). \tag{1.1}$$

Here  $\left[ \frac{\Omega_p(K)^{n+p}}{n^{n+p}V(K)^{n-p}} \right]^{\frac{1}{p}}$  is called the  $L_p$ -affine surface area ratio of  $K$ .

**THEOREM 1.A<sub>b</sub>.** *If  $p \geq 1$  and  $K \in \mathcal{F}_o^n$ , then*

$$\left[ \frac{\Omega_p(K)^{n+p}}{n^{n+p}V(K)^{n-p}} \right]^{\frac{1}{p}} \leq V(K)V(K^*), \tag{1.2}$$

with equality if and only if  $K^*$  and  $\Lambda_p K$  are dilates.

**THEOREM 1.B<sub>a</sub>.** *If  $K \in \mathcal{K}_o^n$ ,  $1 \leq p < q$ , then*

$$\left[ \frac{\Omega_p(K)^{n+p}}{n^{n+p}V(K)^{n-p}} \right]^{\frac{1}{p}} \leq \left[ \frac{\Omega_q(K)^{n+q}}{n^{n+q}V(K)^{n-q}} \right]^{\frac{1}{q}}. \tag{1.3}$$

**THEOREM 1.B<sub>b</sub>.** *If  $K \in \mathcal{F}_o^n$ ,  $1 \leq p < q$ , then*

$$\left[ \frac{\Omega_p(K)^{n+p}}{n^{n+p}V(K)^{n-p}} \right]^{\frac{1}{p}} \leq \left[ \frac{\Omega_q(K)^{n+q}}{n^{n+q}V(K)^{n-q}} \right]^{\frac{1}{q}}, \tag{1.4}$$

with equality if and only if  $\Lambda_p K$  and  $\Lambda_q K$  are dilates.

THEOREM 1.C<sub>a</sub>. If  $K \in \mathcal{K}_o^n$ ,  $1 \leq p < q < r$ , then

$$\Omega_q(K)^{(n+q)(r-p)} \leq \Omega_p(K)^{(n+p)(r-q)} \Omega_r(K)^{(n+r)(q-p)}. \tag{1.5}$$

THEOREM 1.C<sub>b</sub>. If  $K \in \mathcal{F}_o^n$ ,  $1 \leq p < q < r$ , then

$$\Omega_q(K)^{(n+q)(r-p)} \leq \Omega_p(K)^{(n+p)(r-q)} \Omega_r(K)^{(n+r)(q-p)}. \tag{1.6}$$

with equality if and only if  $\Lambda_p K$  and  $\Lambda_r K$  are dilates.

The main purpose of this article is to define the notion of  $L_p$ -mixed affine surface area by Lutwak’s  $L_p$ -mixed quermassintegrals (see [14]). Then by this notion we establish some inequalities which are extensions of parts of Lutwak’s results in [12] and [15].

Associated with the  $L_p$ -mixed quermassintegrals (see [14]), we first give the notion of  $L_p$ -mixed affine surface area as follows:

DEFINITION 1.1. For  $K \in \mathcal{K}_o^n$ ,  $p \geq 1$  and  $i = 0, 1, \dots, n - 1$ , the  $L_p$ -mixed affine surface area,  $\Omega_{p,i}(K)$ , of  $K$  is defined by

$$n^{-\frac{p}{n-i}} \Omega_{p,i}(K)^{\frac{n+p-i}{n-i}} = \inf \{ n W_{p,i}(K, Q^*) \widetilde{W}_i(Q)^{\frac{p}{n-i}} : Q \in \mathcal{S}_o^n \}. \tag{1.7}$$

Here  $W_{p,i}(M, N)$  denotes the  $L_p$ -mixed quermassintegrals of  $M, N \in \mathcal{K}_o^n$ .

Note that above definition is different from the notion of  $i$ th  $L_p$ -mixed affine surface area in ([24])

We easily see that if  $i = 0$  in (1.7), then  $\Omega_{p,i}(K)$  is just the  $L_p$ -affine surface area.

Further, we establish some inequalities for the  $L_p$ -mixed affine surface area which extend the results of Theorems 1.A<sub>a</sub>-1.C<sub>b</sub>. Our main results can be stated as follows:

THEOREM 1.1<sub>a</sub>. If  $K \in \mathcal{K}_o^n$ ,  $p \geq 1$  and  $i = 0, 1, \dots, n - 1$ , then

$$\left[ \frac{\Omega_{p,i}(K)^{n+p-i}}{n^{n+p-i} W_i(K)^{n-p-i}} \right]^{\frac{1}{p}} \leq W_i(K) \widetilde{W}_i(K^*). \tag{1.8}$$

Here  $\left[ \frac{\Omega_{p,i}(K)^{n+p-i}}{n^{n+p-i} W_i(K)^{n-p-i}} \right]^{\frac{1}{p}}$  may be called the  $i$ -th  $L_p$ -mixed affine surface area ratio of  $K$ .

THEOREM 1.1<sub>b</sub>. If  $K \in \mathcal{F}_o^n$ ,  $p \geq 1$  and  $i = 0, 1, \dots, n - 1$ , then

$$\left[ \frac{\Omega_{p,i}(K)^{n+p-i}}{n^{n+p-i} W_i(K)^{n-p-i}} \right]^{\frac{1}{p}} \leq W_i(K) \widetilde{W}_i(K^*), \tag{1.9}$$

with equality if and only if  $K^*$  and  $\Lambda_{p,i} K$  are dilates.

THEOREM 1.2<sub>a</sub>. If  $K \in \mathcal{K}_o^n$ ,  $1 \leq p < q$  and  $i = 0, 1, \dots, n - 1$ , then

$$\left[ \frac{\Omega_{p,i}(K)^{n+p-i}}{n^{n+p-i} W_i(K)^{n-p-i}} \right]^{\frac{1}{p}} \leq \left[ \frac{\Omega_{q,i}(K)^{n+q-i}}{n^{n+q-i} W_i(K)^{n-q-i}} \right]^{\frac{1}{q}}. \tag{1.10}$$

THEOREM 1.2<sub>b</sub>. If  $K \in \mathcal{F}_o^n$ ,  $1 \leq p < q$  and  $i = 0, 1, \dots, n - 1$ , then

$$\left[ \frac{\Omega_{p,i}(K)^{n+p-i}}{n^{n+p-i}W_i(K)^{n-p-i}} \right]^{\frac{1}{p}} \leq \left[ \frac{\Omega_{q,i}(K)^{n+q-i}}{n^{n+q-i}W_i(K)^{n-q-i}} \right]^{\frac{1}{q}}, \tag{1.11}$$

with equality if and only if  $\Lambda_{p,i}K$  and  $\Lambda_{q,i}K$  are dilates.

THEOREM 1.3<sub>a</sub>. If  $K \in \mathcal{K}_o^n$ ,  $1 \leq p < q < r$  and  $i = 0, 1, \dots, n - 1$ , then

$$\Omega_{q,i}(K)^{(n+q-i)(r-p)} \leq \Omega_{p,i}(K)^{(n+p-i)(r-q)}\Omega_{r,i}(K)^{(n+r-i)(q-p)}. \tag{1.12}$$

THEOREM 1.3<sub>b</sub>. If  $K \in \mathcal{F}_o^n$ ,  $1 \leq p < q < r$  and  $i = 0, 1, \dots, n - 1$ , then

$$\Omega_{q,i}(K)^{(n+q-i)(r-p)} \leq \Omega_{p,i}(K)^{(n+p-i)(r-q)}\Omega_{r,i}(K)^{(n+r-i)(q-p)}, \tag{1.13}$$

with equality if and only if  $\Lambda_{p,i}K$  and  $\Lambda_{r,i}K$  are dilates.

The proofs of Theorems 1.1<sub>a</sub>-1.3<sub>b</sub> will be completed in section 3 of this paper.

## 2. Notations and background materials

### 2.1. Support function, radial functions and polar set

If  $K \in \mathcal{K}^n$ , then its support function,  $h_K = h(K, \cdot) : \mathbb{R}^n \rightarrow (-\infty, \infty)$ , is defined by (see [4, 19])

$$h(K, x) = \max\{x \cdot y : y \in K\}, \quad x \in \mathbb{R}^n,$$

where  $x \cdot y$  denotes the standard inner product of  $x$  and  $y$ . Obviously,  $h(\lambda K, \cdot) = \lambda h(K, \cdot)$ , where  $\lambda$  is a positive constant.

If  $K$  is a compact star-shaped (with respect to the origin) in  $\mathbb{R}^n$ , then its radial function,  $\rho_K = \rho(K, \cdot) : \mathbb{R}^n \setminus \{0\} \rightarrow [0, \infty)$ , is defined by (see [4, 19])

$$\rho(K, x) = \max\{\lambda \geq 0 : \lambda x \in K\}, \quad x \in \mathbb{R}^n \setminus \{0\}.$$

If  $\rho_K$  is positive and continuous,  $K$  will be called a star body (with respect to the origin). Two star bodies  $K$  and  $L$  are said to be dilates (of one another) if  $\rho_K(u)/\rho_L(u)$  is independent of  $u \in S^{n-1}$ .

If  $E$  is a nonempty subset and contains the origin in  $\mathbb{R}^n$ , then the polar set,  $E^*$ , of  $E$  is defined by (see [4, 19])

$$E^* = \{x \in \mathbb{R}^n : x \cdot y \leq 1, y \in E\}.$$

It is easily verified that  $(K^*)^* = K$  for all  $K \in \mathcal{K}_o^n$ . Moreover, for  $K \in \mathcal{S}_o^n$  and all  $u \in S^{n-1}$ ,

$$\rho_K(u) = \frac{1}{h_{K^*}(u)}. \tag{2.1}$$

**2.2.  $L_p$ -mixed surface area measure and  $L_p$ -mixed curvature image**

The  $L_p$ -mixed surface area measure of convex bodies is introduced by Lutwak (see[14]). For  $K \in \mathcal{K}_o^n$ , real  $p \geq 1$  and  $i = 0, 1, \dots, n - 1$ , the  $L_p$ -mixed surface area measure,  $S_{p,i}(K, \cdot)$ , of  $K$  is defined by

$$\frac{dS_{p,i}(K, \cdot)}{dS_i(K, \cdot)} = h^{1-p}(K, \cdot). \tag{2.2}$$

Equation (2.2) is the Radon-Nikodym derivative of the  $L_p$ -surface area measure  $S_{p,i}(K, \cdot)$  with the respect to the surface area measure  $S_i(K, \cdot)$ .

For  $i = 0, 1, \dots, n - 1$ , we say  $K \in \mathcal{K}^n$  has a curvature function  $f_i(K, \cdot) : S^{n-1} \rightarrow \mathbb{R}$ , if measure  $S_i(K, \cdot)$  is absolutely continuous with respect to spherical Lebesgue measure  $S$ , and

$$\frac{dS_i(K, \cdot)}{dS} = f_i(K, \cdot). \tag{2.3}$$

Let  $p \geq 1$  and  $i = 0, 1, \dots, n - 1$ . A convex body  $K \in \mathcal{K}_o^n$  is said to have a generalized  $L_p$ -curvature function (see[18]),  $f_{p,i}(K, \cdot) : S^{n-1} \rightarrow \mathbb{R}$ , if measure  $S_{p,i}(K, \cdot)$  is absolutely continuous with respect to spherical Lebesgue measure  $S$ , and

$$\frac{dS_{p,i}(K, \cdot)}{dS} = f_{p,i}(K, \cdot). \tag{2.4}$$

Obviously,  $f_{p,0}(K, \cdot) = f_p(K, \cdot)$ . Here  $f_p(K, \cdot)$  is the  $L_p$ -curvature function of  $K \in \mathcal{K}_o^n$  (see [15]).

Also, from (2.2), (2.3) and (2.4), we know that for  $K \in \mathcal{F}_o^n$ ,

$$f_{p,i}(K, \cdot) = h^{1-p}(K, \cdot)f_i(K, \cdot). \tag{2.5}$$

Meanwhile, according to the definition of  $L_p$ -curvature image, Lu and Wang ([18]) gave the definition of  $L_p$ -mixed curvature image as follows: For  $K \in \mathcal{F}_o^n$ ,  $p \geq 1$  and  $i = 0, 1, \dots, n - 1$ , the  $L_p$ -mixed curvature image,  $\Lambda_{p,i}K \in \mathcal{S}_o^n$ , of  $K$  is defined by

$$\rho(\Lambda_{p,i}K, \cdot)^{n+p-i} = \frac{\widetilde{W}_i(\Lambda_{p,i}K)}{\omega_n} f_{p,i}(K, \cdot). \tag{2.6}$$

If  $i = 0$  in (2.6), then

$$\Lambda_{p,0}K = \Lambda_pK.$$

Here  $\Lambda_pK$  is called  $L_p$ -curvature image that was established by Lutwak (see[15]).

**2.3. Quermassintegrals and  $L_p$ -mixed quermassintegrals**

For  $K \in \mathcal{K}^n$ ,  $i = 0, 1, \dots, n - 1$ , the quermassintegrals,  $W_i(K)$ , of  $K$  are defined by (see [4, 19])

$$W_i(K) = \frac{1}{n} \int_{S^{n-1}} h_K(u) dS_i(K, u).$$

Here  $S_i(K, \cdot)$  ( $i = 0, 1, \dots, n - 1$ ) are the area measure of  $K \in \mathcal{K}^n$ . Obviously,

$$W_0(K) = \frac{1}{n} \int_{S^{n-1}} h_K(u) dS(K, u) = V(K).$$

For  $K, L \in \mathcal{K}_o^n$ ,  $p \geq 1$  and  $\lambda, \mu \geq 0$  (not both zero), the  $L_p$ -Minkowski linear combination (also called Firey combination),  $\lambda \cdot K +_p \mu \cdot L \in \mathcal{K}_o^n$ , of  $K$  and  $L$  is defined by (see[14])

$$h(\lambda \cdot K +_p \mu \cdot L, \cdot)^p = \lambda h(K, \cdot)^p + \mu h(L, \cdot)^p.$$

Here ‘ $+_p$ ’ denotes the  $L_p$ -Minkowski addition and ‘ $\cdot$ ’ denotes the Firey scalar multiplication.

Associated with  $L_p$ -mixed surface area measure, Lutwak ([14]) defined the  $L_p$ -mixed quermassintegrals (also called mixed  $p$ -quermassintegrals). For  $K, L \in \mathcal{K}_o^n$ ,  $p \geq 1$ ,  $i = 0, 1, \dots, n - 1$ , the  $L_p$ -mixed quermassintegrals,  $W_{p,i}(K, L)$ , of  $K$  and  $L$  are given by (see [14])

$$W_{p,i}(K, L) = \frac{1}{n} \int_{S^{n-1}} h_L^p(u) dS_{p,i}(K, u). \tag{2.7}$$

From (2.7), it follows immediately that for each  $K \in \mathcal{K}_o^n$  and all  $p \geq 1$ ,

$$W_{p,i}(K, K) = W_i(K). \tag{2.8}$$

Let  $i = 0$  in (2.7), the  $L_p$ -mixed volume,  $V_p(K, L)$ , of  $K, L \in \mathcal{K}_o^n$  is given by

$$V_p(K, L) = \frac{1}{n} \int_{S^{n-1}} h_L^p(u) dS_p(K, u).$$

If  $i = 0$  and  $p = 1$  in (2.7), then the mixed volume,  $V_1(K, L)$ , of convex bodies  $K$  and  $L$  is defined by

$$V_1(K, L) = \frac{1}{n} \int_{S^{n-1}} h_L(u) dS(K, u).$$

### 2.4. Dual quermassintegrals and $L_p$ -dual mixed quermassintegrals

For  $K \in \mathcal{S}_o^n$  and real  $i$ , the dual quermassintegrals,  $\tilde{W}_i(K)$ , of  $K$  is defined by (see[9])

$$\tilde{W}_i(K) = \frac{1}{n} \int_{S^{n-1}} \rho_K^{n-i}(u) dS(u). \tag{2.9}$$

Obviously, for  $i = 0$ ,

$$\tilde{W}_0(K) = \frac{1}{n} \int_{S^{n-1}} \rho_K^n(u) dS(u) = V(K).$$

For  $K, L \in \mathcal{S}_o^n$ ,  $p \geq 1$  and  $\lambda, \mu \geq 0$  (not both zero), the  $L_p$ -harmonic radial combination,  $\lambda \times K \tilde{+}_{-p} \mu \times L \in \mathcal{S}_o^n$ , of  $K$  and  $L$  is defined by (see[15])

$$\rho(\lambda \times K \tilde{+}_{-p} \mu \times L, \cdot)^{-p} = \lambda \rho(K, \cdot)^{-p} + \mu \rho(L, \cdot)^{-p}.$$

Associated with the  $L_p$ -harmonic radial combination of star bodies, Wang and Leng (see[23]) introduced the notion of  $L_p$ -dual mixed quermassintegrals as follows: For  $K, L \in \mathcal{S}_o^n$ ,  $p \geq 1$  and real  $i \neq n$ , the  $L_p$ -dual mixed quermassintegrals,  $\tilde{W}_{-p,i}(K, L)$ , of  $K$  and  $L$  is defined by

$$\tilde{W}_{-p,i}(K, L) = \frac{1}{n} \int_{S^{n-1}} \rho_K^{n+p-i}(u) \rho_L^{-p}(u) dS(u). \tag{2.10}$$

From (2.9) and (2.10), it follows immediately that for each  $K \in \mathcal{S}_o^n$  and all  $p \geq 1$ ,

$$\tilde{W}_{-p,i}(K, K) = \tilde{W}_i(K). \tag{2.11}$$

The Minkowski’s inequality for the  $L_p$ -dual mixed quermassintegrals is (see[23]):

Let  $K, L \in \mathcal{S}_o^n$ ,  $p \geq 1$  and real  $i \neq n$ , then for  $i < n$  or  $n < i < n + p$ ,

$$\tilde{W}_{-p,i}(K, L) \geq \tilde{W}_i(K)^{\frac{n+p-i}{n-i}} \tilde{W}_i(L)^{-\frac{p}{n-i}}, \tag{2.12}$$

for  $i > n + p$ , the inequality (2.12) is reversed. Equality holds in each inequality if and only if  $K$  and  $L$  are dilates.

### 3. Proofs of Theorems

In this section, we complete the proofs of Theorems 1.1<sub>a</sub>-1.3<sub>b</sub>. Here, we first give a property of the  $L_p$ -mixed affine surface area  $\Omega_{p,i}(K)$  as follows:

**THEOREM 3.1.** *If  $K \in \mathcal{F}_o^n$ ,  $p \geq 1$  and  $i = 0, 1, \dots, n - 1$ , then*

$$\Omega_{p,i}(K)^{n+p-i} = n^{n+p-i} \omega_n^{n-i} \tilde{W}_i(\Lambda_{p,i}K)^p. \tag{3.1}$$

**LEMMA 3.1.** [18] *If  $K \in \mathcal{F}_o^n$ ,  $p \geq 1$  and  $i = 0, 1, \dots, n - 1$ , then for any  $Q \in \mathcal{S}_o^n$ ,*

$$\tilde{W}_{-p,i}(\Lambda_{p,i}K, Q) = \frac{\tilde{W}_i(\Lambda_{p,i}K)}{\omega_n} W_{p,i}(K, Q^*). \tag{3.2}$$

*Proof of Theorem 3.1.* According to (2.12), for  $i < n$ , we have for any  $Q \in \mathcal{S}_o^n$

$$\tilde{W}_{-p,i}(\Lambda_{p,i}K, Q) \geq \tilde{W}_i(\Lambda_{p,i}K)^{\frac{n+p-i}{n-i}} \tilde{W}_i(Q)^{-\frac{p}{n-i}}. \tag{3.3}$$

Using (3.2), (3.3) and definition (1.7), yield

$$\begin{aligned} n^{-\frac{p}{n-i}} \Omega_{p,i}(K)^{\frac{n+p-i}{n-i}} &= \inf \{ n W_{p,i}(K, Q^*) \tilde{W}_i(Q)^{\frac{p}{n-i}} : Q \in \mathcal{S}_o^n \} \\ &= \inf \left\{ \frac{n \omega_n}{\tilde{W}_i(\Lambda_{p,i}K)} \tilde{W}_{-p,i}(\Lambda_{p,i}K, Q) \tilde{W}_i(Q)^{\frac{p}{n-i}} : Q \in \mathcal{S}_o^n \right\} \\ &\geq \inf \left\{ \frac{n \omega_n}{\tilde{W}_i(\Lambda_{p,i}K)} \tilde{W}_i(\Lambda_{p,i}K)^{\frac{n+p-i}{n-i}} \tilde{W}_i(Q)^{-\frac{p}{n-i}} \tilde{W}_i(Q)^{\frac{p}{n-i}} : Q \in \mathcal{S}_o^n \right\} \\ &= \inf \left\{ n \omega_n \tilde{W}_i(\Lambda_{p,i}K)^{\frac{p}{n-i}} : Q \in \mathcal{S}_o^n \right\} \\ &= n \omega_n \tilde{W}_i(\Lambda_{p,i}K)^{\frac{p}{n-i}}. \end{aligned} \tag{3.4}$$

On the other hand, combining with definition (1.7) and (3.2), we obtain for any  $Q \in \mathcal{S}_o^n$

$$\begin{aligned} n^{-\frac{p}{n-i}} \Omega_{p,i}(K)^{\frac{n+p-i}{n-i}} &\leq n W_{p,i}(K, Q^*) \widetilde{W}_i(Q)^{\frac{p}{n-i}} \\ &= \frac{n \omega_n}{\widetilde{W}_i(\Lambda_{p,i}K)} \widetilde{W}_{-p,i}(\Lambda_{p,i}K, Q) \widetilde{W}_i(Q)^{\frac{p}{n-i}}. \end{aligned} \tag{3.5}$$

Taking  $Q$  for  $\Lambda_{p,i}K$  in (3.5) and using (2.11), then

$$n^{-\frac{p}{n-i}} \Omega_{p,i}(K)^{\frac{n+p-i}{n-i}} \leq n \omega_n \widetilde{W}_i(\Lambda_{p,i}K)^{\frac{p}{n-i}}. \tag{3.6}$$

From (3.4) and (3.6), we see that

$$n^{-\frac{p}{n-i}} \Omega_{p,i}(K)^{\frac{n+p-i}{n-i}} = n \omega_n \widetilde{W}_i(\Lambda_{p,i}K)^{\frac{p}{n-i}}.$$

i.e.

$$\Omega_{p,i}(K)^{n+p-i} = n^{n+p-i} \omega_n^{n-i} \widetilde{W}_i(\Lambda_{p,i}K)^p.$$

This yields (3.1).  $\square$

Let  $i = 0$  in Theorem 3.1 to get the following result which was obtained by Lutwak (see [15]).

**COROLLARY 3.1.** *Suppose  $K \in \mathcal{F}_o^n$ ,  $p \geq 1$ , then*

$$\Omega_p(K) = n \omega_n^{\frac{n}{n+p}} V(\Lambda_p K)^{\frac{p}{n+p}}.$$

*Proof of Theorem 1.1a.* From definition (1.7), we obtain for any  $Q \in \mathcal{S}_o^n$ ,

$$\Omega_{p,i}(K)^{\frac{n+p-i}{n-i}} \leq n^{\frac{n+p-i}{n-i}} W_{p,i}(K, Q^*) \widetilde{W}_i(Q)^{\frac{p}{n-i}}, \tag{3.7}$$

Since  $Q^* \in \mathcal{K}_o^n$ , thus taking  $Q^* = K$  in (3.7), then

$$\Omega_{p,i}(K)^{\frac{n+p-i}{n-i}} \leq n^{\frac{n+p-i}{n-i}} W_i(K) \widetilde{W}_i(K^*)^{\frac{p}{n-i}},$$

Hence

$$\left[ \frac{\Omega_{p,i}(K)^{n+p-i}}{n^{n+p-i} W_i(K)^{n-p-i}} \right]^{\frac{1}{p}} \leq W_i(K) \widetilde{W}_i(K^*).$$

This gives (1.8).  $\square$

*Proof of Theorem 1.1b.* Let  $Q^* = K$  in (3.2), and together (2.8) and inequality (2.12), we have that for all  $i < n$ ,

$$\begin{aligned} W_i(K) &= \frac{\omega_n}{\widetilde{W}_i(\Lambda_{p,i}K)} \widetilde{W}_{-p,i}(\Lambda_{p,i}K, K^*) \\ &\geq \frac{\omega_n}{\widetilde{W}_i(\Lambda_{p,i}K)} \widetilde{W}_i(\Lambda_{p,i}K)^{\frac{n+p-i}{n-i}} \widetilde{W}_i(K^*)^{-\frac{p}{n-i}}. \end{aligned}$$



Then

$$\left[ \frac{\omega_n^{n-i} \widetilde{W}_i(\Lambda_{p,i}K)^p}{W_i(K)^{n-p-i}} \right]^{\frac{1}{p}} \leq W_i(K) \widetilde{W}_i(K^*).$$

Using (3.1), we get

$$\left[ \frac{\Omega_{p,i}(K)^{n+p-i}}{n^{n+p-i} W_i(K)^{n-p-i}} \right]^{\frac{1}{p}} \leq W_i(K) \widetilde{W}_i(K^*). \tag{3.8}$$

According to the equality condition of (2.12), we see that equality holds in (3.8) if and only if  $K^*$  and  $\Lambda_{p,i}K$  are dilates. This yields (1.9).  $\square$

In order to prove Theorem 1.2<sub>a</sub>, the following lemma obtained by Wei and Wang (see[29]) is needed.

LEMMA 3.2. *If  $K, L \in \mathcal{K}_o^n$ ,  $1 \leq p < q$  and  $i = 0, 1, \dots, n-1$ , then*

$$\left[ \frac{W_{p,i}(K, L)}{W_i(K)} \right]^{\frac{1}{p}} \leq \left[ \frac{W_{q,i}(K, L)}{W_i(K)} \right]^{\frac{1}{q}},$$

with equality if and only if  $K$  and  $L$  are dilates.

*Proof of Theorem 1.2<sub>a</sub>.* For any  $Q \in \mathcal{S}_o^n$ , we have  $Q^* \in \mathcal{K}_o^n$ . Hence, by Lemma 3.2, we get for  $1 \leq p < q$ ,

$$\left[ \frac{W_{p,i}(K, Q^*)}{W_i(K)} \right]^{\frac{1}{p}} \leq \left[ \frac{W_{q,i}(K, Q^*)}{W_i(K)} \right]^{\frac{1}{q}}, \tag{3.9}$$

with equality if and only if  $K$  is a dilate of  $Q^*$ .

Combining with definition (1.7) and (3.9), if  $i = 0, 1, \dots, n-1$ , then we yield

$$\begin{aligned} \left[ \frac{\Omega_{p,i}(K)^{n+p-i}}{n^{n+p-i} W_i(K)^{n-p-i}} \right]^{\frac{1}{p}} &= \left[ \frac{\Omega_{p,i}(K)^{\frac{n+p-i}{n-i}}}{n^{\frac{n+p-i}{n-i}} W_i(K)^{\frac{n-p-i}{n-i}}} \right]^{\frac{n-i}{p}} \\ &= \inf \left\{ \left[ \frac{W_{p,i}(K, Q^*)}{W_i(K)} \right]^{\frac{n-i}{p}} W_i(K) \widetilde{W}_i(Q) : Q \in \mathcal{S}_o^n \right\} \\ &\leq \inf \left\{ \left[ \frac{W_{q,i}(K, Q^*)}{W_i(K)} \right]^{\frac{n-i}{q}} W_i(K) \widetilde{W}_i(Q) : Q \in \mathcal{S}_o^n \right\} \\ &= \left[ \frac{\Omega_{q,i}(K)^{n+q-i}}{n^{n+q-i} W_i(K)^{n-q-i}} \right]^{\frac{1}{q}}. \end{aligned}$$

Hence

$$\left[ \frac{\Omega_{p,i}(K)^{n+p-i}}{n^{n+p-i} W_i(K)^{n-p-i}} \right]^{\frac{1}{p}} \leq \left[ \frac{\Omega_{q,i}(K)^{n+q-i}}{n^{n+q-i} W_i(K)^{n-q-i}} \right]^{\frac{1}{q}}.$$

This yields (1.10).  $\square$

*Proof of Theorem 1.2 b.* From (3.2), we have

$$\left[ \frac{W_{p,i}(K, Q^*)}{W_i(K)} \right]^{\frac{1}{p}} = \left[ \frac{\omega_n \widetilde{W}_{-p,i}(\Lambda_{p,i}K, Q)}{W_i(K) \widetilde{W}_i(\Lambda_{p,i}K)} \right]^{\frac{1}{p}},$$

$$\left[ \frac{W_{q,i}(K, Q^*)}{W_i(K)} \right]^{\frac{1}{q}} = \left[ \frac{\omega_n \widetilde{W}_{-q,i}(\Lambda_{q,i}K, Q)}{W_i(K) \widetilde{W}_i(\Lambda_{q,i}K)} \right]^{\frac{1}{q}}.$$

Using (3.9), we get that for  $1 \leq p < q$ ,

$$\left[ \frac{\omega_n \widetilde{W}_{-p,i}(\Lambda_{p,i}K, Q)}{W_i(K) \widetilde{W}_i(\Lambda_{p,i}K)} \right]^{\frac{1}{p}} \leq \left[ \frac{\omega_n \widetilde{W}_{-q,i}(\Lambda_{q,i}K, Q)}{W_i(K) \widetilde{W}_i(\Lambda_{q,i}K)} \right]^{\frac{1}{q}}. \tag{3.10}$$

Taking  $Q = \Lambda_{q,i}K$  in (3.10), and using (2.11) and inequality (2.12), we obtain

$$\begin{aligned} \left[ \frac{\omega_n}{W_i(K)} \right]^{\frac{1}{q}} &\geq \left[ \frac{\omega_n \widetilde{W}_{-p,i}(\Lambda_{p,i}K, \Lambda_{q,i}K)}{W_i(K) \widetilde{W}_i(\Lambda_{p,i}K)} \right]^{\frac{1}{p}} \\ &\geq \left[ \frac{\omega_n \widetilde{W}_i(\Lambda_{p,i}K)^{\frac{p}{n-i}} \widetilde{W}_i(\Lambda_{q,i}K)^{-\frac{p}{n-i}}}{W_i(K)} \right]^{\frac{1}{p}}. \end{aligned}$$

So, we get

$$\left[ \frac{\omega_n \widetilde{W}_i(\Lambda_{p,i}K)^{\frac{p}{n-i}}}{W_i(K)} \right]^{\frac{1}{p}} \leq \left[ \frac{\omega_n \widetilde{W}_i(\Lambda_{q,i}K)^{\frac{q}{n-i}}}{W_i(K)} \right]^{\frac{1}{q}}.$$

From (3.1), we have

$$\left[ \frac{\Omega_{p,i}(K)^{n+p-i}}{n^{n+p-i} W_i(K)^{n-p-i}} \right]^{\frac{1}{p}} \leq \left[ \frac{\Omega_{q,i}(K)^{n+q-i}}{n^{n+q-i} W_i(K)^{n-q-i}} \right]^{\frac{1}{q}}. \tag{3.11}$$

According to the equality condition of (2.12), we see that equality holds in (3.11) if and only if  $\Lambda_{p,i}K$  and  $\Lambda_{q,i}K$  are dilates. This yields (1.11).  $\square$

*Proof of Theorem 1.3 a.* Since for any  $Q_1, Q_2 \in \mathcal{S}_o^n$ , there exists  $Q_3 \in \mathcal{S}_o^n$  such that

$$\rho(Q_3, \cdot)^{q(r-p)} = \rho(Q_1, \cdot)^{p(r-q)} \rho(Q_2, \cdot)^{r(q-p)}. \tag{3.12}$$

Then for any  $u \in S^{n-1}$ , this yields

$$\rho_{Q_3}(u)^{n-i} = \rho_{Q_1}(u)^{\frac{(n-i)p(r-q)}{q(r-p)}} \rho_{Q_2}(u)^{\frac{(n-i)r(q-p)}{q(r-p)}}.$$

Since  $1 \leq p < q < r$ , then  $\frac{q(r-p)}{p(r-q)} > 1$ . According to the Hölder's integral inequality

(see[5]) and definition (2.9), we get

$$\begin{aligned}
 & \widetilde{W}_i(Q_1)^{\frac{p(r-q)}{q(r-p)}} \widetilde{W}_i(Q_2)^{\frac{r(q-p)}{q(r-p)}} \\
 &= \left[ \frac{1}{n} \int_{S^{n-1}} \rho_{Q_1}(u)^{n-i} dS(u) \right]^{\frac{p(r-q)}{q(r-p)}} \left[ \frac{1}{n} \int_{S^{n-1}} \rho_{Q_2}(u)^{n-i} dS(u) \right]^{\frac{r(q-p)}{q(r-p)}} \\
 &= \left\{ \frac{1}{n} \int_{S^{n-1}} \left[ \rho_{Q_1}(u)^{\frac{(n-i)p(r-q)}{q(r-p)}} \right]^{\frac{q(r-p)}{p(r-q)}} dS(u) \right\}^{\frac{p(r-q)}{q(r-p)}} \\
 &\quad \times \left\{ \frac{1}{n} \int_{S^{n-1}} \left[ \rho_{Q_2}(u)^{\frac{(n-i)r(q-p)}{q(r-p)}} \right]^{\frac{q(r-p)}{r(q-p)}} dS(u) \right\}^{\frac{r(q-p)}{q(r-p)}} \\
 &\geq \frac{1}{n} \int_{S^{n-1}} \rho_{Q_1}(u)^{\frac{(n-i)p(r-q)}{q(r-p)}} \rho_{Q_2}(u)^{\frac{(n-i)r(q-p)}{q(r-p)}} dS(u) \\
 &= \frac{1}{n} \int_{S^{n-1}} \rho_{Q_3}(u)^{n-i} dS(u) = \widetilde{W}_i(Q_3).
 \end{aligned}$$

Since  $q(r-p) > 0$ , then

$$\widetilde{W}_i(Q_3)^{q(r-p)} \leq \widetilde{W}_i(Q_1)^{p(r-q)} \widetilde{W}_i(Q_2)^{r(q-p)}. \tag{3.13}$$

From (3.12), we see that for any  $u \in S^{n-1}$ ,

$$\rho_{Q_3}(u)^{-q} h_K(u)^{1-q} = [\rho_{Q_1}(u)^{-p} h_K(u)^{1-p}]^{\frac{r-q}{r-p}} [\rho_{Q_2}(u)^{-r} h_K(u)^{1-r}]^{\frac{q-p}{r-p}}. \tag{3.14}$$

Then for  $1 \leq p < q < r$ , i.e.  $\frac{r-p}{r-q} > 1$ , according to the Hölder’s integral inequality, (2.1), (2.7) and (3.14), we get

$$\begin{aligned}
 & W_{p,i}(K, Q_1^*)^{\frac{r-q}{r-p}} W_{r,i}(K, Q_2^*)^{\frac{q-p}{r-p}} \\
 &= \left[ \frac{1}{n} \int_{S^{n-1}} h_{Q_1^*}(u)^p h_K(u)^{1-p} dS_i(K, u) \right]^{\frac{r-q}{r-p}} \\
 &\quad \times \left[ \frac{1}{n} \int_{S^{n-1}} h_{Q_2^*}(u)^r h_K(u)^{1-r} dS_i(K, u) \right]^{\frac{q-p}{r-p}} \\
 &= \left[ \frac{1}{n} \int_{S^{n-1}} [\rho_{Q_1}(u)^{-p} h_K(u)^{1-p}]^{\frac{r-q}{r-p}} [\rho_{Q_2}(u)^{-r} h_K(u)^{1-r}]^{\frac{q-p}{r-p}} dS_i(K, u) \right]^{\frac{r-q}{r-p}} \\
 &\quad \times \left[ \frac{1}{n} \int_{S^{n-1}} [\rho_{Q_2}(u)^{-r} h_K(u)^{1-r}]^{\frac{q-p}{r-p}} [\rho_{Q_1}(u)^{-p} h_K(u)^{1-p}]^{\frac{r-q}{r-p}} dS_i(K, u) \right]^{\frac{q-p}{r-p}} \\
 &\geq \frac{1}{n} \int_{S^{n-1}} \rho_{Q_3}(u)^{-q} h_K(u)^{1-q} dS_i(K, u) \\
 &= W_{q,i}(K, Q_3^*),
 \end{aligned}$$

i.e.

$$W_{q,i}(K, Q_3^*)^{r-p} \leq W_{p,i}(K, Q_1^*)^{r-q} W_{r,i}(K, Q_2^*)^{q-p}. \tag{3.15}$$

Hence, combining with (3.13) and (3.15), we get for  $i = 0, 1, \dots, n - 1$ ,

$$\begin{aligned} & [W_{q,i}(K, Q_3^*) \widetilde{W}_i(Q_3)^{\frac{q}{n-i}}]^{r-p} \\ & \leq [W_{p,i}(K, Q_1^*) \widetilde{W}_i(Q_1)^{\frac{p}{n-i}}]^{r-q} [W_{r,i}(K, Q_2^*) \widetilde{W}_i(Q_2)^{\frac{r}{n-i}}]^{q-p}. \end{aligned}$$

This together with (1.7) yields

$$\Omega_{q,i}(K)^{(n+q-i)(r-p)} \leq \Omega_{p,i}(K)^{(n+p-i)(r-q)} \Omega_{r,i}(K)^{(n+r-i)(q-p)}.$$

This gives (1.12).  $\square$

Finally, we give the proof of Theorem 1.3<sub>b</sub>. The following lemma is required.

LEMMA 3.3. *If  $K \in \mathcal{F}_o^n$ ,  $1 \leq p < q < r$  and  $i = 0, 1, \dots, n - 1$ , then*

$$\widetilde{W}_i(\Lambda_{q,i}K)^{q(r-p)} \leq \widetilde{W}_i(\Lambda_{p,i}K)^{p(r-q)} \widetilde{W}_i(\Lambda_{r,i}K)^{r(q-p)}, \tag{3.16}$$

with equality if and only if  $\Lambda_{p,i}K$  and  $\Lambda_{r,i}K$  are dilates.

*Proof.* From the formula (2.5), it follows that for  $i = 0, 1, \dots, n - 1$ ,

$$f_{q,i}(K, \cdot)^{r-p} = f_{p,i}(K, \cdot)^{r-q} f_{r,i}(K, \cdot)^{q-p}.$$

Thus, by (2.6), we get for any  $u \in S^{n-1}$ ,

$$\begin{aligned} & \widetilde{W}_i(\Lambda_{q,i}K)^{p-r} \rho(\Lambda_{q,i}K, u)^{(n+q-i)(r-p)} \\ & = [\widetilde{W}_i(\Lambda_{p,i}K)^{q-r} \rho(\Lambda_{p,i}K, u)^{(n+p-i)(r-q)}] [\widetilde{W}_i(\Lambda_{r,i}K)^{p-q} \rho(\Lambda_{r,i}K, u)^{(n+r-i)(q-p)}], \end{aligned}$$

that is

$$\begin{aligned} & \widetilde{W}_i(\Lambda_{q,i}K)^{\frac{(p-r)(n-i)}{(n+q-i)(r-p)}} \rho(\Lambda_{q,i}K, u)^{n-i} \\ & = [\widetilde{W}_i(\Lambda_{p,i}K)^{\frac{(q-r)(n-i)}{(n+p-i)(r-p)}} (\rho(\Lambda_{p,i}K, u)^{n-i})^{\frac{(n+p-i)(r-q)}{(n+q-i)(r-p)}}] \\ & \quad \times [\widetilde{W}_i(\Lambda_{r,i}K)^{\frac{(p-q)(n-i)}{(n+r-i)(q-p)}} (\rho(\Lambda_{r,i}K, u)^{n-i})^{\frac{(n+r-i)(q-p)}{(n+q-i)(r-p)}}]. \end{aligned} \tag{3.17}$$

Using the Hölder inequality and (2.9) in (3.17), we obtain

$$\widetilde{W}_i(\Lambda_{q,i}K)^{q(r-p)} \leq \widetilde{W}_i(\Lambda_{p,i}K)^{p(r-q)} \widetilde{W}_i(\Lambda_{r,i}K)^{r(q-p)}. \tag{3.18}$$

From the equality condition of the Hölder inequality, we see that equality holds in (3.18) if and only if  $\Lambda_{p,i}K$  and  $\Lambda_{r,i}K$  are dilates. This yields (3.16).  $\square$

*Proof of Theorem 1.3<sub>b</sub>.* From (3.1) and (3.16), we have

$$\begin{aligned} \Omega_{q,i}(K)^{(n+q-i)(r-p)} & = [n^{n+q-i} \omega_n^{n-i}]^{r-p} \widetilde{W}_i(\Lambda_{q,i}K)^{q(r-p)} \\ & \leq [n^{n+q-i} \omega_n^{n-i}]^{r-p} \widetilde{W}_i(\Lambda_{p,i}K)^{p(r-q)} \widetilde{W}_i(\Lambda_{r,i}K)^{r(q-p)} \\ & = \Omega_{p,i}(K)^{(n+p-i)(r-q)} \Omega_{r,i}(K)^{(n+r-i)(q-p)}. \end{aligned} \tag{3.19}$$

According to the equality condition of (3.16), we know that equality holds in (3.19) if and only if  $\Lambda_{p,i}K$  and  $\Lambda_{r,i}K$  are dilates. This gives (1.13).  $\square$

*Acknowledgements.* The authors would like to sincerely thank the referees for very valuable and helpful comments and suggestions, which made the paper more accurate and readable.

## REFERENCES

- [1] W. BLASCHKE, *Über affine geometrie XXIX: affininimalflächen*, Math. Z., **12**, (1922), 262–273.
- [2] W. S. CHEUNG AND C. ZHAO, *Width-integrals and affine surface area of convex bodies*, Banach J. Math. Anal., **2**, 1 (2008), 70–77.
- [3] Y. FENG AND W. WANG, *Blaschke-Minkowski homomorphisms and affine surface area*, Publ. Math. Debrecen, **85**, 3–4 (2014), 297–308.
- [4] R. J. GARDNER, *Geometric Tomography*, Cambridge Univ. Press, Cambridge, UK, 2nd edition, 2006.
- [5] G. H. HARDY, J. E. LITTLEWOOD AND G. PÓLYA, *Inequalities*, Cambridge Univ. Press, Cambridge, 1952.
- [6] K. LEICHTWEISS, *Über eine formel Blaschkes zur affinoberfläche*, Studia Sci. Math. Hungar., (1986), 453–474.
- [7] K. LEICHTWEISS, *Bemerkungen zur Definition einer erweiterten affinoberfläche von E. Lutwak*, Manuscripta. Math., **65** (1989), 181–197.
- [8] K. LEICHTWEISS, *On the history of the affine surface area for convex bodies*, Results Math., **20** (1991), 650–656.
- [9] E. LUTWAK, *Dual mixed volumes*, Pacific J. Math., **58**, (1975), 531–538.
- [10] E. LUTWAK, *On the Blaschke-Santaló inequality*, Ann. New York Acad. Sci., **440** (1985), 106–112.
- [11] E. LUTWAK, *On some affine isoperimetric inequalities*, J. Differential Geom., **56** (1986), 1–13.
- [12] E. LUTWAK, *Mixed affine surface area*, J. Math. Anal. Appl., **125** (1987), 351–360.
- [13] E. LUTWAK, *Extend affine surface area*, Adv. Math., **85** (1991), 39–68.
- [14] E. LUTWAK, *The Brunn-Minkowski-Firey theory I: mixed volumes and the Minkowski problem*, J. Differential Geom., **38** (1993), 131–150.
- [15] E. LUTWAK, *The Brunn-Minkowski-Firey theory II: affine and geominimal surface areas*, Adv. Math., **118**, 2 (1996), 244–294.
- [16] M. LUDWIG AND M. REITZNER, *A characterization of affine surface area*, Adv. Math., **147** (1999), 138–172.
- [17] F. H. LU, W. D. WANG AND G. S. LENG, *On inequalities for mixed affine surface area*, Indian J. pure appl. Math., **38**, 3 (2007), 153–161.
- [18] F. H. LU AND W. D. WANG, *Inequalities for  $L_p$ -mixed curvature image*, Acta Mathematica Scientia., **30B**, 4 (2010), 1044–1052.
- [19] R. SCHNEIDER, *Convex Bodies: The Brunn-Minkowski theory*, Cambridge University Press, Cambridge, 2nd edition, 2014.
- [20] R. SCHNEIDER, *Affine surface area and convex bodies of elliptic type*, Periodica Mathematica Hungarica, **69**, 2 (2014), 120–125.
- [21] C. SCHÜTT AND E. WERNER, *Surface bodies and  $p$ -affine surface area*, Adv. Math., **187** (2004), 98–145.
- [22] W. D. WANG AND Y. B. FENG, *A general  $L_p$ -version of Prtty's affine projection inequality*, Taiwan J. Math., **17**, 2 (2013), 517–528.
- [23] W. D. WANG AND G. S. LENG,  *$L_p$ -dual mixed quermassintegrals*, Indian J. Pure Appl. Math., **36**, 4 (2005), 177–188.
- [24] W. D. WANG AND G. S. LENG,  *$L_p$ -mixed affine surface area*, J. Math. Anal. Appl., **335** (2007), 341–354.
- [25] W. D. WANG AND G. S. LENG, *Some affine isoperimetric inequalities associated with  $L_p$ -affine surface area*, Houston J. Math., **34**, 2 (2008), 443–453.
- [26] E. WERNER, *On  $L_p$  affine surface areas*, Indiana Univ. Math. J., **56** (2007), 2305–2324.

- [27] E. WERNER, *Rényi divergence and  $L_p$ -affine surface area for convex bodies*, Adv. Math., **230** (2012), 1040–1059.
- [28] M. MEYER AND E. WERNER, *On the  $p$ -affine surface area*, Adv. Math., **152**, 2 (2000), 288–313.
- [29] B. WEI AND W. D. WANG, *Some inequalities for the  $L_p$ -mixed quermassintegrals*, Wuhan University Journal of Natural Sciences, **18**, 3 (2013), 233–236.
- [30] E. WERNER AND D. P. YE, *New  $L_p$ -affine isoperimetric inequalities*, Adv. Math., **218** (2008), 762–780.
- [31] E. WERNER AND D. P. YE,  *$L_p$  Inequalities for mixed  $p$ -affine surface area*, Math. Ann., **347** (2010), 703–737.
- [32] D. P. YE, *Inequalities for general mixed affine surface areas*, J. London Math. Soc., **85** (2012), 101–120.
- [33] D. P. YE, B. C. ZHU AND J. Z. ZHOU, *The mixed  $L_p$  geominimal surface areas for multiple convex bodies*, arXiv: 1311.5180v1.
- [34] C. J. ZHAO AND G. S. LENG, *Width-integrals of projection bodies and affine surface area*, Chinese Annals of Mathematics Series A, **2**, 2 (2005), 275–282.
- [35] C. J. ZHAO AND B. MIHÁLY, *Width-integrals of mixed projection bodies and mixed affine surface area*, General Mathematics, **19**, 1 (2011), 123–133.
- [36] C. J. ZHAO, *On quotients of  $i$ th affine surface areas*, Turk. J. Math., **37**, 6 (2013), 1022–1029.

(Received July 18, 2016)

Tian Li

Department of Mathematics  
China Three Gorges University  
Yichang, 443002, P. R. China  
e-mail: 13997717035@163.com

Weidong Wang

Department of Mathematics  
China Three Gorges University  
Yichang, 443002, P. R. China  
e-mail: wdwxh722@163.com