

## ESTIMATES FOR TSALLIS RELATIVE OPERATOR ENTROPY

HAMID REZA MORADI, SHIGERU FURUICHI AND NICUȘOR MINCULETE

(Communicated by J.-C. Bourin)

*Abstract.* We give the tight bounds of Tsallis relative operator entropy by using Hermite-Hadamard's inequality. Some reverse inequalities related to Young's inequality are also given. In addition, operator inequalities for normalized positive linear map with Tsallis relative operator entropy are given.

### 1. Introduction and preliminaries

The operator theory related to inequalities in Hilbert space is studied in many papers. Let  $A, B$  be two operators in a Hilbert space  $\mathcal{H}$ . An operator  $A$  is said to be strictly positive (denoted by  $A > 0$ ) if  $A$  is positive and invertible. For two strictly positive operators  $A, B$  and  $p \in [0, 1]$ ,  $p$ -power mean  $A\#_p B$  is defined by

$$A\#_p B := A^{\frac{1}{2}} \left( A^{-\frac{1}{2}} B A^{-\frac{1}{2}} \right)^p A^{\frac{1}{2}},$$

and we remark that  $A\#_p B = A^{1-p} B^p$  if  $A$  commutes with  $B$ . We also use the symbol  $A\sharp_r B := A^{\frac{1}{2}} \left( A^{-\frac{1}{2}} B A^{-\frac{1}{2}} \right)^r A^{\frac{1}{2}}$  for  $r \in \mathbb{R}$ . The weighted operator arithmetic mean is defined by

$$A\nabla_p B := (1-p)A + pB,$$

for any  $p \in [0, 1]$ .

The relative operator entropy  $S(A|B)$  in [5] is defined by

$$S(A|B) := A^{\frac{1}{2}} \left( \log A^{-\frac{1}{2}} B A^{-\frac{1}{2}} \right) A^{\frac{1}{2}},$$

where  $A$  and  $B$  are two invertible positive operators on a Hilbert space. As a parametric extension of the relative operator entropy, Yanagi, Kuriyama and Furuichi [19] defined Tsallis relative operator entropy which is an operator version of Tsallis relative entropy in quantum system due to Abe [1], also see [9]:

$$T_p(A|B) := \frac{A^{\frac{1}{2}} \left( A^{-\frac{1}{2}} B A^{-\frac{1}{2}} \right)^p A^{\frac{1}{2}} - A}{p}, \quad p \in (0, 1].$$

*Mathematics subject classification* (2010): 47A63, 46L05, 47A60.

*Keywords and phrases:* Relative operator entropy, Tsallis relative operator entropy, operator inequality, Hermite-Hadamard's inequality, Young's inequality.

The authors would like to thank editor and referee for providing valuable comments to improve our manuscript. The author (S. F.) was partially supported by JSPS KAKENHI Grant Number 16K05257.

Notice that  $T_p(A|B)$  can be written by using the notation of  $A\#_p B$  as follows:

$$T_p(A|B) := \frac{A\#_p B - A}{p}, \quad p \in (0, 1].$$

The relation between relative operator entropy  $S(A|B)$  and Tsallis relative operator entropy  $T_p(A|B)$  was considered in [9, 20], as follows:

$$A - AB^{-1}A \leq T_{-p}(A|B) \leq S(A|B) \leq T_p(A|B) \leq B - A. \tag{1}$$

Some deeper properties of Tsallis relative operator entropy were proved in [6, 11, 13, 20].

The main result of the present paper is a set of bounds that are complementary to (1). Some of our inequalities improve well-known ones. Among other inequalities, it is shown that if  $A, B$  are invertible positive operators and  $p \in (0, 1]$ , then

$$\begin{aligned} A^{\frac{1}{2}} \left( \frac{A^{-\frac{1}{2}}BA^{-\frac{1}{2}} + I}{2} \right)^{p-1} \left( A^{-\frac{1}{2}}BA^{-\frac{1}{2}} - I \right) A^{\frac{1}{2}} &\leq T_p(A|B) \\ &\leq \frac{1}{2} (A\#_p B - A\sharp_{p-1} B + B - A), \end{aligned}$$

which is a considerable refinement of (1), where  $I$  is the identity operator. We also prove a reverse inequality involving Tsallis relative operator entropy  $T_p(A|B)$ .

**2. Refinements of the inequalities (1) via Hermite-Hadamard inequality**

An important ingredient in our approach is the following:

Let  $f$  be a convex function on  $[a, b] \subseteq \mathbb{R}$ ; the well-known Hermite-Hadamard’s inequality can be expressed as

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(t) dt \leq \frac{f(a)+f(b)}{2}. \tag{2}$$

**THEOREM 1.** *For any invertible positive operators  $A$  and  $B$  such that  $A \leq B$ , and  $p \in (0, 1]$  we have*

$$\begin{aligned} A^{\frac{1}{2}} \left( \frac{A^{-\frac{1}{2}}BA^{-\frac{1}{2}} + I}{2} \right)^{p-1} \left( A^{-\frac{1}{2}}BA^{-\frac{1}{2}} - I \right) A^{\frac{1}{2}} &\leq T_p(A|B) \\ &\leq \frac{1}{2} (A\#_p B - A\sharp_{p-1} B + B - A). \end{aligned} \tag{3}$$

*Proof.* Consider the function  $f(t) = t^{p-1}$ ,  $p \in (0, 1]$ . It is easy to check that  $f(t)$  is convex on  $[1, \infty)$ . Bearing in mind the fact

$$\int_1^x t^{p-1} dt = \frac{x^p - 1}{p},$$

and utilizing the left-hand side of Hermite-Hadamard inequality, one can see that

$$\left(\frac{x+1}{2}\right)^{p-1} (x-1) \leq \frac{x^p-1}{p}, \quad (4)$$

where  $x \geq 1$  and  $p \in (0, 1]$ . On the other hand, it follows from the right-hand side of Hermite-Hadamard inequality that

$$\frac{x^p-1}{p} \leq \left(\frac{x^{p-1}+1}{2}\right) (x-1), \quad (5)$$

for each  $x \geq 1$  and  $p \in (0, 1]$ .

Replacing  $x$  by  $A^{-\frac{1}{2}}BA^{-\frac{1}{2}}$  in (4) and (5), and multiplying  $A^{\frac{1}{2}}$  on both sides, we get the desired result (3).  $\square$

REMARK 1. Simple calculation gives for all  $x \geq 1$  and  $p \in (0, 1]$ ,

$$0 \leq 1 - \frac{1}{x} \leq \left(\frac{x+1}{2}\right)^{p-1} (x-1) \leq \frac{x^p-1}{p} \leq \left(\frac{x^{p-1}+1}{2}\right) (x-1) \leq x-1, \quad (6)$$

which means

$$\begin{aligned} 0 \leq A - AB^{-1}A &\leq A^{\frac{1}{2}} \left( \frac{A^{-\frac{1}{2}}BA^{-\frac{1}{2}} + I}{2} \right)^{p-1} (A^{-\frac{1}{2}}BA^{-\frac{1}{2}} - I) A^{\frac{1}{2}} \\ &\leq T_p(A|B) \\ &\leq \frac{1}{2} (A \#_p B - A \natural_{p-1} B + B - A) \\ &\leq B - A, \end{aligned}$$

for any invertible positive operators  $A$  and  $B$  such that  $A \leq B$ , and  $p \in (0, 1]$ . Therefore, our inequalities (3) improve the inequalities (1) for the case  $A \leq B$ .

PROPOSITION 1. For  $x \geq 1$  and  $\frac{1}{2} \leq p \leq 1$ ,

$$\frac{x-1}{\sqrt{x}} \leq \left(\frac{x+1}{2}\right)^{p-1} (x-1). \quad (7)$$

*Proof.* In order to prove (7), we set the function  $f_p(x) \equiv \left(\frac{x+1}{2}\right)^{p-1} - \frac{1}{\sqrt{x}}$  where  $x \geq 1$  and  $\frac{1}{2} \leq p \leq 1$ . Since  $\frac{df_p(x)}{dp} = \left(\frac{x+1}{2}\right)^{p-1} \ln\left(\frac{x+1}{2}\right)$ ,  $\frac{df_p(x)}{dp} \geq 0$  for  $x \geq 1$ . Thus, we have  $f_p(x) \geq f_{1/2}(x) = \frac{\sqrt{2x} - \sqrt{x+1}}{\sqrt{x(x+1)}} \geq 0$  for  $x \geq 1$ . Therefore, we have the inequality (7).  $\square$

REMARK 2. The first inequality (3) gives tight lower bound for the Tsallis relative operator entropy  $T_p(A|B)$  more than the eighth inequality in [8, Theorem 2.8 (i)], due to Proposition 1.

COROLLARY 1. For any invertible positive operators  $A$  and  $B$  such that  $A \geq B$ , and  $p \in (0, 1]$ , we have

$$\begin{aligned} A\#_p B - A\sharp_{p-1} B &\leq \frac{1}{2} (A\#_p B - A\sharp_{p-1} B + B - A) \leq T_p(A|B) \\ &\leq A^{\frac{1}{2}} \left( \frac{A^{-\frac{1}{2}} B A^{-\frac{1}{2}} + I}{2} \right)^{p-1} \left( A^{-\frac{1}{2}} B A^{-\frac{1}{2}} - I \right) A^{\frac{1}{2}} \\ &\leq A\sharp_{p+1} B - A\#_p B \leq 0. \end{aligned}$$

*Proof.* Put  $t = \frac{1}{x} \leq 1$  in the inequalities (6).  $\square$

REMARK 3. By numerical computations, we have  $\left(\frac{x^{p-1}+1}{2}\right)(x-1) - \frac{2(x-1)}{x+1} \simeq -0.83219$  for  $p = \frac{2}{3}, x = 0.01$ , and we also have  $\left(\frac{x^{p-1}+1}{2}\right)(x-1) - \frac{2(x-1)}{x+1} \simeq 0.216868$  for  $p = \frac{2}{3}, x = 0.1$ . Thus we do not have ordering between  $\left(\frac{x^{p-1}+1}{2}\right)(x-1)$  and  $\frac{2(x-1)}{x+1}$  for  $0 < x \leq 1$  and  $\frac{1}{2} \leq p \leq 1$ . So there is no ordering between the second inequality in Corollary 1 and the sixth inequality in [8, Theorem 2.8 (ii)].

THEOREM 2. For any invertible positive operators  $A$  and  $B$  such that  $A \leq B$  and  $p \in (0, 1]$ , we have the following inequalities,

$$L_p(A, B) + K_p(A, B) \leq T_p(A|B) \leq R_p(A, B) + K_p(A, B), \tag{8}$$

and

$$J_p(A, B) - 2R_p(A, B) \leq T_p(A|B) \leq J_p(A, B) - 2L_p(A, B), \tag{9}$$

where

$$\begin{aligned} K_p(A, B) &\equiv A^{1/2} \left( \frac{A^{-1/2} B A^{-1/2} + I}{2} \right)^{p-1} (A^{-1/2} B A^{-1/2} - I) A^{1/2}, \\ J_p(A, B) &\equiv \frac{1}{2} (A\#_p B - A\sharp_{p-1} B + B - A), \\ L_p(A, B) &\equiv \frac{1}{24} (p-1)(p-2) (A\#_p B - 3A\sharp_{p-1} B + 3A\sharp_{p-2} B - A\sharp_{p-3} B), \\ R_p(A, B) &\equiv \frac{1}{24} (p-1)(p-2) (A\sharp_3 B - 3A\sharp_2 B + 3B - A). \end{aligned}$$

*Proof.* According to [18, Theorem 1], if  $f : [a, b] \rightarrow \mathbb{R}$  is a twice differentiable function that there exists real constants  $m$  and  $M$  so that  $m \leq f'' \leq M$ , then

$$m \frac{(b-a)^2}{24} \leq \frac{1}{b-a} \int_a^b f(t) dt - f\left(\frac{a+b}{2}\right) \leq M \frac{(b-a)^2}{24}, \tag{10}$$

$$m \frac{(b-a)^2}{12} \leq \frac{f(a)+f(b)}{2} - \frac{1}{b-a} \int_a^b f(t) dt \leq M \frac{(b-a)^2}{12}. \tag{11}$$

Putting  $f(t) = t^{p-1}$  with  $p \in (0, 1]$  and  $a = 1, b = x$  in the above inequalities, then we have the desired results by a similar way to the proof of Theorem 1.  $\square$

REMARK 4. The first inequality of (8) and the second inequality of (9) give tighter bounds of Tsallis relative entropy  $T_p(A|B)$  than those in the inequalities (3), because of the following reasons.

- (i) The first inequality of (10) gives tight lower bound more than the first inequality of Hermite-Hadamard's inequality (2).
- (ii) The first inequality of (11) gives tight upper bound more than the second inequality of Hermite-Hadamard's inequality (2).

### 3. Some reverse inequalities via Young type inequalities

The scalar Young's inequality says that if  $a, b > 0$ , then

$$a^{1-p}b^p \leq (1-p)a + pb, \quad p \in [0, 1].$$

The following inequalities provide a refinement and a multiplicative reverse for Young's inequality with Kantorovich constant:

$$K^r(h, 2)a^{1-p}b^p \leq (1-p)a + pb \leq K^R(h, 2)a^{1-p}b^p, \quad (12)$$

where  $a, b > 0$ ,  $p \in [0, 1]$ ,  $r = \min\{p, 1-p\}$ ,  $R = \max\{p, 1-p\}$ ,  $K(h, 2) = \frac{(h+1)^2}{4h}$  and  $h = \frac{b}{a}$ . Notice that the first inequality in (12) was obtained by Zou *et al.* in [21, Corollary 3] while the second one was obtained by Liao *et al.* [17, Corollary 2.2].

In [14, 15], Kittaneh and Manasrah obtained another refinement and reverse of Young's inequality:

$$r(\sqrt{a} - \sqrt{b})^2 + a^{1-p}b^p \leq (1-p)a + pb \leq R(\sqrt{a} - \sqrt{b})^2 + a^{1-p}b^p, \quad (13)$$

where  $a, b > 0$ ,  $p \in [0, 1]$ ,  $r = \min\{p, 1-p\}$ ,  $R = \max\{p, 1-p\}$ . Further refinements and generalizations of Young's inequality have been obtained in [2, 7].

More interesting things happen when we apply these considerations to the operators. For instance, from the inequality (12) it follows that:

PROPOSITION 2. Let  $A, B$  be two invertible positive operators such that  $I < h'I \leq A^{-\frac{1}{2}}BA^{-\frac{1}{2}} \leq hI$  or  $0 < hI \leq A^{-\frac{1}{2}}BA^{-\frac{1}{2}} \leq h'I < I$ , then

$$K^r(h', 2)A\#_pB \leq A\nabla_pB \leq K^R(h, 2)A\#_pB, \quad (14)$$

where  $p \in [0, 1]$ ,  $r = \min\{p, 1-p\}$ ,  $R = \max\{p, 1-p\}$ .

*Proof.* The choice  $a = 1$  and  $b = x$  reduces (12) to the inequality

$$K^r(x, 2)x^p \leq (1-p) + px \leq K^R(x, 2)x^p.$$

Whence

$$\min_{1 < h' \leq x \leq h} K^r(x, 2)T^p \leq (1-p)I + pT \leq \max_{1 < h' \leq x \leq h} K^R(x, 2)T^p, \quad (15)$$

for any positive operator  $T$  such that  $I < h'I \leq T \leq hI$ . On choosing  $T = A^{-\frac{1}{2}}BA^{-\frac{1}{2}}$  in (15) we get

$$\begin{aligned} \min_{1 < h' \leq x \leq h} K^r(x, 2) \left( A^{-\frac{1}{2}}BA^{-\frac{1}{2}} \right)^p &\leq (1-p)I + pA^{-\frac{1}{2}}BA^{-\frac{1}{2}} \\ &\leq \max_{1 < h' \leq x \leq h} K^R(x, 2) \left( A^{-\frac{1}{2}}BA^{-\frac{1}{2}} \right)^p. \end{aligned}$$

Since  $K(x, 2)$  is increasing for  $x > 1$  we can write

$$K^r(h', 2) \left( A^{-\frac{1}{2}}BA^{-\frac{1}{2}} \right)^p \leq (1-p)I + pA^{-\frac{1}{2}}BA^{-\frac{1}{2}} \leq K^R(h, 2) \left( A^{-\frac{1}{2}}BA^{-\frac{1}{2}} \right)^p. \tag{16}$$

Multiplying both sides by  $A^{\frac{1}{2}}$  to inequality (16), we obtain the required inequality (14).

Another case follows from the fact that  $K(x, 2)$  is decreasing for  $x < 1$ .  $\square$

Ando’s inequality [3, Theorem 3] says that if  $A, B$  are positive operators and  $\Phi$  is a positive linear mapping, then

$$\Phi(A\#_pB) \leq \Phi(A)\#_p\Phi(B), \quad p \in [0, 1]. \tag{17}$$

Concerning inequality (17), we have the following corollary:

**COROLLARY 2.** *Let  $A, B$  be two invertible positive operators such that  $I < h'I \leq A^{-\frac{1}{2}}BA^{-\frac{1}{2}} \leq hI$  or  $0 < hI \leq A^{-\frac{1}{2}}BA^{-\frac{1}{2}} \leq h'I < I$ . Let  $\Phi$  be positive linear map on  $\mathcal{B}(\mathcal{H})$ , then*

$$\begin{aligned} \frac{K^r(h', 2)}{K^R(h, 2)} \Phi(A\#_pB) &\leq \frac{1}{K^R(h, 2)} \Phi(A\nabla_pB) \leq \Phi(A)\#_p\Phi(B) \\ &\leq \frac{1}{K^r(h', 2)} \Phi(A\nabla_pB) \leq \frac{K^R(h, 2)}{K^r(h', 2)} \Phi(A\#_pB), \end{aligned} \tag{18}$$

where  $p \in [0, 1]$ ,  $r = \min\{p, 1-p\}$  and  $R = \max\{p, 1-p\}$ .

*Proof.* If we apply positive linear map  $\Phi$  in (14) we infer

$$K^r(h', 2) \Phi(A\#_pB) \leq \Phi(A\nabla_pB) \leq K^R(h, 2) \Phi(A\#_pB). \tag{19}$$

On the other hand, if we take  $A = \Phi(A)$  and  $B = \Phi(B)$  in (14) we can write

$$K^r(h', 2) \Phi(A)\#_p\Phi(B) \leq \Phi(A\nabla_pB) \leq K^R(h, 2) \Phi(A)\#_p\Phi(B). \tag{20}$$

Now, combining inequality (19) and inequality (20), we deduce the desired inequalities (18).  $\square$

**REMARK 5.** It is well-known that the generalized Kantorovich constant  $K(h, p)$  [12, Definition 1] is defined by

$$K(h, p) := \frac{h^p - h}{(p-1)(h-1)} \left( \frac{p-1}{p} \frac{h^p-1}{h^p-h} \right)^p, \tag{21}$$

for all  $p \in \mathbb{R}$ . By virtue of a generalized Kantorovich constant, in the matrix setting, Bourin *et al.* in [4, Theorem 6] gave the following reverse of Ando's inequality for a positive linear map:

Let  $A$  and  $B$  be positive operators such that  $mA \leq B \leq MA$ , and let  $\Phi$  be a positive linear map. Then

$$\Phi(A) \#_p \Phi(B) \leq \frac{1}{K(h, p)} \Phi(A \#_p B), \quad p \in [0, 1] \quad (22)$$

where  $h = \frac{M}{m}$ . The above result naturally extends one proved in Lee [16, Theorem 4] for  $p = \frac{1}{2}$ .

Of course the constant  $\frac{K^R(h, 2)}{K^r(h', 2)}$  is not better than  $\frac{1}{K(\frac{1}{h'}, p)}$ . Concerning the sharpness of the estimate (22), see [4, Lemma 7]. However our bounds on  $\Phi(A) \#_p \Phi(B)$  are calculated by the original Kantorovich constant  $K(h, 2)$  without the generalized one  $K(h, p)$ . It is also interesting our bounds on  $\Phi(A) \#_p \Phi(B)$  are expressed by  $\Phi(A \nabla_p B)$  with only one constant either  $h$  or  $h'$ .

After discussion on inequalities related to the operator mean with positive linear map, we give a result on Tsallis relative operator entropy (which is the main theme in this paper) with a positive linear map. It is well-known that Tsallis relative operator entropy has the following information monotonicity:

$$\Phi(T_p(A|B)) \leq T_p(\Phi(A) | \Phi(B)). \quad (23)$$

Utilizing (13), we have the following counterpart of (23):

**THEOREM 3.** *Let  $A, B$  be two invertible positive operators. Let  $\Phi$  be normalized positive linear map on  $\mathcal{B}(\mathcal{H})$ , then*

$$\begin{aligned} & \frac{2r}{p} (\Phi(A \nabla B) - \Phi(A) \# \Phi(B)) + T_p(\Phi(A) | \Phi(B)) \\ & \leq \Phi(B - A) \leq \frac{2R}{p} (\Phi(A \nabla B) - \Phi(A \# B)) + \Phi(T_p(A|B)), \end{aligned} \quad (24)$$

where  $p \in (0, 1]$ ,  $r = \min\{p, 1 - p\}$  and  $R = \max\{p, 1 - p\}$ .

*Proof.* Using the method of the proof of Proposition 2 and Corollary 2 for the inequality (13), we can obtain that

$$\begin{aligned} & 2r (\Phi(A \nabla B) - \Phi(A) \# \Phi(B)) + \Phi(A) \#_p \Phi(B) \\ & \leq \Phi(A \nabla_p B) \leq 2R (\Phi(A \nabla B) - \Phi(A \# B)) + \Phi(A \#_p B). \end{aligned}$$

A simple calculation shows that

$$\begin{aligned} & \frac{2r}{p} (\Phi(A \nabla B) - \Phi(A) \# \Phi(B)) + \frac{\Phi(A) \#_p \Phi(B) - \Phi(A)}{p} \\ & \leq \Phi(B - A) \leq \frac{2R}{p} (\Phi(A \nabla B) - \Phi(A \# B)) + \frac{\Phi(A \#_p B) - \Phi(A)}{p}. \end{aligned}$$

Apparently, the above inequality is equivalent to inequality (24). The proof is completed.  $\square$

The following example may render the above statement clearer.

EXAMPLE 1. Let  $\Phi(X) = U^*XU$  ( $X \in \mathcal{M}_2$ ) where  $U = \begin{pmatrix} \frac{\sqrt{2}}{2} & \frac{\sqrt{2}}{2} \\ -\frac{\sqrt{2}}{2} & \frac{\sqrt{2}}{2} \end{pmatrix}$ . If  $A = \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix}$  and  $B = \begin{pmatrix} 6 & 2 \\ 2 & 4 \end{pmatrix}$  and  $p = \frac{1}{4}$  we get

$$\frac{2r}{p} (\Phi(A\nabla B) - \Phi(A)\#\Phi(A)) + (T_p(\Phi(A)|\Phi(A))) = \begin{pmatrix} 0.486 & 0.443 \\ 0.443 & 5.638 \end{pmatrix}$$

$$\Phi(B - A) = \begin{pmatrix} 0.5 & 0.5 \\ 0.5 & 6.5 \end{pmatrix}$$

and

$$\frac{2R}{p} (\Phi(A\nabla B) - \Phi(A\#B)) + \Phi(T_p(A|B)) = \begin{pmatrix} 0.562 & 0.951 \\ 0.951 & 13.521 \end{pmatrix}$$

which shows that inequality (24) is true.

Tsallis relative entropy  $D_p(A||B)$  for two positive operators  $A$  and  $B$  is defined by:

$$D_p(A||B) := \frac{Tr[A] - Tr[A^{1-p}B^p]}{p}, \quad p \in (0, 1].$$

In information theory, relative entropy (divergence) is usually defined for density operators which are positive operators with unit trace. However we consider Tsallis relative entropy defined for positive operators to derive the relation with Tsallis relative operator entropy. If  $A$  and  $B$  are positive operators, then

$$Tr[A - B] \leq D_p(A||B) \leq -Tr[T_p(A|B)], \quad p \in (0, 1]. \tag{25}$$

Note that the first inequality of (25) is due to Furuta [13, Proposition F] and the second inequality is due to Furuichi *et al.* [10, Theorem 2.2].

As a direct consequence of Theorem 3, we have the following interesting relation.

COROLLARY 3. *Let  $A, B$  be two positive operators on a finite dimensional Hilbert space  $\mathcal{H}$ , then*

$$\begin{aligned} & \frac{2R}{p} \left( Tr[A\#B] - \frac{Tr[A+B]}{2} \right) - Tr[T_p(A|B)] \\ & \leq Tr[A - B] \leq \frac{2r}{p} \left( \sqrt{Tr[A]Tr[B]} - \frac{Tr[A+B]}{2} \right) + D_p(A||B), \end{aligned} \tag{26}$$

where  $p \in (0, 1]$ ,  $r = \min\{p, 1 - p\}$  and  $R = \max\{p, 1 - p\}$ .



*Proof.* Taking  $\Phi(X) = \frac{1}{\dim \mathcal{H}} \text{Tr}[X]$  in (24), we have

$$\begin{aligned} & \frac{2r}{p} \left( \frac{\text{Tr}[A+B]}{2} - \sqrt{\text{Tr}[A]\text{Tr}[B]} \right) + \frac{(\text{Tr}[A])^{1-p}(\text{Tr}[B])^p - \text{Tr}[A]}{p} \\ & \leq \text{Tr}[B-A] \leq \frac{2R}{p} \left( \frac{\text{Tr}[A+B]}{2} - \text{Tr}[A\#B] \right) + \text{Tr}[T_p(A|B)]. \end{aligned} \tag{27}$$

It is not too difficult to see that (27) can be also reformulated in the following way

$$\begin{aligned} & \frac{2R}{p} \left( \text{Tr}[A\#B] - \frac{\text{Tr}[A+B]}{2} \right) - \text{Tr}[T_p(A|B)] \\ & \leq \text{Tr}[A-B] \leq \frac{2r}{p} \left( \sqrt{\text{Tr}[A]\text{Tr}[B]} - \frac{\text{Tr}[A+B]}{2} \right) + \frac{\text{Tr}[A] - (\text{Tr}[A])^{1-p}(\text{Tr}[B])^p}{p}. \end{aligned}$$

Now, having in mind that  $\text{Tr}[A^{1-p}B^p] \leq (\text{Tr}[A])^{1-p}(\text{Tr}[B])^p$ , we infer

$$\begin{aligned} & \frac{2R}{p} \left( \text{Tr}[A\#B] - \frac{\text{Tr}[A+B]}{2} \right) - \text{Tr}[T_p(A|B)] \\ & \leq \text{Tr}[A-B] \leq \frac{2r}{p} \left( \sqrt{\text{Tr}[A]\text{Tr}[B]} - \frac{\text{Tr}[A+B]}{2} \right) + \frac{\text{Tr}[A] - \text{Tr}[A^{1-p}B^p]}{p}, \end{aligned}$$

which, in turn, leads to (26).  $\square$

REMARK 6. If  $A$  and  $B$  are density operators, then the second inequality of (26) implies the non-negativity of Tsallis relative entropy,  $D_p(A||B) \geq 0$ . If  $p \in [\frac{1}{2}, 1]$ , then the first inequality of (26) implies

$$\text{Tr}[T_{1/2}(A|B)] \leq \text{Tr}[T_p(A|B)].$$

If  $p \in (0, \frac{1}{2}]$ , then the first inequality of (26) also implies

$$(1-p)\text{Tr}[T_{1/2}(A|B)] + (2p-1)\text{Tr}[B-A] \leq p\text{Tr}[T_p(A|B)].$$

REFERENCES

[1] S. ABE, *Monotonic decrease of the quantum non-additive divergence by projective measurements*, Phys. Lett. A., **312** (5–6) (2003), 336–338.  
 [2] Y. AL-MANASRAH, F. KITTANEH, *A generalization of two refined Young inequalities*, Positivity., **19** (4) (2015), 757–768.  
 [3] T. ANDO, *Concavity of certain maps on positive definite matrices and applications to Hadamard products*, Linear Algebra Appl., **26** (1979), 203–241.  
 [4] J. C. BOURIN, E.-Y. LEE, M. FUJII, Y. SEO, *A matrix reverse Holder inequality*, Linear Algebra Appl., **431** (11) (2009), 2154–2159.  
 [5] J. I. FUJII, E. KAMEI, *Relative operator entropy in non-commutative information theory*, Math. Japon., **34** (1989), 341–348.  
 [6] S. FURUICHI, *Inequalities for Tsallis relative entropy and generalized skew information*, Linear Multilinear Algebra., **59** (10) (2011), 1143–1158.

- [7] S. FURUICHI, N. MINCULETE, *Alternative reverse inequalities for Young's inequality*, J. Math Inequal., **5** (4) (2011), 595–600.
- [8] S. FURUICHI, *Precise estimates of bounds on relative operator entropies*, Math. Inequal. Appl., **18** (3) (2015), 869–877.
- [9] S. FURUICHI, K. YANAGI AND K. KURIYAMA, *A note on operator inequalities of Tsallis relative operator entropy*, Linear Algebra Appl., **407** (2005), 19–31.
- [10] S. FURUICHI, K. YANAGI, K. KURIYAMA, *Fundamental properties of Tsallis relative entropy*, J. Math. Phys., **45** (2004), 4868–4877.
- [11] S. FURUICHI, N. MINCULETE, F. C. MITROI, *Some inequalities on generalized entropies*, J. Inequal. Appl., **1** (2012), 226.
- [12] T. FURUTA, *Basic properties of the generalized Kantorovich constant  $K(p) = \frac{h^p-h}{(p-1)(h-1)}$*  ·  $\left(\frac{p-1}{p} \frac{h^p-1}{h^p-h}\right)^p$ , Acta Sci. Math (Szeged), **70** (2004), 319–337.
- [13] T. FURUTA, *Two reverse inequalities associated with Tsallis relative operator entropy via generalized Kantorovich constant and their applications*, Linear Algebra Appl., **412** (2) (2006), 526–537.
- [14] F. KITTANEH AND Y. MANASRAH, *Improved Young and Heinz inequalities for matrix*, J. Math. Anal. Appl., **361** (2010), 262–269.
- [15] F. KITTANEH AND Y. MANASRAH, *Reverse Young and Heinz inequalities for matrices*, Linear Multilinear Algebra., **59** (2011), 1031–1037.
- [16] E.-Y. LEE, *A matrix reverse Cauchy-schwarz inequality*, Linear Algebra Appl., **430** (2) (2009), 805–810.
- [17] W. LIAO, J. WU, J. ZHAO, *New versions of reverse Young and Heinz mean inequalities with the Kantorovich constant*, Taiwanese J. Math., **19** (2) (2015), 467–479.
- [18] C. P. NICULESCU, L. E. PERSSON, *Old and new on the Hermite-Hadamard inequality*, Real Anal. Exchange., **29** (2) (2004), 663–686.
- [19] K. YANAGI, K. KURIYAMA, S. FURUICHI, *Generalized Shannon inequalities based on Tsallis relative operator entropy*, Linear Algebra Appl., **394** (2005), 109–118.
- [20] L. ZOU, *Operator inequalities associated with Tsallis relative operator entropy*, Math. Inequal. Appl., **18** (2) (2015), 401–406.
- [21] H. ZUO, G. SHI, M. FUJII, *Refined Young inequality with Kantorovich constant*, J. Math. Inequal., **5** (4) (2011), 551–556.

(Received March 4, 2017)

Hamid Reza Moradi  
Young Researchers and Elite Club, Mashhad Branch  
Islamic Azad University  
Mashhad, Iran  
e-mail: hrmoradi@mshdiau.ac.ir

Shigeru Furuichi  
Department of Information Science  
College of Humanities and Sciences, Nihon University  
3-25-40, Sakurajyousui, Setagaya-ku, Tokyo, 156-8550, Japan  
e-mail: furuichi@chs.nihon-u.ac.jp

Nicușor Minculete  
Faculty of Mathematics and Computer Science  
Transilvania University of Braşov  
Str. Iuliu Maniu, nr. 50, Braşov, 500091, Romania  
e-mail: minculeteten@yahoo.com