

## A COUNTEREXAMPLE TO A QUESTION OF BAPAT & SUNDER

STEPHEN DRURY

(Communicated by J.-C. Bourin)

*Abstract.* The objective of this article is to provide a counterexample to a question of Bapat and Sunder concerning the relative magnitudes of the permanent of a positive semidefinite matrix and the largest eigenvalue of a related matrix. We also discuss the significance of this result in connection with the eigenvalues of the Schur matrix.

### 1. Introduction

The permanent of a  $n \times n$  matrix  $A = (a_{jk})$  is defined as the quantity

$$\text{per}(A) = \sum_{\sigma \in S_n} \prod_{j=1}^n a_{j, \sigma(j)}.$$

It is an important concept useful in combinatorial applications. For a recent survey of permanent inequalities and open questions the reader is referred to [11] and the references therein.

Let  $A$  be a  $n \times n$  positive semidefinite matrix. Denote by  $A(i, j)$  the  $(n-1) \times (n-1)$  submatrix of  $A$  obtained by deleting the  $i^{\text{th}}$  row and  $j^{\text{th}}$  column of  $A$ . Now define a  $n \times n$  matrix  $B$  by

$$b_{ij} = a_{ij} \text{per}(A(i, j)). \tag{1}$$

Then it is clear that  $B$  is again a positive semidefinite matrix and it follows from the Laplace expansion of the permanent that all its row and column sums are equal to  $\text{per}(A)$ . Thus,  $\text{per}(A)$  is an eigenvalue of  $B$  and  $\mathbb{1}$  is the corresponding eigenvector.

In [2, Conjecture 3], Bapat and Sunder raise the question of whether  $\text{per}(A)$  is necessarily the largest eigenvalue of  $B$ . We provide a counterexample to this question.

*Mathematics subject classification* (2010): 15A15.

*Keywords and phrases:* Permanent, Hadamard product, Oppenheim's inequality, permanent on top.

## 2. The counterexample

With  $n = 8$ , we take

$$X = \begin{pmatrix} -7+4i & 9-3i & -6+2i & 3+4i & 7+6i & 4-4i & i & 5-8i \\ 4-5i & 1+4i & -8-2i & -7+4i & 1-4i & 1-8i & 8-6i & 1-3i \end{pmatrix}$$

and set  $A = X^*X$ . Then we obtain

$$A = \begin{pmatrix} 106 & -91+6i & 28-38i & -53-59i & -1-81i & -15i & 66+9i & -48+29i \\ -91-6i & 107 & -76+30i & 24+77i & 30+67i & 17-36i & -19-29i & 58-64i \\ 28+38i & -76-30i & 108 & 38-76i & -30-16i & -24+82i & -50+58i & -48+64i \\ -53+59i & 24-77i & 38+76i & 90 & 22+14i & -43+24i & -76+13i & -36-27i \\ -1+81i & 30-67i & -30+16i & 22-14i & 102 & 37-56i & 38+33i & -85i \\ 15i & 17+36i & -24-82i & -43-24i & 37+56i & 97 & 52+62i & 77-7i \\ 66-9i & -19+29i & -50-58i & -76-13i & 38-33i & 52-62i & 101 & 18-23i \\ -48-29i & 58+64i & -48-64i & -36+27i & 85i & 77+7i & 18+23i & 99 \end{pmatrix}$$

a rank two positive semidefinite matrix. Calculations show that

$$\text{per}(A) = 2977257622144118400$$

and that the largest eigenvalue of  $B$  exceeds 3028080150918724811.

This example was found using a hill-climbing computer search and the example found was rounded for easy presentation. The ratio  $\lambda_1(B)/\text{per}(A)$  found in the search was approximately 1.01956. No example was found for  $n = 7$ .

## 3. The Schur matrix

For a positive semidefinite  $n \times n$  matrix  $A$ , define the convolution operator  $\Pi(A)$  on the symmetric group  $S_n$  by its matrix

$$\Pi(A)_{\sigma,\rho} = \prod_{j=1}^n a_{\sigma(j),\rho(j)}.$$

This (usually huge) matrix is known as the Schur matrix. As is well-known,  $\Pi(A)$  is unitarily similar to a block diagonal matrix indexed by the set of irreducible representations of  $S_n$ . The diagonal block corresponding to the irreducible representation  $\pi$  is a matrix multiplication operator by a hermitian matrix  $\widehat{\Pi(A)}(\pi)$ . Thus, the eigenvalues of  $\Pi(A)$  coming from the diagonal block corresponding to  $\pi$  are the eigenvalues of  $\widehat{\Pi(A)}(\pi)$  each repeated  $d_\pi$  times where  $d_\pi$  is the dimension of  $\pi$ . The reader may consult [5] for the Fourier analysis of compact (and hence finite) groups.

The permanent on top conjecture was originally formulated by G. Soules in his Ph.D. dissertation 1966 [10] and published in [7]. It asks if the largest eigenvalue of  $\Pi(A)$  is  $\text{per}(A)$ , namely the eigenvalue arising from the trivial representation. In 2016, Shchesnovich [9] presented an example of a  $5 \times 5$  positive semidefinite matrix  $A$  and a unit column vector  $X$  indexed by  $S_5$  such that  $X^*\Pi(A)X > \text{per}(A)$  thereby demolishing Soule's conjecture.

The irreducible representations of  $S_n$  are well-known to be in one-to-one correspondence with the Ferrers diagrams with  $n$  entries. Details can be found in [6, 8]. Shchesnovich does not identify in his paper the representation that is responsible for his counterexample, but calculations reveal that it is the Ferrers diagram (3, 2) that has 3 entries in the first row and 2 in the second.

It is well-known that the representation  $\sigma \mapsto P(\sigma)$ , the representation that takes each permutation to its permutation matrix decomposes as the direct sum of the trivial representation (Ferrers diagram  $(n)$ ) and the representation  $\pi_1$  with Ferrers diagram  $(n - 1, 1)$ . We have, denoting  $\varepsilon$  the identity permutation,

$$\sum_{\rho \in S_n} \Pi(A)_{\rho, \varepsilon} P_{j,k}(\rho) = \sum_{\rho \in S_n} \left( \prod_{i=1}^n a_{\rho i, i} \right) \delta_{j, \rho k} = a_{jk} \text{per}(A(j, k)) = b_{jk}.$$

It follows that the eigenvalues of  $B$  are  $\text{per}(A)$  together with the eigenvalues of  $\widehat{\Pi(A)}(\pi_1)$ . Thus for  $n = 8$  we have yet another counterexample to the permanent on top conjecture [9, 3, 4].

#### 4. A new question

So, the relevant question is now:

QUESTION 1. *For a given  $n$ , which irreducible representations  $\pi$  of  $S_n$  have the property that the largest eigenvalue of  $\widehat{\Pi(A)}(\pi)$  is bounded above by  $\text{per}(A)$  for every positive definite  $n \times n$  hermitian matrix  $A$ ?*

The branching rule is a rule that determines how a given representation of  $S_n$  decomposes when it is restricted to  $S_m$ , the subgroup of  $S_n$  of permutations of  $\{1, 2, \dots, m\}$  for  $m < n$ . It is a consequence of the branching rule [8, §2.8] that if the Ferrers diagram of  $\pi$  contains either of the Ferrers diagrams (3, 2) or (7, 1) then the representation does not have the property of Question 1. Here, we are using the word ‘contains’ in a very loose sense. A Ferrers diagram  $\alpha$  contains another  $\beta$  if each row count of  $\beta$  is dominated by the corresponding row count of  $\alpha$ . This means that  $\beta$  ‘fits inside’  $\alpha$ .

CONJECTURE 1. *A representation satisfies the property asked in the question if it contains neither of the Ferrers diagrams (3, 2) or (7, 1).*

Some other conjectures that we believe might be true and that do not appear in [11] are:

CONJECTURE 2. *If  $A$  is a real rank two correlation matrix (i.e.  $a_{jk} = \cos(\theta_j - \theta_k)$  with the  $\theta_j$  real) then  $\text{per}(A \circ A) \leq \text{per}(A)$ . Here  $\circ$  denotes the Hadamard (entry-wise) product.*

This is a special case of a question raised in [1].

CONJECTURE 3. *If  $A$  is a real positive semidefinite matrix then  $\lambda_1(B) = \text{per}(A)$ . Here  $B$  is defined by (1) and  $\lambda_1(B)$  is its largest eigenvalue.*

## REFERENCES

- [1] R. B. BAPAT, V. S. SUNDER, *On majorization and Schur products*, Linear Algebra Appl. **72** (1985), 107–117.
- [2] R. B. BAPAT, V. S. SUNDER, *An extremal property of the permanent and the determinant*, Linear Algebra Appl. **76** (1986), 153–163.
- [3] S. W. DRURY, *A Counterexample to a Question of Bapat and Sunder*, Electronic Journal of Linear Algebra **31** (2016), 69–70.
- [4] S. W. DRURY, *A real counterexample to two inequalities involving permanents*, Mathematical Inequalities and Applications **20** (2017), 349–352.
- [5] C. F. DUNKL, D. E. RAMIREZ, *Topics in harmonic analysis*, Appleton–Century–Crofts, New York, 1971.
- [6] G. D. JAMES, *The Representation Theory of Symmetric Groups*, Lecture Notes in Mathematics, vol. 682, Springer–Verlag, New York, 1978.
- [7] H. MINC, *Theory of permanents 1978–1981*, Linear Multilinear Algebra **12** (1983), 227–263.
- [8] B. E. SAGAN, *The Symmetric Group*, Wadsworth & Brooks/Cole, Pacific Grove, 1991.
- [9] V. S. SHCHESNOVICH, *The permanent-on-top conjecture is false*, Linear Algebra Appl. **490** (2016), 196–201.
- [10] G. SOULES, *Matrix functions and the Laplace expansion theorem*, Ph. D. Dissertation, University of California – Santa Barbara, July, 1966.
- [11] F. ZHANG, *An update on a few permanent conjectures*, Special Matrices **4** (2016), 305–316.

(Received September 11, 2017)

*Stephen Drury*  
*Department of Mathematics and Statistics*  
*McGill University*  
*Montreal, Canada H3A 0B9*  
*e-mail: stephen.drury@mcgill.ca*