

A HILBERT-TYPE FRACTIONAL INTEGRAL INEQUALITY WITH THE KERNEL OF MITTAG-LEFFLER FUNCTION AND ITS APPLICATIONS

QIONG LIU

(Communicated by J. Pečarić)

Abstract. By using the theory of the local fractional calculus and the methods of weight function, a Hilbert-type fractional integral inequality with the kernel of Mittag-Leffler function and its equivalent form are given. Their constant factors are proved being the best possible, and its applications are discussed briefly.

1. Introduction

Assuming that $f, g \geq 0$, $0 < \int_0^\infty f^2(x)dx < \infty$, $0 < \int_0^\infty g^2(y)dy < \infty$, we have the famous Hilbert's integral inequality and its equivalent form (cf. [1, 2]):

$$\int_0^\infty \int_0^\infty \frac{f(x)g(y)}{x+y} dx dy < \pi \left\{ \int_0^\infty f^2(x)dx \right\}^{\frac{1}{2}} \left\{ \int_0^\infty g^2(y)dy \right\}^{\frac{1}{2}}, \quad (1)$$

$$\int_0^\infty \left[\int_0^\infty \frac{f(x)}{x+y} dx \right]^2 dy < \pi^2 \int_0^\infty f^2(x)dx, \quad (2)$$

where the constant π and π^2 are the best possible. There have been a number of improvements and extensions on inequalities (1) and (2) (cf. [3–10]), which are important in the mathematical analysis and its applications (cf. [1, 2, 8]). In 2010, Yang [11] presented a new Hilbert-type integral inequality and its equivalent form, as follows:

If $p > 1$, $\frac{1}{p} + \frac{1}{q} = 1$, $f, g > 0$, $0 < \int_0^\infty \frac{1}{x} f^p(x)dx < \infty$, $0 < \int_0^\infty \frac{1}{y} g^q(y)dy < \infty$, then

$$\int_0^\infty \int_0^\infty e^{-xy} f(x)g(y) dx dy < \left\{ \int_0^\infty \frac{1}{x} f^p(x)dx \right\}^{\frac{1}{p}} \left\{ \int_0^\infty \frac{1}{y} g^q(y)dy \right\}^{\frac{1}{q}}, \quad (3)$$

$$\int_0^\infty y \left[\int_0^\infty e^{-xy} f(x)dx \right]^p dy < \int_0^\infty \frac{1}{x} f^p(x)dx. \quad (4)$$

In recent years, the theory of fractional calculus has been developed rapidly, and it has been widely used in the fields of science and engineering. Some researchers have used the fractal theory to discuss and generalize some classical inequalities (cf. [12–14]), but the research on the Hilbert-type fractional integral inequalities are still not involved. In this paper, by using the theory of local fractional calculus and the method of weight function to make a meaningful attempt, a Hilbert-type fractional integral inequality and its equivalent form are established.

Mathematics subject classification (2010): 26D15, 26D10.

Keywords and phrases: Fractal space, Hilbert-type fractional integral inequality, weight function.

This work was supported by the National Natural Science Foundations of China (No: 11171280) and Scientific Support Project of Hunan Province Education Department of China (No: 10C1186).

2. Preliminaries

DEFINITION 1. (cf. [15, 16]) The α -type set of set Ω are defined as the set Ω^α , where $0 < \alpha \leq 1$. This set Ω^α is called a fractional set. The set Ω is called base set of fractional set.

Let \mathbb{R}^α ($0 < \alpha \leq 1$) be the α -type set of real line numbers, if $a^\alpha, b^\alpha, c^\alpha \in \mathbb{R}^\alpha$, then (cf. [15, 16])

- 1° $a^\alpha + b^\alpha \in \mathbb{R}^\alpha, a^\alpha b^\alpha \in \mathbb{R}^\alpha$;
- 2° $a^\alpha + b^\alpha = b^\alpha + a^\alpha = (a+b)^\alpha = (b+a)^\alpha$;
- 3° $a^\alpha + (b+c)^\alpha = (a+b)^\alpha + c^\alpha$;
- 4° $a^\alpha b^\alpha = b^\alpha a^\alpha = (ab)^\alpha = (ba)^\alpha$;
- 5° $a^\alpha (b^\alpha c^\alpha) = (a^\alpha b^\alpha) c^\alpha$;
- 6° $a^\alpha (b^\alpha + c^\alpha) = a^\alpha b^\alpha + a^\alpha c^\alpha$;
- 7° $a^\alpha + 0^\alpha = a^\alpha, a^\alpha 1^\alpha = a^\alpha, (-a)^\alpha = -a^\alpha$;
- 8° If $a < b$, then $a^\alpha < b^\alpha$. If $a^\alpha < b^\alpha$, then $a < b$.

DEFINITION 2. (cf. [15, 16]) A non-differentiable function $f : \mathbb{R} \rightarrow \mathbb{R}^\alpha$ ($0 < \alpha \leq 1$), $x \rightarrow f(x)$ is called to be local fractional continuous at x_0 , if for any $\varepsilon > 0$, there exists $\delta > 0$, such that $|f(x) - f(x_0)| < \varepsilon$ whenever $|x - x_0| < \delta$. If $f(x)$ is local fractional continuous on the interval (a, b) , we denote $f(x) \in C_\alpha(a, b)$.

DEFINITION 3. (cf. [15, 16]) The local fractional derivative of $f(x)$ of order α ($0 < \alpha \leq 1$) at x_0 is defined by

$$f^{(\alpha)}(x_0) = \frac{d^\alpha f(x)}{dx^\alpha} \Big|_{x=x_0} = \lim_{x \rightarrow x_0} \frac{\Gamma(1 + \alpha)(f(x) - f(x_0))}{(x - x_0)^\alpha}.$$

If for all $x \in I \subset \mathbb{R}$, there exists $f^{(k+1)\alpha}(x) = \overbrace{D_x^\alpha \cdots D_x^\alpha}^{k+1} f(x)$, then we denote $f \in D_{(k+1)\alpha}(I)$, where $k = 0, 1, 2, \dots$.

DEFINITION 4. (cf. [15, 16]) Let $f(x) \in C_\alpha(a, b)$. Then the local fractional integral is defined by

$${}_a I_b^\alpha f(x) = \frac{1}{\Gamma(1 + \alpha)} \int_a^b f(t)(dt)^\alpha = \frac{1}{\Gamma(1 + \alpha)} \lim_{\Delta t \rightarrow 0} \sum_{i=1}^N f(t_i)(\Delta t_i)^\alpha,$$

with $\Delta t_i = t_i - t_{i-1}$ ($i = 1, \dots, N$), $\Delta t = \max_{1 \leq i \leq N} \{\Delta t_i\}$, and $a = t_0 < t_1 < \dots < t_{N-1} < t_N = b$ is partition of interval $[a, b]$. Here, it follows that ${}_a I_b^\alpha f(x) = 0$ if $a = b$, ${}_a I_b^\alpha f(x) = -{}_b I_a^\alpha f(x)$ if $a < b$.

DEFINITION 5. (cf. [15, 16]) Mittag-Leffler function defined on fractal set of fractal dimension α is by the expression:

$$E_\alpha(x^\alpha) := \sum_{k=0}^\infty \frac{x^{\alpha k}}{\Gamma(1 + \alpha k)}, \quad x \in \mathbb{R}, \quad \text{and } 0 < \alpha \leq 1.$$

DEFINITION 6. If $s \in \mathbb{C}$, let $f(x)$ denote a function which vanishes for negative values of x . Local fractional Laplace's transform $L_\alpha\{f(x)\}$ of order α is defined by the following expression, if it is finite:

$$L_\alpha\{f(x)\}(s) := \frac{1}{\Gamma(1 + \alpha)} \int_0^\infty E_\alpha(-s^\alpha x^\alpha) f(x) (dx)^\alpha,$$

DEFINITION 7. Gamma function on fractal set of fractal dimension α is defined by the following expression:

$$\Gamma_\alpha(x) := \frac{1}{\Gamma(1 + \alpha)} \int_0^\infty E_\alpha(-t^\alpha) t^{(x-1)\alpha} (dt)^\alpha, \quad 0 < \alpha \leq 1$$

LEMMA 1. If $m > -1$, $Re(s) > 0$, then we have

$$L_\alpha\{x^{m\alpha}\}(s) = \frac{\Gamma_\alpha(m + 1)}{s^{(m+1)\alpha}}.$$

Proof. Setting $t = sx$, by Definition 6 and Definition 7, we obtain

$$\begin{aligned} L_\alpha\{x^{m\alpha}\}(s) &= \frac{1}{\Gamma(1 + \alpha)} \int_0^\infty E_\alpha(-s^\alpha x^\alpha) x^{m\alpha} (dx)^\alpha \\ &= \frac{1}{s^{(m+1)\alpha} \Gamma(1 + \alpha)} \int_0^\infty E_\alpha(-t^\alpha) t^{m\alpha} (dt)^\alpha \\ &= \frac{\Gamma_\alpha(m + 1)}{s^{(m+1)\alpha}}. \quad \square \end{aligned}$$

LEMMA 2. (cf. [18, 19, 20]) If $f, g (\geq 0) \in C_\alpha(a, b)$, $F, G, h (\geq 0) \in C_\alpha(S^{(\beta)})$, $p > 1$, $\frac{1}{p} + \frac{1}{q} = 1$, $S^{(\beta)}$ is a fractal surface, then we have

1° Hölder inequality on the fractal set

$${}_a I_b^\alpha f(x) g(x) \leq \left\{ {}_a I_b^\alpha f^p(x) \right\}^{\frac{1}{p}} \left\{ {}_a I_b^\alpha g^q(x) \right\}^{\frac{1}{q}},$$

2° Hölder weighted inequality on the fractal set

$$\begin{aligned} &\frac{1}{\Gamma^2(1 + \alpha)} \iint_{S^{(\beta)}} h(x, y) F(x, y) G(x, y) (dx)^\alpha (dy)^\alpha \\ &\leq \left\{ \frac{1}{\Gamma^2(1 + \alpha)} \iint_{S^{(\beta)}} h(x, y) F^p(x, y) (dx)^\alpha (dy)^\alpha \right\}^{\frac{1}{p}} \\ &\quad \times \left\{ \frac{1}{\Gamma^2(1 + \alpha)} \iint_{S^{(\beta)}} h(x, y) G^q(x, y) (dx)^\alpha (dy)^\alpha \right\}^{\frac{1}{q}}. \end{aligned}$$

The inequality keeps the form of equality, then there exist constants A and B , such that they are not all zero and $AF^p(x, y) = BG^q(x, y)$.a.e. on $S^{(\beta)}$.

LEMMA 3. If $0 < \alpha \leq 1$, $\beta, \lambda > 0$, we define the weight function $\omega(\alpha, \beta, \lambda, x)$ as follows:

$$\omega(\alpha, \beta, \lambda, x) := \frac{1}{\Gamma(1+\alpha)} \int_0^\infty E_\alpha(-\lambda^\alpha(xy)^\alpha) x^{\beta\alpha} y^{(\beta-1)\alpha} (dy)^\alpha, \quad x \in (0, +\infty),$$

then we have

$$\omega(\alpha, \beta, \lambda, x) = \frac{\Gamma_\alpha(\beta)}{\lambda^{\beta\alpha}}, \quad (5)$$

Proof. Setting $xy = u$, by Lemma 1, we obtain

$$\begin{aligned} \omega(\alpha, \beta, \lambda, x) &= \frac{1}{\Gamma(1+\alpha)} \int_0^\infty E_\alpha(-\lambda^\alpha(xy)^\alpha) x^{\beta\alpha} y^{(\beta-1)\alpha} (dy)^\alpha \\ &= \frac{1}{\Gamma(1+\alpha)} \int_0^\infty E_\alpha(-\lambda^\alpha u^\alpha) u^{(\beta-1)\alpha} (du)^\alpha \\ &= L_\alpha\{u^{(\beta-1)\alpha}\}(\lambda) \\ &= \frac{\Gamma_\alpha(\beta)}{\lambda^{\beta\alpha}} \quad \square \end{aligned}$$

LEMMA 4. If $p > 1$, $\frac{1}{p} + \frac{1}{q} = 1$, $0 < \alpha \leq 1$, $\beta, \lambda > 0$, and ε small enough, we define two functions $\tilde{f}(x)$, $\tilde{g}(y)$ as follows:

$$\begin{aligned} \tilde{f}(x) &:= \begin{cases} 0, & x \in (0, 1) \\ x^{(\beta-1)\alpha - \frac{\varepsilon\alpha}{p}}, & x \in [1, \infty) \end{cases}, \\ \tilde{g}(y) &:= \begin{cases} 0, & y \in (1, \infty) \\ y^{(\beta-1)\alpha + \frac{\varepsilon\alpha}{q}}, & y \in (0, 1] \end{cases}, \end{aligned}$$

then we have

$$\tilde{J}\varepsilon^\alpha = \left\{ {}_0I_\infty^\alpha(x^{[p(1-\beta)-1]\alpha} \tilde{f}^p(x)) \right\}^{\frac{1}{p}} \left\{ {}_0I_\infty^\alpha(y^{[q(1-\beta)-1]\alpha} \tilde{g}^q(y)) \right\}^{\frac{1}{q}} \varepsilon^\alpha = \frac{1}{\Gamma(1+\alpha)}, \quad (6)$$

$$\tilde{I}\varepsilon^\alpha = \varepsilon^\alpha \cdot {}_0I_\infty^\alpha \left[{}_0I_\infty^\alpha(E_\alpha(-\lambda^\alpha(xy)^\alpha) \tilde{f}(x) \tilde{g}(y)) \right] > \frac{\Gamma_\alpha(\beta)}{\lambda^{\beta\alpha} \Gamma(1+\alpha)} (1 - o(1)) \quad (\varepsilon \rightarrow 0^+). \quad (7)$$

Proof. We easily obtain:

$$\begin{aligned} \tilde{J}\varepsilon^\alpha &= \left\{ {}_0I_\infty^\alpha(x^{[p(1-\beta)-1]\alpha} \tilde{f}^p(x)) \right\}^{\frac{1}{p}} \left\{ {}_0I_\infty^\alpha(y^{[q(1-\beta)-1]\alpha} \tilde{g}^q(y)) \right\}^{\frac{1}{q}} \varepsilon^\alpha \\ &= \left\{ {}_1I_\infty^\alpha(x^{-(1+\varepsilon)\alpha}) \right\}^{\frac{1}{p}} \left\{ {}_0I_1^\alpha(y^{-(1-\varepsilon)\alpha} (dy)^\alpha) \right\}^{\frac{1}{q}} \varepsilon^\alpha \\ &= \frac{1}{\Gamma(1+\alpha)}. \end{aligned}$$

Since $H(u) = u^{(\beta+1)\alpha} E_\alpha(-\lambda^\alpha(xy)^\alpha)$ is continuous in $[x, \infty)$ and $\lim_{u \rightarrow +\infty} H(u) = 0$, therefore exists a constant $M > 0$, satisfying $H(u) \leq M$. In addition, let $u^{-1+\frac{\varepsilon}{q}} = t$, then $u^{(-2+\frac{\varepsilon}{q})\alpha} (du)^\alpha = -\frac{1}{(1-\frac{\varepsilon}{q})^\alpha} (dt)^\alpha$, and we can write

$$\begin{aligned} & {}_1I_\infty^\alpha(x^{-(1+\varepsilon)\alpha}) {}_xI_\infty^\alpha(E_\alpha(-\lambda^\alpha u^\alpha) u^{(\beta-1+\frac{\varepsilon}{q})\alpha}) \\ &= \frac{1}{\Gamma^2(1+\alpha)} \int_1^\infty x^{-(1+\varepsilon)\alpha} (dx)^\alpha \int_x^\infty E_\alpha(-\lambda^\alpha u^\alpha) u^{(\beta-1+\frac{\varepsilon}{q})\alpha} (du)^\alpha \\ &< \frac{M}{\Gamma^2(1+\alpha)} \int_1^\infty x^{-\alpha} (dx)^\alpha \int_x^\infty u^{(-2+\frac{\varepsilon}{q})\alpha} (du)^\alpha \\ &= \frac{-M}{(1-\frac{\varepsilon}{q})^\alpha \Gamma^2(1+\alpha)} \int_1^\infty x^{-\alpha} (dx)^\alpha \int_{x^{-1+\frac{\varepsilon}{q}}}^0 (dt)^\alpha \\ &= \frac{M}{(1-\frac{\varepsilon}{q})^\alpha \Gamma^2(1+\alpha)} \int_1^\infty x^{-(2-\frac{\varepsilon}{q})\alpha} (dx)^\alpha \\ &= \frac{-M}{(1-\frac{\varepsilon}{q})^{2\alpha} \Gamma^2(1+\alpha)} \int_1^0 (ds)^\alpha \\ &= \frac{M}{(1-\frac{\varepsilon}{q})^{2\alpha} \Gamma^2(1+\alpha)} \end{aligned}$$

Further, setting $xy = u$, and by Lemma 1, we have

$$\begin{aligned} \tilde{I}\varepsilon^\alpha &= \varepsilon^\alpha \cdot {}_0I_\infty^\alpha \left[{}_0I_\infty^\alpha(E_\alpha(-\lambda^\alpha x^\alpha y^\alpha) \tilde{f}(x) \tilde{g}(y)) \right] \\ &= \frac{\varepsilon^\alpha}{\Gamma(1+\alpha)} \int_1^\infty x^{(\beta-1)\alpha - \frac{\varepsilon\alpha}{p}} (dx)^\alpha \left[\frac{1}{\Gamma(1+\alpha)} \int_0^1 E_\alpha(-\lambda^\alpha(xy)^\alpha) y^{(\beta-1)\alpha + \frac{\varepsilon\alpha}{q}} (dy)^\alpha \right] \\ &= \frac{\varepsilon^\alpha}{\Gamma(1+\alpha)} \int_1^\infty x^{-(1+\varepsilon)\alpha} (dx)^\alpha \left[\frac{1}{\Gamma(1+\alpha)} \int_0^x E_\alpha(-\lambda^\alpha u^\alpha) u^{(\beta-1+\frac{\varepsilon}{q})\alpha} (du)^\alpha \right] \\ &= \frac{1}{\Gamma^2(1+\alpha)} \int_0^\infty E_\alpha(-\lambda^\alpha u^\alpha) u^{(\beta-1+\frac{\varepsilon}{q})\alpha} (du)^\alpha \\ &\quad - \varepsilon^\alpha \cdot {}_1I_\infty^\alpha(x^{-(1+\varepsilon)\alpha}) {}_xI_\infty^\alpha(E_\alpha(-\lambda^\alpha u^\alpha) u^{(\beta-1+\frac{\varepsilon}{q})\alpha}) \\ &= \frac{\Gamma_\alpha(\beta + \frac{\varepsilon}{q})}{\lambda^{(\beta+\frac{\varepsilon}{q})\alpha} \Gamma^2(1+\alpha)} - \varepsilon^\alpha \cdot {}_1I_\infty^\alpha(x^{-(1+\varepsilon)\alpha}) {}_xI_\infty^\alpha(E_\alpha(-\lambda^\alpha u^\alpha) u^{(\beta-1+\frac{\varepsilon}{q})\alpha}) \\ &> \frac{\Gamma_\alpha(\beta + \frac{\varepsilon}{q})}{\lambda^{(\beta+\frac{\varepsilon}{q})\alpha} \Gamma(1+\alpha)} - \frac{M\varepsilon^\alpha}{(1-\frac{\varepsilon}{q})^{2\alpha} \Gamma^2(1+\alpha)} \\ &= \frac{\Gamma_\alpha(\beta)}{\lambda^{\beta\alpha} \Gamma(1+\alpha)} (1 - o(1)) \quad (\varepsilon \rightarrow 0^+). \quad \square \end{aligned}$$

3. Main results

For convenience, we use notation of the double fractional integral as (see [16]):

$${}_0I_\infty^\alpha [{}_0I_\infty^\alpha F(x,y)] = \frac{1}{\Gamma^2(1+\alpha)} \int_0^\infty \int_0^\infty F(x,y)(dx)^\alpha (dy)^\alpha.$$

THEOREM 1. *If $p > 1$, $\frac{1}{p} + \frac{1}{q} = 1$, $0 < \alpha \leq 1$, $\beta, \lambda > 0$, $f, g(\geq 0) \in C_\alpha(0, \infty)$, $0 < {}_0I_\infty^\alpha (x^{[p(1-\beta)-1]\alpha} f^p(x)) < \infty$, $0 < {}_0I_\infty^\alpha (y^{[q(1-\beta)-1]\alpha} g^q(y)) < \infty$, then we have*

$${}_0I_\infty^\alpha [{}_0I_\infty^\alpha E_\alpha(-\lambda^\alpha(xy)^\alpha) f(x)g(y)] < \frac{\Gamma_\alpha(\beta)}{\lambda^{\beta\alpha}} \left\{ {}_0I_\infty^\alpha (x^{[p(1-\beta)-1]\alpha} f^p(x)) \right\}^{\frac{1}{p}} \left\{ {}_0I_\infty^\alpha (y^{[q(1-\beta)-1]\alpha} g^q(y)) \right\}^{\frac{1}{q}}, \tag{8}$$

where the constant factor $\frac{\Gamma_\alpha(\beta)}{\lambda^{\beta\alpha}}$ is the best possible.

Proof. By Hölder weighted inequality on fractal set and Lemma 3, we obtain

$$\begin{aligned} & {}_0I_\infty^\alpha [{}_0I_\infty^\alpha E_\alpha(-\lambda^\alpha(xy)^\alpha) f(x)g(y)] \\ &= {}_0I_\infty^\alpha [{}_0I_\infty^\alpha E_\alpha(-\lambda^\alpha(xy)^\alpha) \left[\frac{y^{\frac{(\beta-1)\alpha}{p}}}{x^{\frac{(\beta-1)\alpha}{q}}} \frac{x^{\frac{(\beta-1)\alpha}{q}}}{y^{\frac{(\beta-1)\alpha}{p}}} \right] f(x)g(y)] \\ &\leq \left\{ {}_0I_\infty^\alpha [{}_0I_\infty^\alpha E_\alpha(-\lambda^\alpha(xy)^\alpha) f^p(x) \frac{y^{(\beta-1)\alpha}}{x^{(p-1)(\beta-1)\alpha}}] \right\}^{\frac{1}{p}} \\ &\quad \times \left\{ {}_0I_\infty^\alpha [{}_0I_\infty^\alpha E_\alpha(-\lambda^\alpha(xy)^\alpha) g^q(y) \frac{x^{(\beta-1)\alpha}}{y^{(q-1)(\beta-1)\alpha}}] \right\}^{\frac{1}{q}} \\ &= \left\{ {}_0I_\infty^\alpha [\omega(\alpha, \beta, \lambda, x) x^{p(1-\beta)-1\alpha} f^p(x)] \right\}^{\frac{1}{p}} \left\{ {}_0I_\infty^\alpha [\omega(\alpha, \beta, \lambda, y) y^{q(1-\beta)-1\alpha} g^q(y)] \right\}^{\frac{1}{q}} \\ &= \frac{\Gamma_\alpha(\beta)}{\lambda^{\beta\alpha}} \left\{ {}_0I_\infty^\alpha (x^{[p(1-\beta)-1]\alpha} f^p(x)) \right\}^{\frac{1}{p}} \left\{ {}_0I_\infty^\alpha (y^{[q(1-\beta)-1]\alpha} g^q(y)) \right\}^{\frac{1}{q}} \end{aligned} \tag{9}$$

If (9) keeps the form of equality for some $y \in (0, \infty)$, then, there exist constants A and B , such that they are not all zero, and

$$A \frac{y^{(\beta-1)\alpha}}{x^{(p-1)(\beta-1)\alpha}} f^p(x) = B \frac{x^{(\beta-1)\alpha}}{y^{(q-1)(\beta-1)\alpha}} g^q(y) \quad \text{a.e. in } (0, \infty) \times (0, \infty).$$

Assuming that $A \neq 0$, then there exist constant $C \neq 0$, such that $x^{[p(1-\beta)-1]\alpha} f^p(x) = \frac{C}{Ax^\alpha}$ a.e. in $x \in (0, \infty)$. We find that $\frac{1}{\Gamma(1+\alpha)} \int_0^\infty \frac{C}{Ax^\alpha} (dx)^\alpha = \frac{C}{A} Ln_\alpha x^\alpha|_0^\infty$ is diffuse, which contradicts the fact that $0 < {}_0I_\infty^\alpha (x^{[p(1-\beta)-1]\alpha} f^p(x)) < \infty$. Thus (9) takes the form of strict inequality. So we obtain (8).

If the constant factor $\frac{\Gamma_\alpha(\beta)}{\lambda^{\beta\alpha}}$ appearing on the right hand sides of (8) is not the best possible, then exists a positive constant $K < \frac{\Gamma_\alpha(\beta)}{\lambda^{\beta\alpha}}$, such that inequality (8) is still

valid when replacing $\frac{\Gamma_\alpha(\beta)}{\lambda\beta\alpha}$ by K . Then by (6) and (7), we have $\frac{\Gamma_\alpha(\beta)}{\lambda\beta\alpha}(1 - o(1)) < K$. Letting $\varepsilon \rightarrow 0^+$, we get $K \geq \frac{\Gamma_\alpha(\beta)}{\lambda\beta\alpha}$, which contradicts the fact that $K < \frac{\Gamma_\alpha(\beta)}{\lambda\beta\alpha}$. Hence, the constant factor $\frac{\Gamma_\alpha(\beta)}{\lambda\beta\alpha}$ of (8) is the best possible. The theorem is proved. \square

THEOREM 2. *Under the conditions of Theorem 1, we have*

$${}_0I_\infty^\alpha \left\{ y^{\frac{(q(\beta-1)+1)\alpha}{q-1}} [{}_0I_\infty^\alpha E_\alpha(-\lambda^\alpha(xy)^\alpha)f(x)]^p \right\} < \left[\frac{\Gamma_\alpha(\beta)}{\lambda\beta\alpha} \right]^p {}_0I_\infty^\alpha (x^{[p(1-\beta)-1]\alpha} f^p(x)), \tag{10}$$

where the constant factor $\left[\frac{\Gamma_\alpha(\beta)}{\lambda\beta\alpha} \right]^p$ is the best possible, and inequality (10) is equivalent to inequality (8).

Proof. Letting $[f(x)]_n := \min\{n, f(x)\}$. Since $0 < {}_0I_\infty^\alpha (x^{[p(1-\beta)-1]\alpha} f^p(x)) < \infty$, there exists positive integer $n_0 \in \mathbb{N}$, such that $0 < \frac{1}{n} I_n^\alpha (x^{[p(1-\beta)-1]\alpha} f^p(x)) < \infty$ ($n \geq n_0$). Setting $g_n(y) := y^{\frac{(q(\beta-1)+1)\alpha}{q-1}} \left[\frac{1}{n} I_n^\alpha (E_\alpha(-\lambda^\alpha(xy)^\alpha)[f(x)]_n) \right]^{\frac{p}{q}}$ ($\frac{1}{n} < y < n, n \geq n_0$), when $n \geq n_0$, by (8) we find

$$\begin{aligned} 0 &< \frac{1}{n} I_n^\alpha (y^{[q(1-\beta)-1]\alpha} g_n^q(y)) \\ &= \frac{1}{n} I_n^\alpha (y^{[q(1-\beta)-1]\alpha} g_n^{(q-1)}(y) g_n(y)) \\ &= \frac{1}{n} I_n^\alpha \left\{ \frac{1}{n} I_n^\alpha (E_\alpha(-\lambda^\alpha(xy)^\alpha)[f(x)]_n) y^{\frac{(q(\beta-1)+1)\alpha}{q-1}} \left[\frac{1}{n} I_n^\alpha (E_\alpha(-\lambda^\alpha(xy)^\alpha)[f(x)]_n) \right]^{\frac{p}{q}} \right\} \\ &= \frac{1}{n} I_n^\alpha \left\{ \frac{1}{n} I_n^\alpha (E_\alpha(-\lambda^\alpha(xy)^\alpha)[f(x)]_n g_n(y)) \right\} \\ &< \frac{\Gamma_\alpha(\beta)}{\lambda\beta\alpha} \left\{ \frac{1}{n} I_n^\alpha (x^{[p(1-\beta)-1]\alpha} [f(x)]_n^p) \right\}^{\frac{1}{p}} \left\{ \frac{1}{n} I_n^\alpha (y^{[q(1-\beta)-1]\alpha} g_n^q(y)) \right\}^{\frac{1}{q}}. \end{aligned} \tag{11}$$

Moreover, by (11) we have

$$\begin{aligned} 0 &< \frac{1}{n} I_n^\alpha (y^{[q(1-\beta)-1]\alpha} g_n^q(y)) = \frac{1}{n} I_n^\alpha \left\{ y^{\frac{(q(\beta-1)+1)\alpha}{q-1}} \left[\frac{1}{n} I_n^\alpha (E_\alpha(-\lambda^\alpha(xy)^\alpha)[f(x)]_n) \right]^p \right\} \\ &< \left[\frac{\Gamma_\alpha(\beta)}{\lambda\beta\alpha} \right]^p \frac{1}{n} I_n^\alpha (x^{[p(1-\beta)-1]\alpha} [f(x)]_n^p) < \infty. \end{aligned} \tag{12}$$

It follows that $0 < {}_0I_\infty^\alpha (y^{[q(1-\beta)-1]\alpha} g_\infty^q(y)) < \infty$. For $n \rightarrow \infty$, by (8), both (11) and (12) still keep the form of strict inequalities. Hence we have inequality (10).

On the other hand, by Hölder's inequality on fractal set and (10), we have

$$\begin{aligned} &{}_0I_\infty^\alpha [{}_0I_\infty^\alpha E_\alpha(-\lambda^\alpha(xy)^\alpha)f(x)g(y)] \\ &= {}_0I_\infty^\alpha \left\{ \left[y^{\frac{(q(\beta-1)+1)\alpha}{p(q-1)}} {}_0I_\infty^\alpha (E_\alpha(-\lambda^\alpha(xy)^\alpha)f(x)) \right] \left[y^{\frac{(q(1-\beta)-1)\alpha}{p(q-1)}} g(y) \right] \right\} \\ &\leq \left\{ {}_0I_\infty^\alpha \left(y^{\frac{(q(\beta-1)+1)\alpha}{(q-1)}} [{}_0I_\infty^\alpha (E_\alpha(-\lambda^\alpha(xy)^\alpha)f(x))]^p \right) \right\}^{\frac{1}{p}} \left\{ {}_0I_\infty^\alpha (y^{[q(1-\beta)-1]\alpha} g^q(y)) \right\}^{\frac{1}{p}} \\ &< \frac{\Gamma_\alpha(\beta)}{\lambda\beta\alpha} \left\{ {}_0I_\infty^\alpha (x^{[p(1-\beta)-1]\alpha} f^p(x)) \right\}^{\frac{1}{p}} \left\{ {}_0I_\infty^\alpha (y^{[q(1-\beta)-1]\alpha} g^q(y)) \right\}^{\frac{1}{q}}. \end{aligned}$$

The above inequality is (8), thus inequality (10) is equivalent to inequality (8).

If the constant factor $\left[\frac{\Gamma_\alpha(\beta)}{\lambda\beta\alpha}\right]^p$ appearing on the right hand sides of (10) is not best possible, then by (10), we can get a contradiction that the constant factor $\frac{\Gamma_\alpha(\beta)}{\lambda\beta\alpha}$ appearing on the right hand sides of (8) is not the best possible. Thus the constant factor $\left[\frac{\Gamma_\alpha(\beta)}{\lambda\beta\alpha}\right]^p$ in (10) is the best possible. \square

4. Simple applications

Selecting α values and appropriate β, λ values in (8) and (10), by Lemma 3, some Hilbert-type fractional integral inequalities and their equivalent forms are obtained.

EXAMPLE 1. Letting $\alpha = \beta = \lambda = 1$, we obtain (3) and (4).

EXAMPLE 2. Letting $\alpha = 0.5, \beta = \lambda = 1, p = q = 2$. If $f, g (> 0) \in C_{0.5}(0, \infty)$, $0 < {}_0I_\infty^{0.5}(x^{-0.5}f^2(x)) < \infty, 0 < {}_0I_\infty^{0.5}(y^{-0.5}g^2(y)) < \infty$, then we have the following equivalence inequalities:

$${}_0I_\infty^{0.5} \left[{}_0I_\infty^{0.5} \left((E_{0.5}(-(xy)^{0.5})f(x)g(y)) \right) \right] < \left\{ {}_0I_\infty^{0.5} (x^{-0.5}f^2(x)) \right\}^{\frac{1}{2}} \left\{ {}_0I_\infty^{0.5} (y^{-0.5}g^2(y)) \right\}^{\frac{1}{2}}, \quad (13)$$

$${}_0I_\infty^{0.5} \left\{ y^{0.5} \left[{}_0I_\infty^{0.5} (E_{0.5}(-(xy)^{0.5})f(x)) \right]^2 \right\} < {}_0I_\infty^{0.5} (x^{-0.5}f^2(x)). \quad (14)$$

REFERENCES

- [1] G. H. HARDY, J. E. LITTLEWOOD, G. PÖLYA, *Inequalities*, Cambridge University Press, Cambridge, 1934.
- [2] D. S. MITRINOVIĆ, J. E. PEČARIĆ, AND A. M. FINK, *Inequalities involving functions and their integrals and derivatives*, Kluwer Academic, Boston, 1991.
- [3] G. H. HARDY, *Note on a theorem of Hilbert concernint series of positive terms*, London Math. Soc. **23** (1925), Records of Proc., 45–46.
- [4] B. C. YANG, *On Hilbert's integral inequality*, J. Math. Anal. Appl. **220** (1998), 778–785.
- [5] M. KRNIĆ, J. PEČARIĆ, *General Hilbert's and Hardy's inequalities*, Mathematical inequalities and applications **8**, 1 (2005), 29–51.
- [6] I. BRNETIC, J. PEČARIĆ, *Generalization of Hilbert's integral inequality*, Mathematical inequalities and applications **7**, 2 (2004), 199–205.
- [7] M. KRNIĆ, M. GAO, J. PEČARIĆ, AND X. GAO, *On the besr constant in Hilbert's inequality*, Mathematical inequalities and applications **8**, 2 (2005), 317–329.
- [8] B. C. YANG, *The norm of operator and Hilbert-type inequalities*, Science press, 2009.
- [9] Q. LIU, D. Z CHEN, *A Hilbert-type integral inequality with a hybrid kernel and its applications*, Colloquium mathematicum **143**, 2 (2016), 193–207.
- [10] M. T. RASSIAS, B. C. YANG, *A Hilbert-type integral inequality in the whole plane related to the hypergeometric function and the beta function*, Journal of mathematical analysis and applications **428** (2015), 1286–1308.
- [11] B. C. YANG, *Hilbert-type integral inequality with non-homogeneous kernel*, Journal of Shanghai University: Natural Science **17**, 5 (2011), 603–605.
- [12] H. MO, X. SUI, D. YU, *Generalized convex functions on fractal sets and two related inequalities*, Abstract and Applied Analysis, Article ID636751 (2014).

- [13] E. SAMET, Z. S. MEHMET, *Generalized pompeiu-type inequalities for local fractional integrals and its applications*, Applied Mathematics and Computation **274** (2016), 282–291.
- [14] W. B. SUN, Q. LIU, *New inequalities of Hermite-Hadamard type for generalised convex function on fractal sets and its applications*, Journal of Zhejiang University: Science Edition **44**, 1 (2017), 47–52.
- [15] X. J. YANG, *Advanced local fractional calculus and its applications*, New York: World Science Publisher, 2012.
- [16] X. J. YANG, *Local fractional functional analysis and its applications*, Progress in Nonlinear Science **4** (2011), 1–225.
- [17] J. GUY, *Laplace's transform of fractional order via the Mittag-Leffler function and modified Riemann-Liouville derivative*, Applied Mathematics Letters **22** (2009), 1659–1664.
- [18] G. S. CHEN, *Generalizations of Hölder's and some related integral inequalities on fractal space*, Math. GM **26** (2011), [arXiv: 1109.5567v1](https://arxiv.org/abs/1109.5567v1).
- [19] G. S. CHEN, H. M. SRIVASTAVA, P. WANG, *Some further generalizations of Hölder's inequality and related results on fractal space*, Abstract and Applied Analysis, Article ID 832802 (2014).
- [20] J. C. KUANG, *Applied inequalities* (3rd ed.), Shandong Science and Technology Press, Jinan, 2004.

(Received May 27, 2017)

Qiong Liu
College of Science
Shao Yang University
Shao Yang, China 422000, P. R. China
e-mail: liuqiongxx13@163.com