

## THE BOUNDEDNESS AND ESSENTIAL NORM OF THE DIFFERENCE OF COMPOSITION OPERATORS FROM WEIGHTED BERGMAN SPACES INTO THE BLOCH SPACE

YANHUA ZHANG AND LIXU ZHANG

(Communicated by S. Stević)

*Abstract.* A new characterization for the boundedness of the difference of composition operators  $C_\varphi - C_\psi$  from weighted Bergman spaces into the Bloch space in terms of the Bloch norm of the quantities  $\varphi^n - \psi^n$ ,  $n \in \mathbb{N}$ , is given, as well as an asymptotic estimate for the essential norm of the operator.

### 1. Introduction

Let  $\mathbb{D}$  be the open unit disk in the complex plane  $\mathbb{C}$  and  $H(\mathbb{D})$  be the space of holomorphic functions in  $\mathbb{D}$ . For  $a, z \in \mathbb{D}$ , let  $\sigma_a(z) = \frac{a-z}{1-\bar{a}z}$  be the Möbius transformation of  $\mathbb{D}$  which interchanges 0 and  $a$ . For  $z, w \in \mathbb{D}$ , the pseudo-hyperbolic distance between  $z$  and  $w$  is given by

$$\rho(z, w) = |\sigma_w(z)| = \left| \frac{z-w}{1-\bar{w}z} \right|.$$

It is well known that  $\rho(z, w) \leq 1$ . We denote  $\rho(\varphi(z), \psi(z))$  by  $\rho(z)$ .

For  $0 < p < \infty$  and  $\alpha > -1$ , the weighted Bergman space  $A_\alpha^p = A_\alpha^p(\mathbb{D})$  consists of all  $f \in H(\mathbb{D})$  such that

$$\|f\|_{A_\alpha^p}^p = (\alpha + 1) \int_{\mathbb{D}} |f(z)|^p (1 - |z|^2)^\alpha dA(z) < \infty,$$

where  $dA$  denote the normalized Lebesgue area measure on  $\mathbb{D}$  such that  $A(\mathbb{D}) = 1$ .

Recall that the classical Bloch space  $\mathcal{B} = \mathcal{B}(\mathbb{D})$  consists of all  $f \in H(\mathbb{D})$  such that

$$\|f\|_{\mathcal{B}} = \sup_{z \in \mathbb{D}} (1 - |z|^2) |f'(z)| < \infty.$$

Then the norm  $\|f\|_{\mathcal{B}} = |f(0)| + \|f\|_{\mathcal{B}}$  makes  $\mathcal{B}$  a Banach space.

*Mathematics subject classification* (2010): 30H30, 47B33.

*Keywords and phrases:* Bloch space, weighted Bergman space, difference of composition operators, boundedness, essential norm.

Let  $\varphi$  be an analytic self-map of the unit disk  $\mathbb{D}$ . Associated with  $\varphi$  is the composition operator  $C_\varphi$  defined by

$$C_\varphi(f)(z) = f(\varphi(z)), \quad z \in \mathbb{D},$$

for  $f \in H(\mathbb{D})$ . We denote by  $S(\mathbb{D})$  the set of all analytic self-maps of  $\mathbb{D}$ . The main subject in the study of composition operators is to describe operator theoretic properties of  $C_\varphi$  in terms of function theoretic properties of  $\varphi$ . We refer to books [3, 34] for the basic theory of composition operators on a wide variety of topics.

Recall that  $L : X \rightarrow Y$  is compact if it maps bounded sets into relatively compact sets and  $X, Y$  are Banach spaces. The essential norm of a continuous linear operator  $L$  is its distance to compact operators, that is,

$$\|L\|_{e, X \rightarrow Y} = \inf \{ \|L - K\|_{X \rightarrow Y} : K \text{ is compact} \},$$

where  $\|\cdot\|_{X \rightarrow Y}$  is the operator norm. Clearly,  $L$  is compact if and only if  $\|L\|_{e, X \rightarrow Y} = 0$ .

There has been a considerable interest in estimating of essential norms of composition, weighted composition and other concrete operators involving the composition ones on spaces of analytic functions (see, e.g., [3, 8, 11, 13, 16, 18, 21, 22, 24, 25, 27, 29, 30, 33]).

It is a simple consequence of the Schwarz-Pick inequality that any composition operator  $C_\varphi$  is bounded on  $\mathcal{B}$ . The compactness and essential norm of  $C_\varphi$  on  $\mathcal{B}$  was studied in [12, 13, 31]. Wulan, Zheng and Zhu proved that  $C_\varphi : \mathcal{B} \rightarrow \mathcal{B}$  is compact if and only if  $\lim_{n \rightarrow \infty} \|\varphi^n\|_{\mathcal{B}} = 0$  in [32]. Soon after that, Zhao [33] obtained an exact value of the essential norm for  $C_\varphi : \mathcal{B} \rightarrow \mathcal{B}$ , i.e.

$$\|C_\varphi\|_{e, \mathcal{B} \rightarrow \mathcal{B}} = \frac{e}{2} \limsup_{n \rightarrow \infty} \|\varphi^n\|_{\mathcal{B}}.$$

In [8], it was proved that  $C_\varphi : A_\alpha^p \rightarrow \mathcal{B}$ ,  $1 \leq p < \infty$ ,  $\alpha > -1$ ,  $\varphi \in S(\mathbb{D})$ , is bounded if and only if  $\sup_{n \geq 1} n^{(\alpha+2)/p} \|\varphi^n\|_{\mathcal{B}} < \infty$  (for earlier characterizations, their extensions and related operators, see, e.g., [9, 10, 19, 20, 23, 26]). Moreover, under the assumption that  $C_\varphi : A_\alpha^p \rightarrow \mathcal{B}$  is bounded, they showed that

$$\|C_\varphi\|_{e, A_\alpha^p \rightarrow \mathcal{B}} \approx \limsup_{n \rightarrow \infty} n^{(\alpha+2)/p} \|\varphi^n\|_{\mathcal{B}}.$$

Many authors have investigated in the last few decades the difference of composition operators on various analytic function spaces in order to study their topological structure. The study of the difference of two composition operators was started in [1, 15]. We refer to [2, 4, 5, 6, 7, 14, 16, 17, 24, 28] and related references therein for more information of the difference of composition operators between different spaces of analytic functions.

Let  $\varphi, \psi \in S(\mathbb{D})$  and  $1 \leq p < \infty$ ,  $\alpha > -1$ . We define

$$\mathcal{D}_\varphi(z) := \frac{(1 - |z|^2)\varphi'(z)}{(1 - |\varphi(z)|^2)^{\frac{2+\alpha+p}{p}}}, \quad \mathcal{D}_\psi(z) := \frac{(1 - |z|^2)\psi'(z)}{(1 - |\psi(z)|^2)^{\frac{2+\alpha+p}{p}}}.$$

In [35], Zhu and Yang studied the boundedness and compactness of the difference of two composition operators from  $A_\alpha^p$  to  $\mathcal{B}$ . For example, they showed that  $C_\varphi - C_\psi : A_\alpha^p \rightarrow \mathcal{B}$  is bounded if and only if  $\sup_{z \in \mathbb{D}} |\mathcal{D}_\varphi(z) - \mathcal{D}_\psi(z)| < \infty$  and

$$\sup_{z \in \mathbb{D}} |\mathcal{D}_\varphi(z)|\rho(z) < \infty \text{ (or } \sup_{z \in \mathbb{D}} |\mathcal{D}_\psi(z)|\rho(z) < \infty).$$

Motivated by the results in [8] and [35], in the present paper, we study the boundedness, compactness and essential norm of the operator  $C_\varphi - C_\psi : A_\alpha^p \rightarrow \mathcal{B}$  by using the sequence  $n^{(\alpha+2)/p} \|\varphi^n - \psi^n\|_{\mathcal{B}}$ ,  $n \in \mathbb{N}$ .

For two quantities  $P$  and  $Q$  which may depend on  $\varphi$  and  $\psi$ , we use the abbreviation  $P \lesssim Q$  whenever there is a positive constant  $c$  (independent of  $\varphi$  and  $\psi$ ) such that  $P \leq cQ$ . We write  $P \approx Q$ , if  $P \lesssim Q \lesssim P$ .

### 2. Boundedness of $C_\varphi - C_\psi : A_\alpha^p \rightarrow \mathcal{B}$

In this section we characterize the boundedness of the operator  $C_\varphi - C_\psi : A_\alpha^p \rightarrow \mathcal{B}$ . In order to prove the main result in this section, we need the following lemmas.

LEMMA 2.1. [9] *Let  $1 \leq p < \infty$ ,  $\alpha > -1$ . If  $f \in A_\alpha^p$ , then*

$$|f(z)| \leq \frac{C\|f\|_{A_\alpha^p}}{(1 - |z|^2)^{\frac{2+\alpha}{p}}} \text{ and } |f'(z)| \leq \frac{C\|f\|_{A_\alpha^p}}{(1 - |z|^2)^{\frac{2+\alpha+p}{p}}}.$$

LEMMA 2.2. [35] *Let  $1 \leq p < \infty$ ,  $\alpha > -1$ . Then for all  $z, w \in \mathbb{D}$ ,*

$$\sup_{\|f\|_{A_\alpha^p} \leq 1} |(1 - |z|^2)^{\frac{2+\alpha+p}{p}} f'(z) - (1 - |w|^2)^{\frac{2+\alpha+p}{p}} f'(w)| \lesssim \rho(z, w).$$

For any  $a \in \mathbb{D}$ , we define the following two families of test functions:

$$f_a(z) = \int_0^z \frac{(1 - |a|^2)^{\frac{2+\alpha+p}{p}}}{(1 - \bar{a}u)^{2\frac{2+\alpha+p}{p}}} du,$$

$$g_a(z) = \int_0^z \frac{(1 - |a|^2)^{\frac{2+\alpha+p}{p}}}{(1 - \bar{a}u)^{2\frac{2+\alpha+p}{p}}} \cdot \sigma_a(u) du, \quad z \in \mathbb{D}.$$

LEMMA 2.3. *Let  $\varphi, \psi \in S(\mathbb{D})$ ,  $1 \leq p < \infty$ ,  $\alpha > -1$ . Then*

(i)

$$\sup_{a \in \mathbb{D}} \|(C_\varphi - C_\psi)f_a\|_{\mathcal{B}} \lesssim \sup_{n \in \mathbb{N}} n^{\frac{2+\alpha}{p}} \|\varphi^n - \psi^n\|_{\mathcal{B}}.$$

(ii)

$$\sup_{a \in \mathbb{D}} \|(C_\varphi - C_\psi)g_a\|_{\mathcal{B}} \lesssim \sup_{n \in \mathbb{N}} n^{\frac{2+\alpha}{p}} \|\varphi^n - \psi^n\|_{\mathcal{B}}.$$

*Proof.* It is easy to see that  $\|g_a\|_{A_\alpha^p} \leq \|f_a\|_{A_\alpha^p} \lesssim 1$ . Moreover,

$$f_a(z) = (1 - |a|^2)^t \sum_{k=0}^\infty \frac{\Gamma(k + 2t)}{(k + 1)! \Gamma(2t)} \bar{a}^k z^{k+1}, \quad z \in \mathbb{D}.$$

Here  $t = \frac{2+\alpha+p}{p}$ . Then by Stirling's formula we get

$$f_a(z) \approx (1 - |a|^2)^t \sum_{k=0}^\infty k^{2t-2} \bar{a}^k z^{k+1}, \quad z \in \mathbb{D}.$$

Therefore,

$$\begin{aligned} \|(C_\varphi - C_\psi)f_a\|_{\mathcal{B}} &\lesssim (1 - |a|^2)^t \sum_{k=1}^\infty k^{2t-2} |a|^k \|\varphi^{k+1} - \psi^{k+1}\|_{\mathcal{B}} \\ &\lesssim (1 - |a|^2)^t \sum_{k=1}^\infty k^{t-1} |a|^k (k + 1)^{t-1} \|\varphi^{k+1} - \psi^{k+1}\|_{\mathcal{B}} \\ &\lesssim (1 - |a|^2)^t \sum_{k=1}^\infty k^{t-1} |a|^k \sup_{n \in \mathbb{N}} n^{\frac{2+\alpha}{p}} \|\varphi^n - \psi^n\|_{\mathcal{B}} \\ &\lesssim \sup_{n \in \mathbb{N}} n^{\frac{2+\alpha}{p}} \|\varphi^n - \psi^n\|_{\mathcal{B}}. \end{aligned}$$

Since  $a \in \mathbb{D}$  is arbitrary, we see that (i) holds.

Similarly, we have

$$\begin{aligned} g_a(z) &= (1 - |a|^2)^t \int_0^z \left( \sum_{k=0}^\infty \frac{\Gamma(k + 2t)}{k! \Gamma(2t)} \bar{a}^k u^k \right) \left( a - (1 - |a|^2) \sum_{k=0}^\infty \bar{a}^k u^{k+1} \right) du \\ &= a f_a(z) - (1 - |a|^2)^{t+1} \int_0^z \sum_{k=1}^\infty \left( \sum_{l=0}^{k-1} \frac{\Gamma(l + 2t)}{l! \Gamma(2t)} \right) \bar{a}^{k-1} u^k du. \end{aligned}$$

Hence,

$$\begin{aligned} \|(C_\varphi - C_\psi)g_a\|_{\mathcal{B}} &\lesssim \|(C_\varphi - C_\psi)f_a\|_{\mathcal{B}} + (1 - |a|^2)^{t+1} \sum_{k=1}^\infty k^{2t-1} |a|^{k-1} \|\varphi^{k+1} - \psi^{k+1}\|_{\mathcal{B}} \\ &\lesssim \|(C_\varphi - C_\psi)f_a\|_{\mathcal{B}} + (1 - |a|^2)^{t+1} \sum_{k=1}^\infty k^t |a|^{k-1} \sup_{n \geq 2} n^{t-1} \|\varphi^n - \psi^n\|_{\mathcal{B}} \\ &\lesssim \sup_{n \in \mathbb{N}} n^{\frac{2+\alpha}{p}} \|\varphi^n - \psi^n\|_{\mathcal{B}}, \end{aligned}$$

for any  $a \in \mathbb{D}$ . Therefore (ii) holds. The proof is complete.  $\square$

By modifying the proof of Corollary 4.20 in [34], we can easily obtain the following result.

LEMMA 2.4. *Let  $1 \leq p < \infty$ ,  $\alpha > -1$ . Then*

$$\|z^n\|_{A_\alpha^p} \approx \frac{1}{n^{\frac{2+\alpha}{p}}}.$$

THEOREM 2.1. *Let  $1 \leq p < \infty$  and  $-1 < \alpha < \infty$ . Let  $\varphi$  and  $\psi$  be analytic self-maps of  $\mathbb{D}$ . Then  $C_\varphi - C_\psi : A_\alpha^p \rightarrow \mathcal{B}$  is bounded if and only if*

$$\sup_{n \in \mathbb{N}} n^{\frac{2+\alpha}{p}} \|\varphi^n - \psi^n\|_{\mathcal{B}} < \infty. \tag{1}$$

*Proof.* First, we assume that (1) holds. For any  $z \in \mathbb{D}$ , we have

$$\begin{aligned} & \| (C_\varphi - C_\psi) f_{\varphi(z)} \|_{\mathcal{B}} \\ & \geq |((C_\varphi - C_\psi) f_{\varphi(z)})'(z)| (1 - |z|^2) \\ & = |f'_{\varphi(z)}(\varphi(z))\varphi'(z) - f'_{\varphi(z)}(\psi(z))\psi'(z)| (1 - |z|^2) \\ & = \left| \mathcal{D}_\varphi(z) - \frac{(1 - |\varphi(z)|^2)^{\frac{2+\alpha+p}{p}} (1 - |\psi(z)|^2)^{\frac{2+\alpha+p}{p}}}{(1 - \overline{\varphi(z)}\psi(z))^{\frac{2(2+\alpha+p)}{p}}} \mathcal{D}_\psi(z) \right| \\ & \geq |\mathcal{D}_\varphi(z)| - \frac{(1 - |\varphi(z)|^2)^{\frac{2+\alpha+p}{p}} (1 - |\psi(z)|^2)^{\frac{2+\alpha+p}{p}}}{|1 - \overline{\varphi(z)}\psi(z)|^{\frac{2(2+\alpha+p)}{p}}} |\mathcal{D}_\psi(z)| \end{aligned}$$

and

$$\begin{aligned} \| (C_\varphi - C_\psi) g_{\varphi(z)} \|_{\mathcal{B}} & \geq |g'_{\varphi(z)}(\varphi(z))\varphi'(z) - g'_{\varphi(z)}(\psi(z))\psi'(z)| (1 - |z|^2) \\ & = \frac{(1 - |\varphi(z)|^2)^{\frac{2+\alpha+p}{p}} (1 - |\psi(z)|^2)^{\frac{2+\alpha+p}{p}}}{|1 - \overline{\varphi(z)}\psi(z)|^{\frac{2(2+\alpha+p)}{p}}} |\mathcal{D}_\psi(z)| \rho(z). \end{aligned}$$

Hence

$$\begin{aligned} |\mathcal{D}_\varphi(z)| \rho(z) & \leq \| (C_\varphi - C_\psi) f_{\varphi(z)} \|_{\mathcal{B}} \rho(z) + \| (C_\varphi - C_\psi) g_{\varphi(z)} \|_{\mathcal{B}} \\ & \leq \| (C_\varphi - C_\psi) f_{\varphi(z)} \|_{\mathcal{B}} + \| (C_\varphi - C_\psi) g_{\varphi(z)} \|_{\mathcal{B}}, \end{aligned}$$

and consequently

$$\begin{aligned} \sup_{z \in \mathbb{D}} |\mathcal{D}_\varphi(z)| \rho(z) & \leq \sup_{z \in \mathbb{D}} \| (C_\varphi - C_\psi) f_{\varphi(z)} \|_{\mathcal{B}} + \sup_{z \in \mathbb{D}} \| (C_\varphi - C_\psi) g_{\varphi(z)} \|_{\mathcal{B}} \\ & \leq \sup_{a \in \mathbb{D}} \| (C_\varphi - C_\psi) f_a \|_{\mathcal{B}} + \sup_{a \in \mathbb{D}} \| (C_\varphi - C_\psi) g_a \|_{\mathcal{B}}. \end{aligned} \tag{2}$$

Similarly,

$$|\mathcal{D}_\psi(z)| \rho(z) \leq \| (C_\varphi - C_\psi) f_{\psi(z)} \|_{\mathcal{B}} + \| (C_\varphi - C_\psi) g_{\psi(z)} \|_{\mathcal{B}}. \tag{3}$$

By Lemma 2.2, we have

$$\begin{aligned}
 & \| (C_\varphi - C_\psi) f_{\varphi(z)} \|_{\mathcal{B}} \\
 \geq & \left| \mathcal{D}_\varphi(z) - \frac{(1 - |\varphi(z)|^2)^{\frac{2+\alpha+p}{p}} (1 - |\psi(z)|^2)^{\frac{2+\alpha+p}{p}}}{(1 - \overline{\varphi(z)}\psi(z))^{\frac{2(2+\alpha+p)}{p}}} \mathcal{D}_\psi(z) \right| \\
 \geq & \left| \mathcal{D}_\varphi(z) - \mathcal{D}_\psi(z) \right| - \left| 1 - \frac{(1 - |\varphi(z)|^2)^{\frac{2+\alpha+p}{p}} (1 - |\psi(z)|^2)^{\frac{2+\alpha+p}{p}}}{(1 - \overline{\varphi(z)}\psi(z))^{\frac{2(2+\alpha+p)}{p}}} \right| |\mathcal{D}_\psi(z)| \\
 \geq & |\mathcal{D}_\varphi(z) - \mathcal{D}_\psi(z)| \\
 & - \left| (1 - |\varphi(z)|^2)^{\frac{2+\alpha+p}{p}} f'_{\varphi(z)}(\varphi(z)) - (1 - |\psi(z)|^2)^{\frac{2+\alpha+p}{p}} f'_{\varphi(z)}(\psi(z)) \right| |\mathcal{D}_\psi(z)| \\
 \gtrsim & |\mathcal{D}_\varphi(z) - \mathcal{D}_\psi(z)| - |\mathcal{D}_\psi(z)| \rho(z).
 \end{aligned}$$

Thus, by (3) we get

$$\begin{aligned}
 & |\mathcal{D}_\varphi(z) - \mathcal{D}_\psi(z)| \\
 \lesssim & \| (C_\varphi - C_\psi) f_{\varphi(z)} \|_{\mathcal{B}} + |\mathcal{D}_\psi(z)| \rho(z) \\
 \lesssim & \| (C_\varphi - C_\psi) f_{\varphi(z)} \|_{\mathcal{B}} + \| (C_\varphi - C_\psi) f_{\psi(z)} \|_{\mathcal{B}} + \| (C_\varphi - C_\psi) g_{\psi(z)} \|_{\mathcal{B}}.
 \end{aligned}$$

Hence

$$\sup_{z \in \mathbb{D}} |\mathcal{D}_\varphi(z) - \mathcal{D}_\psi(z)| \lesssim \sup_{a \in \mathbb{D}} \| (C_\varphi - C_\psi) f_a \|_{\mathcal{B}} + \sup_{a \in \mathbb{D}} \| (C_\varphi - C_\psi) g_a \|_{\mathcal{B}}. \tag{4}$$

Combining (2), (4), and Lemma 2.3, we have

$$\begin{aligned}
 & \sup_{z \in \mathbb{D}} |\mathcal{D}_\varphi(z)| \rho(z) + \sup_{z \in \mathbb{D}} |\mathcal{D}_\varphi(z) - \mathcal{D}_\psi(z)| \\
 \lesssim & \sup_{a \in \mathbb{D}} \| (C_\varphi - C_\psi) f_a \|_{\mathcal{B}} + \sup_{a \in \mathbb{D}} \| (C_\varphi - C_\psi) g_a \|_{\mathcal{B}} \\
 \lesssim & \sup_{n \in \mathbb{N}} n^{\frac{2+\alpha}{p}} \| \varphi^n - \psi^n \|_{\mathcal{B}} < \infty.
 \end{aligned}$$

By Theorem 1 of [35], we see that  $C_\varphi - C_\psi : A_\alpha^p \rightarrow \mathcal{B}$  is bounded.

Conversely, suppose that  $C_\varphi - C_\psi : A_\alpha^p \rightarrow \mathcal{B}$  is bounded. For any  $n \in \mathbb{N}$ , let  $f_n(z) = z^n / \|z^n\|_{A_\alpha^p}$ . Then  $\|f_n\|_{A_\alpha^p} = 1$ . Thus, by the boundedness of  $C_\varphi - C_\psi$  and Lemma 2.4, we obtain

$$\begin{aligned}
 \infty & > \| C_\varphi - C_\psi \|_{A_\alpha^p \rightarrow \mathcal{B}} \geq \| (C_\varphi - C_\psi) f_n \|_{\mathcal{B}} = \frac{\| \varphi^n - \psi^n \|_{\mathcal{B}}}{\| z^n \|_{A_\alpha^p}} \\
 & \gtrsim n^{\frac{2+\alpha}{p}} \| \varphi^n - \psi^n \|_{\mathcal{B}},
 \end{aligned}$$

which implies that (1) holds. The proof is complete.  $\square$

### 3. Essential norm estimates

An estimate for the essential norm of  $C_\varphi - C_\psi$  from  $A_\alpha^p$  to  $\mathcal{B}$  will be given in this section by using  $n^{\frac{2+\alpha}{p}} \|\varphi^n - \psi^n\|_{\mathcal{B}}$ . We state some auxiliary results firstly.

LEMMA 3.1. [31] *Let  $X, Y$  be two Banach spaces of analytic functions on  $\mathbb{D}$ . Suppose that the following statements hold.*

- (1) *The point evaluation functionals on  $Y$  are continuous.*
- (2) *The closed unit ball of  $X$  is a compact subset of  $X$  in the topology of uniform convergence on compact sets.*
- (3)  *$T : X \rightarrow Y$  is continuous when  $X$  and  $Y$  are given the topology of uniform convergence on compact sets.*

Then,  $T$  is a compact operator if and only if given a bounded sequence  $\{f_n\}$  in  $X$  such that  $f_n \rightarrow 0$  uniformly on compact sets, then the sequence  $\{Tf_n\}$  converges to zero in the norm of  $Y$ .

LEMMA 3.2. *Let  $1 \leq p < \infty$  and  $-1 < \alpha < \infty$ . Let  $\varphi$  and  $\psi$  be analytic self-maps of  $\mathbb{D}$  such that  $C_\varphi - C_\psi : A_\alpha^p \rightarrow \mathcal{B}$  is bounded. Then the following inequalities hold.*

(i)

$$\limsup_{|a| \rightarrow 1} \|(C_\varphi - C_\psi)f_a\|_{\mathcal{B}} \lesssim \limsup_{n \rightarrow \infty} n^{\frac{2+\alpha}{p}} \|\varphi^n - \psi^n\|_{\mathcal{B}}$$

(ii)

$$\limsup_{|a| \rightarrow 1} \|(C_\varphi - C_\psi)g_a\|_{\mathcal{B}} \lesssim \limsup_{n \rightarrow \infty} n^{\frac{2+\alpha}{p}} \|\varphi^n - \psi^n\|_{\mathcal{B}}.$$

*Proof.* Let  $t = \frac{2+\alpha+p}{p}$ . For any  $N \in \mathbb{N}$ , from the proof of Lemma 2.3, we obtain

$$\begin{aligned} & \limsup_{|a| \rightarrow 1} \|(C_\varphi - C_\psi)f_a\|_{\mathcal{B}} \\ & \leq \limsup_{|a| \rightarrow 1} (1 - |a|^{2t}) \sum_{k=0}^N \frac{\Gamma(k+2t)}{(k+1)! \Gamma(2t)} (k+1)^{1-t} |a|^k (k+1)^{\frac{2+\alpha}{p}} \|\varphi^{k+1} - \psi^{k+1}\|_{\mathcal{B}} \\ & \quad + \limsup_{|a| \rightarrow 1} (1 - |a|^{2t}) \sum_{k=N+1}^{\infty} \frac{\Gamma(k+2t)}{(k+1)! \Gamma(2t)} (k+1)^{1-t} |a|^k \\ & \quad \times \sup_{n \geq N+1} (n+1)^{\frac{2+\alpha}{p}} \|\varphi^{n+1} - \psi^{n+1}\|_{\mathcal{B}} \\ & \lesssim \sup_{n \geq N+2} n^{\frac{2+\alpha}{p}} \|\varphi^n - \psi^n\|_{\mathcal{B}}, \end{aligned}$$

which implies that (i) holds.

Similarly, from the proof of Lemma 2.3 and (i), we have

$$\begin{aligned} \limsup_{|a| \rightarrow 1} \|(C_\varphi - C_\psi)g_a\|_{\mathcal{B}} &\lesssim \limsup_{|a| \rightarrow 1} \|(C_\varphi - C_\psi)f_a\|_{\mathcal{B}} + \sup_{n \geq N+2} n^{\frac{2+\alpha}{p}} \|\varphi^n - \psi^n\|_{\mathcal{B}} \\ &\lesssim \sup_{n \geq N+2} n^{\frac{2+\alpha}{p}} \|\varphi^n - \psi^n\|_{\mathcal{B}}. \end{aligned}$$

The proof is complete.  $\square$

**THEOREM 3.1.** *Let  $1 \leq p < \infty$  and  $\alpha > -1$ . Let  $\varphi$  and  $\psi$  be analytic self-maps of  $\mathbb{D}$ . If  $C_\varphi, C_\psi : A_\alpha^p \rightarrow \mathcal{B}$  are bounded, then*

$$\|C_\varphi - C_\psi\|_{e, A_\alpha^p \rightarrow \mathcal{B}} \approx \limsup_{n \rightarrow \infty} n^{\frac{2+\alpha}{p}} \|\varphi^n - \psi^n\|_{\mathcal{B}}.$$

*Proof.* First, we consider the upper estimate. We adopt the method from [17]. For  $h \in (0, 1)$ , let  $K_h f(z) = f_h(z) = f(hz)$ . Then  $K_h : A_\alpha^p \rightarrow A_\alpha^p$  is compact with  $\|K_h\|_{A_\alpha^p \rightarrow A_\alpha^p} \leq 1$ . Let  $\{h_n\} \subset (0, 1)$  be a sequence such that  $h_n \rightarrow 1$  as  $n \rightarrow \infty$ . Since each  $K_{h_n}$  is compact on  $A_\alpha^p$ ,  $C_\varphi - C_\psi$  is bounded from  $A_\alpha^p$  to  $\mathcal{B}$ ,  $(C_\varphi - C_\psi)K_{h_n}$  is also compact from  $A_\alpha^p$  to  $\mathcal{B}$ . Then, we have

$$\begin{aligned} \|C_\varphi - C_\psi\|_{e, A_\alpha^p \rightarrow \mathcal{B}} &\leq \limsup_{n \rightarrow \infty} \|C_\varphi - C_\psi - (C_\varphi - C_\psi)K_{h_n}\| \\ &= \limsup_{n \rightarrow \infty} \|(C_\varphi - C_\psi)(I - K_{h_n})\| \\ &\leq \limsup_{n \rightarrow \infty} \sup_{\|f\|_{A_\alpha^p} \leq 1} \|(C_\varphi - C_\psi)(I - K_{h_n})f\|_{\mathcal{B}}, \end{aligned}$$

which is bounded by

$$\begin{aligned} &\limsup_{n \rightarrow \infty} \sup_{\|f\|_{A_\alpha^p} \leq 1} |(I - K_{h_n})(f(\varphi(0)) - f(\psi(0)))| + \limsup_{n \rightarrow \infty} \sup_{\|f\|_{A_\alpha^p} \leq 1} \sup_{z \in \mathbb{D}} \\ &|((I - K_{h_n})f)'(\varphi(z))\varphi'(z) - ((I - K_{h_n})f)'(\psi(z))\psi'(z)|(1 - |z|^2). \end{aligned} \tag{5}$$

Since  $f - f_{h_n}$  uniformly on compact subsets of  $\mathbb{D}$  as  $n \rightarrow \infty$  and the sets  $\{\varphi(0)\}$  and  $\{\psi(0)\}$  are compact, we have

$$\limsup_{n \rightarrow \infty} \sup_{\|f\|_{A_\alpha^p} \leq 1} |(I - K_{h_n})(f(\varphi(0)) - f(\psi(0)))| = 0. \tag{6}$$

Let

$$P_n = \sup_{\|f\|_{A_\alpha^p} \leq 1} \sup_{z \in \mathbb{D}} |((I - K_{h_n})f)'(\varphi(z))\varphi'(z) - ((I - K_{h_n})f)'(\psi(z))\psi'(z)|(1 - |z|^2),$$

$f \in A_\alpha^p$  with  $\|f\|_{A_\alpha^p} \leq 1$  and fix an arbitrary  $s \in (0, 1)$ . We set

$$\mathcal{Q}_n^f(z) := |((I - K_{h_n})f)'(\varphi(z))\varphi'(z) - ((I - K_{h_n})f)'(\psi(z))\psi'(z)|(1 - |z|^2)$$



and

$$\mathbb{D}_1 := \{z \in \mathbb{D} : |\varphi(z)| > s, |\psi(z)| > s\}, \mathbb{D}_2 := \{z \in \mathbb{D} : |\varphi(z)| > s, |\psi(z)| \leq s\},$$

$$\mathbb{D}_3 := \{z \in \mathbb{D} : |\varphi(z)| \leq s, |\psi(z)| > s\}, \mathbb{D}_4 := \{z \in \mathbb{D} : |\varphi(z)| \leq s, |\psi(z)| \leq s\}.$$

Then

$$\begin{aligned} \limsup_{n \rightarrow \infty} P_n &= \limsup_{n \rightarrow \infty} \sup_{\|f\|_{A_\alpha^p} \leq 1} \sup_{z \in \mathbb{D}} \mathcal{Q}_n^f = \max_{1 \leq i \leq 4} \limsup_{n \rightarrow \infty} \sup_{\|f\|_{A_\alpha^p} \leq 1} \sup_{z \in \mathbb{D}_i} \mathcal{Q}_n^f \\ &= \max_{1 \leq i \leq 4} \{\limsup_{n \rightarrow \infty} P_n^{(i)}\}, \end{aligned}$$

where  $P_n^{(i)} = \sup_{\|f\|_{A_\alpha^p} \leq 1} \sup_{z \in \mathbb{D}_i} \mathcal{Q}_n^f$ . In addition, we have

$$\begin{aligned} \mathcal{Q}_n^f(z) &\leq |((I - K_{h_n})f)'(\psi(z))|(1 - |\psi(z)|^2)^{\frac{2+\alpha+p}{p}} |\mathcal{D}_\varphi(z) - \mathcal{D}_\psi(z)| \\ &\quad + |((I - K_{h_n})f)'(\varphi(z))|(1 - |\varphi(z)|^2)^{\frac{2+\alpha+p}{p}} \\ &\quad - ((I - K_{h_n})f)'(\psi(z))(1 - |\psi(z)|^2)^{\frac{2+\alpha+p}{p}} \|\mathcal{D}_\varphi(z)\| \\ &\lesssim |((I - K_{h_n})f)'(\psi(z))|(1 - |\psi(z)|^2)^{\frac{2+\alpha+p}{p}} |\mathcal{D}_\varphi(z) - \mathcal{D}_\psi(z)| + |\mathcal{D}_\varphi(z)|\rho(z) \end{aligned}$$

and

$$\mathcal{Q}_n^f(z) \lesssim |((I - K_{h_n})f)'(\varphi(z))|(1 - |\varphi(z)|^2)^{\frac{2+\alpha+p}{p}} |\mathcal{D}_\varphi(z) - \mathcal{D}_\psi(z)| + |\mathcal{D}_\psi(z)|\rho(z).$$

Using the fact that

$$\limsup_{n \rightarrow \infty} \|(I - K_{h_n})f\|_{A_\alpha^p} \leq \limsup_{n \rightarrow \infty} \|I - K_{h_n}\|_{A_\alpha^p \rightarrow A_\alpha^p} \|f\|_{A_\alpha^p} \leq 1,$$

we get

$$\begin{aligned} \limsup_{n \rightarrow \infty} P_n^{(1)} &\lesssim \limsup_{n \rightarrow \infty} \sup_{\|f\|_{A_\alpha^p} \leq 1} \sup_{z \in \mathbb{D}_1} \left( |((I - K_{h_n})f)'(\varphi(z))|(1 - |\varphi(z)|^2)^{\frac{2+\alpha+p}{p}} \right. \\ &\quad \left. \times |\mathcal{D}_\varphi(z) - \mathcal{D}_\psi(z)| + |\mathcal{D}_\psi(z)|\rho(z) \right) \\ &\lesssim \limsup_{n \rightarrow \infty} \sup_{\|f\|_{A_\alpha^p} \leq 1} \sup_{\substack{|\varphi(z)| > s \\ |\psi(z)| > s}} \|(I - K_{h_n})f\|_{A_\alpha^p} |\mathcal{D}_\varphi(z) - \mathcal{D}_\psi(z)| \\ &\quad + \sup_{|\psi(z)| > s} |\mathcal{D}_\psi(z)|\rho(z) \\ &\lesssim \sup_{\substack{|\varphi(z)| > s \\ |\psi(z)| > s}} |\mathcal{D}_\varphi(z) - \mathcal{D}_\psi(z)| + \sup_{|\psi(z)| > s} |\mathcal{D}_\psi(z)|\rho(z). \end{aligned}$$

Thus,

$$\limsup_{n \rightarrow \infty} P_n^{(1)} \leq \limsup_{\substack{|\varphi(z)| \rightarrow 1 \\ |\psi(z)| \rightarrow 1}} |\mathcal{D}_\varphi(z) - \mathcal{D}_\psi(z)| + \limsup_{|\psi(z)| \rightarrow 1} |\mathcal{D}_\psi(z)|\rho(z).$$

Since  $C_\varphi$  and  $C_\psi$  are bounded and  $\sup_{z \in \mathbb{D}} |\mathcal{D}_\varphi(z) - \mathcal{D}_\psi(z)| < \infty$ , we obtain

$$\begin{aligned} & \limsup_{n \rightarrow \infty} P_n^{(2)} \\ & \leq \limsup_{n \rightarrow \infty} \sup_{\|f\|_{A_\alpha^p} \leq 1} \sup_{z \in \mathbb{D}_2} \left( |((I - K_{h_n})f)'(\psi(z))|(1 - |\psi(z)|^2)^{\frac{2+\alpha+p}{p}} \right. \\ & \quad \left. \times |\mathcal{D}_\varphi(z) - \mathcal{D}_\psi(z)| + |\mathcal{D}_\varphi(z)|\rho(z) \right) \\ & \lesssim \limsup_{n \rightarrow \infty} \sup_{\|f\|_{A_\alpha^p} \leq 1} \sup_{|\psi(z)| \leq s} |((I - K_{h_n})f)'(\psi(z))|(1 - |\psi(z)|^2)^{\frac{2+\alpha+p}{p}} \\ & \quad \times |\mathcal{D}_\varphi(z) - \mathcal{D}_\psi(z)| + \sup_{|\varphi(z)| > s} |\mathcal{D}_\varphi(z)|\rho(z) \\ & \lesssim \sup_{|\varphi(z)| > s} |\mathcal{D}_\varphi(z)|\rho(z). \end{aligned}$$

Since  $s$  is arbitrary, we have

$$\limsup_{n \rightarrow \infty} P_n^{(2)} \lesssim \limsup_{|\varphi(z)| \rightarrow 1} |\mathcal{D}_\varphi(z)|\rho(z).$$

Similarly, we obtain

$$\limsup_{n \rightarrow \infty} P_n^{(3)} \lesssim \limsup_{|\psi(z)| \rightarrow 1} |\mathcal{D}_\psi(z)|\rho(z).$$

Finally, by Lemma 3.1, we obtain

$$\begin{aligned} & \limsup_{n \rightarrow \infty} P_n^{(4)} = \limsup_{n \rightarrow \infty} \sup_{\|f\|_{A_\alpha^p} \leq 1} \sup_{z \in \mathbb{D}_4} \mathcal{Q}_n^f \\ & \leq \limsup_{n \rightarrow \infty} \sup_{\|f\|_{A_\alpha^p} \leq 1} \sup_{|\varphi(z)| \leq s} |((I - K_{h_n})f)'(\varphi(z))|\phi'(z)|(1 - |z|^2) \\ & \quad + \limsup_{n \rightarrow \infty} \sup_{\|f\|_{A_\alpha^p} \leq 1} \sup_{|\psi(z)| \leq s} |((I - K_{h_n})f)'(\psi(z))|\psi'(z)|(1 - |z|^2) \\ & = 0. \end{aligned}$$

Therefore,

$$\begin{aligned} \limsup_{n \rightarrow \infty} P_n & = \max\{\limsup_{n \rightarrow \infty} P_n^{(1)}, \limsup_{n \rightarrow \infty} P_n^{(2)}, \limsup_{n \rightarrow \infty} P_n^{(3)}, \limsup_{n \rightarrow \infty} P_n^{(4)}\} \\ & \lesssim \limsup_{|\varphi(z)| \rightarrow 1} |\mathcal{D}_\varphi(z)|\rho(z) + \limsup_{|\psi(z)| \rightarrow 1} |\mathcal{D}_\psi(z)|\rho(z) \\ & \quad + \limsup_{\substack{|\varphi(z)| \rightarrow 1 \\ |\psi(z)| \rightarrow 1}} |\mathcal{D}_\varphi(z) - \mathcal{D}_\psi(z)|. \end{aligned} \tag{7}$$

Set

$$\mathcal{A}_{fg} := \limsup_{|a| \rightarrow 1} \|(C_\varphi - C_\psi)fa\|_{\mathcal{B}} + \limsup_{|a| \rightarrow 1} \|(C_\varphi - C_\psi)ga\|_{\mathcal{B}}.$$

From the proof of Theorem 2.1, we obtain

$$\limsup_{|\varphi(z)| \rightarrow 1} |\mathcal{D}_\varphi(z)|\rho(z) \lesssim \mathcal{A}_{fg}, \quad \limsup_{|\psi(z)| \rightarrow 1} |\mathcal{D}_\psi(z)|\rho(z) \lesssim \mathcal{A}_{fg}$$

and

$$\limsup_{\substack{|\varphi(z)| \rightarrow 1 \\ |\psi(z)| \rightarrow 1}} |\mathcal{D}_\varphi(z) - \mathcal{D}_\psi(z)| \lesssim \mathcal{A}_{fg}.$$

By (5), (6), (7) and Lemma 3.2, we have

$$\begin{aligned} & \|C_\varphi - C_\psi\|_{e.A_\alpha^p \rightarrow \mathcal{B}} \\ & \lesssim \limsup_{|\varphi(z)| \rightarrow 1} |\mathcal{D}_\varphi(z)|\rho(z) + \limsup_{|\psi(z)| \rightarrow 1} |\mathcal{D}_\psi(z)|\rho(z) + \limsup_{\substack{|\varphi(z)| \rightarrow 1 \\ |\psi(z)| \rightarrow 1}} |\mathcal{D}_\varphi(z) - \mathcal{D}_\psi(z)| \\ & \lesssim \mathcal{A}_{fg} \lesssim \limsup_{n \rightarrow \infty} n^{\frac{2+\alpha}{p}} \|\varphi^n - \psi^n\|_{\mathcal{B}}, \end{aligned}$$

as desired.

Next, we consider the lower estimate. Let  $n$  be any positive integer. Set  $y_n(z) = z^n / \|z^n\|_{A_\alpha^p}$ . Then  $\|y_n\|_{A_\alpha^p} = 1$  and by Lemma 2.4,  $y_n \rightarrow 0$  uniformly on compact subsets of  $\mathbb{D}$ . Let  $K : A_\alpha^p \rightarrow \mathcal{B}$  be a compact operator. By Lemma 3.1 we have  $\lim_{n \rightarrow \infty} \|Ky_n\|_{\mathcal{B}} = 0$ . Hence,

$$\|C_\varphi - C_\psi - K\| \geq \limsup_{n \rightarrow \infty} \|(C_\varphi - C_\psi - K)y_n\|_{\mathcal{B}} \geq \limsup_{n \rightarrow \infty} \|(C_\varphi - C_\psi)y_n\|_{\mathcal{B}}.$$

By Lemma 2.4, we have

$$\begin{aligned} \limsup_{n \rightarrow \infty} n^{\frac{2+\alpha}{p}} \|\varphi^n - \psi^n\|_{\mathcal{B}} & \lesssim \limsup_{n \rightarrow \infty} \frac{1}{\|z^n\|_{A_\alpha^p}} \|(C_\varphi - C_\psi)z^n\|_{\mathcal{B}} \\ & = \limsup_{n \rightarrow \infty} \|(C_\varphi - C_\psi)y_n\|_{\mathcal{B}} \\ & \leq \|C_\varphi - C_\psi\|_{e.A_\alpha^p \rightarrow \mathcal{B}}. \end{aligned}$$

The proof is complete.  $\square$

From Theorem 3.1, we immediately get the following corollary.

**COROLLARY 3.1.** *Let  $1 \leq p < \infty$  and  $\alpha > -1$ . Let  $\varphi$  and  $\psi$  be analytic self-maps of  $\mathbb{D}$  such that  $C_\varphi, C_\psi$  are bounded from  $A_\alpha^p$  to  $\mathcal{B}$ . Then  $C_\varphi - C_\psi$  is compact from  $A_\alpha^p$  to  $\mathcal{B}$  if and only if*

$$\limsup_{n \rightarrow \infty} n^{\frac{2+\alpha}{p}} \|\varphi^n - \psi^n\|_{\mathcal{B}} = 0.$$

## REFERENCES

- [1] E. BERKSON, *Composition operators isolated in the uniform operator topology*, Proc. Amer. Math. Soc. **81** (1981), 230–232.
- [2] J. BONET, M. LINDSTRÖM AND E. WOLF, *Differences of composition operators between weighted Banach spaces of holomorphic functions*, J. Austral. Math. Soc. **84** (2008), 9–20.
- [3] C. COWEN AND B. MACCLUER, *Composition Operators on Spaces of Analytic Functions*, CRC Press, Boca Raton, FL, 1995.
- [4] T. HOSOKAWA AND S. OHNO, *Differences of composition operators on the Bloch spaces*, J. Operator Theory **57** (2007), 229–242.
- [5] Q. HU, S. LI AND Y. SHI, *A new characterization of differences of weighted composition operators on weighted-type spaces*, Comput. Methods Funct. Theory **17** (2017), 303–318.
- [6] Z. JIANG AND S. STEVIĆ, *Compact differences of weighted composition operators from weighted Bergman spaces to weighted-type spaces*, Appl. Math. Comput. **217** (2010), 3522–3530.
- [7] S. LI, *Differences of generalized composition operators on the Bloch space*, J. Math. Anal. Appl. **394** (2012), 706–711.
- [8] S. LI, R. QIAN AND J. ZHOU, *Essential norm and a new characterization of weighted composition operators from weighted Bergman spaces and Hardy spaces into the Bloch space*, Czechoslovak Math. J. **67** (2017), 629–643.
- [9] S. LI AND S. STEVIĆ, *Weighted composition operators from Bergman-type spaces into Bloch spaces*, Proc. Indian Acad. Sci. Math. Sci. **117** (2007), 371–385.
- [10] S. LI AND S. STEVIĆ, *Products of composition and integral type operators from  $H^\infty$  to the Bloch space*, Complex Var. Elliptic Equ. **53** (5) (2008), 463–474.
- [11] X. LIU AND S. LI, *Norm and essential norm of a weighted composition operator on the Bloch space*, Integr. Equ. Oper. Theory **87** (2017), 309–325.
- [12] K. MADIGAN AND A. MATHESON, *Compact composition operators on the Bloch space*, Trans. Amer. Math. Soc. **347** (1995), 2679–2687.
- [13] A. MONTES-RODRIGUEZ, *The essential norm of a composition operator on Bloch spaces*, Pacific J. Math. **188** (1999), 339–351.
- [14] P. NIEMINEN, *Compact differences of composition operators on Bloch and Lipschitz spaces*, Comput. Method Funct. Theory **7** (2007), 325–344.
- [15] J. SHAPIRO AND C. SUNDBERG, *Isolation amongst the composition operators*, Pacific J. Math. **145** (1990), 117–152.
- [16] Y. SHI AND S. LI, *Essential norm of the differences of composition operators on the Bloch space*, Math. Ineq. Appl. **20** (2017), 543–555.
- [17] Y. SHI AND S. LI, *Differences of composition operators on Bloch type spaces*, Complex Anal. Oper. Theory. **11** (2017), 227–242.
- [18] S. STEVIĆ, *Essential norms of weighted composition operators from the  $\alpha$ -Bloch space to a weighted-type space on the unit ball*, Abstr. Appl. Anal. **2008** (2008), Article ID 279691, 11 pages.
- [19] S. STEVIĆ, *On a new integral-type operator from the weighted Bergman space to the Bloch-type space on the unit ball*, Discrete Dyn. Nat. Soc. **2008** (2008), Article ID 154263, 14 pages.
- [20] S. STEVIĆ, *Integral-type operators from a mixed norm space to a Bloch-type space on the unit ball*, Siberian Math. J. **50** (6) (2009), 1098–1105.
- [21] S. STEVIĆ, *Norm and essential norm of composition followed by differentiation from  $\alpha$ -Bloch spaces to  $H_\mu^\infty$* , Appl. Math. Comput. **207** (2009), 225–229.
- [22] S. STEVIĆ, *On a new integral-type operator from the Bloch space to Bloch-type spaces on the unit ball*, J. Math. Anal. Appl. **354** (2009), 426–434.
- [23] S. STEVIĆ, *On an integral-type operator from logarithmic Bloch-type and mixed-norm spaces to Bloch-type spaces*, Nonlinear Anal. TMA **71** (2009), 6323–6342.
- [24] S. STEVIĆ, *Essential norm of differences of weighted composition operators between weighted-type spaces on the unit ball*, Appl. Math. Comput. **217** (2010), 1811–1824.
- [25] S. STEVIĆ, *Norm and essential norm of an integral-type operator from the Dirichlet space to the Bloch-type space on the unit ball*, Abstr. Appl. Anal. **2010** (2010), Article ID 134969, 9 pages.
- [26] S. STEVIĆ, *Weighted differentiation composition operators from the mixed-norm space to the  $n$ th weighted-type space on the unit disk*, Abstr. Appl. Anal. **2010** (2010), Article ID 246287, 15 pages.

- [27] S. STEVIĆ, *Essential norm of some extensions of the generalized composition operators between  $k$ -th weighted-type spaces*, J. Inequal. Appl. **2017** (2017), Article No. 220, 13 pages.
- [28] S. STEVIĆ AND Z. JIANG, *Compactness of the differences of weighted composition operators from weighted Bergman spaces to weighted-type spaces on the unit ball*, Taiwanese J. Math. **15** (2011), 2647–2665.
- [29] S. STEVIĆ AND A. SHARMA, *Essential norm of composition operators between weighted Hardy spaces*, Appl. Math. Comput. **217** (2011), 6192–6197.
- [30] S. STEVIĆ, A. SHARMA AND A. BHAT, *Essential norm of products of multiplication composition and differentiation operators on weighted Bergman spaces*, Appl. Math. Comput. **218** (2011), 2386–2397.
- [31] M. TJANI, *Compact composition operators on some Möbius invariant Banach space*, PhD dissertation, Michigan State University, 1996.
- [32] H. WULAN, D. ZHENG AND K. ZHU, *Compact composition operators on BMOA and the Bloch space*, Proc. Amer. Math. Soc. **137** (2009), 3861–3868.
- [33] R. ZHAO, *Essential norms of composition operators between Bloch type spaces*, Proc. Amer. Math. Soc. **138** (2010), 2537–2546.
- [34] K. ZHU, *Operator Theory in Function Spaces*, American Mathematical Society, Providence, RI, 2007.
- [35] X. ZHU AND W. YANG, *Difference of composition operators from weighted Bergman spaces to the Bloch space*, Filomat **28** (2014), 1935–1941.

(Received January 9, 2018)

Yanhua Zhang  
Department of Mathematics  
Qufu Normal University  
273165, Qufu, ShanDong, China  
e-mail: qfuzhangyanhua@163.com

Lixu Zhang  
Faculty of Information Technology  
Macau University of Science and Technology  
Avenida Wai Long, Taipa, Macau  
e-mail: zhanglixu46@163.com