

## FRACTIONAL DIFFERENTIAL OPERATORS IN VECTOR-VALUED SPACES AND APPLICATIONS

VELI SHAKHMUROV

(Communicated by M. Bohner)

*Abstract.* Fractional differential operator equations with parameter are studied. Uniform  $L_p$ -separability properties and sharp resolvent estimates are obtained for elliptic equations in terms of fractional derivatives. Moreover, maximal regularity properties of the fractional abstract parabolic equation are established. Particularly, it is proven that the operators generated by these equations are positive and also are generators of analytic semigroups. As an application, the anisotropic parameter dependent fractional differential equations and the system of fractional differential equations are studied.

### 1. Introduction, notations and background

In the last years, the maximal regularity properties of boundary value problems (BVPs) for differential-operator equations (DOEs) have found many applications in PDE and pseudo DE with applications in physics (see [1, 5, 8-15] and the references therein). Pseudo-differential equations (PsDE) were treated e.g. in [17-18]. DOEs have found many applications in fractional differential equations (FDEs), pseudo-differential equations (PsDE) and PDEs (see e.g. [1–3], [5], [11], [12], [16–19], [24]). The regularity properties of PsDE have been studied extensively by many researchers (see e.g. [6, 10], [21-22] and the references therein). The boundedness of PsDEs in Sobolev spaces have been treated e.g. in [10], [14], [22]. Moreover, the smoothness of PsDE with bounded operator coefficients have been explored e.g. in [8], [15]. In contrast to [8], [15] the FDE considered here, contain unbounded operators and parameters. In particular, the main objective of the present paper is to discuss the uniform  $L_p(\mathbb{R}^n; E)$ -maximal regularity of the elliptic fractional differential operator equation (FDOE) with parameters

$$P_\varepsilon(D)u + Au + \sum_{|\alpha| < m} \varepsilon(\alpha) A_\alpha(x) D^\alpha u + \lambda u = f(x), \quad x \in \mathbb{R}^n, \quad (1.1)$$

where  $P_\varepsilon(D)$  is a fractional differential operator,  $A, A_\alpha(x)$  are linear operators in a Banach space  $E$  for  $\alpha_i \in [0, \infty)$  and  $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_n)$ . Here,  $D^\alpha = D_1^{\alpha_1} D_2^{\alpha_2} \dots D_n^{\alpha_n}$

*Mathematics subject classification* (2010): 47GXX, 35JXX, 47FXX, 47DXX, 43AXX — Please, give more precise subject classification, Thanks!.

*Keywords and phrases:* Fractional-differential equations, Sobolev-Lions spaces, differential-operator equations, maximal  $L_p$  regularity, abstract parabolic equations, operator-valued multipliers.

are the Liouville derivatives,  $m$  is a positive number,  $\varepsilon_k$  are positive,  $\lambda$  is a complex parameter,  $\varepsilon = (\varepsilon_1, \varepsilon_2, \dots, \varepsilon_n)$  and  $\varepsilon(\alpha) = \prod_{k=1}^n \varepsilon_k^{\frac{\alpha_k}{m}}$ .

Here,  $L_p(\Omega; E)$  denotes the space of strongly measurable  $E$ -valued functions that are defined on the measurable subset  $\Omega \subset \mathbb{R}^n$  with the norm given by

$$\|f\|_{L_p(\Omega; E)} = \left( \int_{\Omega} \|f(x)\|_E^p dx \right)^{\frac{1}{p}}, \quad 1 \leq p < \infty,$$

$$\|f\|_{L^\infty} = \operatorname{ess\,sup}_{x \in \Omega} \|f(x)\|_E.$$

We prove that problem (1.1) has a maximal regular unique solution and the following uniform coercive estimate holds

$$\sum_{|\alpha| \leq m} \varepsilon(\alpha) |\lambda|^{1 - \frac{|\alpha|}{m}} \|D^\alpha u\|_{L_p(\mathbb{R}^n; E)} + \|Au\|_{L_p(\mathbb{R}^n; E)} \leq C \|f\|_{L_p(\mathbb{R}^n; E)} \tag{1.2}$$

for  $f \in L_p(\mathbb{R}^n; E)$ ,  $\lambda \in S_\varphi$ , where  $S_\varphi$  is a set of complex numbers that is related with the spectrum of the operator  $A$ . The estimate (1.2) implies that the operator  $O_\varepsilon$  generated by (1.1) has a bounded inverse from  $L_p(\mathbb{R}^n; E)$  into the space  $H_p^m(\mathbb{R}^n; E(A), E)$ , which will be defined subsequently. Particularly, from the estimate (1.2), we obtain that the operator  $O_\varepsilon$  is uniformly positive in  $L_p(\mathbb{R}^n; E)$ . By using this property we prove the uniform well posedness of the Cauchy problem for the parabolic FDOE with parameter

$$\frac{\partial u}{\partial t} + P_\varepsilon(D)u + Au = f(t, x), \quad u(0, x) = 0, \tag{1.3}$$

in  $E$ -valued mixed spaces  $L_{\mathbf{p}}$  for  $\mathbf{p} = (p, p_1)$ . In other words, we show that problem (1.3) has a unique solution  $u \in W_{\mathbf{p}}^{1,m}(\mathbb{R}_+^{n+1}; E(A), E)$  for  $f \in L_{\mathbf{p}}(\mathbb{R}_+^{n+1}; E)$  satisfying the following uniform coercive estimate

$$\left\| \frac{\partial u}{\partial t} \right\|_{L_{\mathbf{p}}(\mathbb{R}_+^{n+1}; E)} + \|P_\varepsilon(D)u\|_{L_{\mathbf{p}}(\mathbb{R}_+^{n+1}; E)} + \|Au\|_{L_{\mathbf{p}}(\mathbb{R}_+^{n+1}; E)} \leq M \|f\|_{L_{\mathbf{p}}(\mathbb{R}_+^{n+1}; E)}. \tag{1.4}$$

Note that, the constants  $C, M$  in (1.2) and (1.4) are independent of parameters. As an application, in this paper the following are established: (a) maximal regularity properties of the anisotropic elliptic FDOE in mixed  $L_{\mathbf{p}}$ ,  $\mathbf{p} = (p_1, p)$  spaces; (b) coercive properties of the system of FDOEs of infinite many order in  $L_p$  spaces.

The Banach space  $E$  is called an UMD-space if the Hilbert operator

$$(Hf)(x) = \lim_{\varepsilon \rightarrow 0} \int_{|x-y| > \varepsilon} \frac{f(y)}{x-y} dy$$

is bounded in  $L^p(\mathbb{R}; E)$ ,  $p \in (1, \infty)$  (see. e.g [4]). UMD spaces include e.g.  $L^p$ ,  $l_p$  and Lorentz spaces  $L_{pq}$  for  $p, q \in (1, \infty)$ .

Let  $\mathbb{C}$  denote the set of complex numbers and

$$S_\varphi = \{\lambda; \lambda \in \mathbb{C}, |\arg \lambda| \leq \varphi\} \cup \{0\}, 0 \leq \varphi < \pi.$$

A linear operator  $A$  is said to be  $\varphi$ -positive (or positive) in a Banach space  $E$  if  $D(A)$  is dense on  $E$  and

$$\left\| (A + \lambda I)^{-1} \right\|_{B(E)} \leq M(1 + |\lambda|)^{-1}$$

for any  $\lambda \in S_\varphi$ , where  $\varphi \in [0, \pi)$ ,  $I$  is the identity operator in  $E$ ,  $B(E)$  is the space of bounded linear operators in  $E$ . Sometimes  $A + \lambda I$  will be written  $A + \lambda$  and will be denoted by  $A_\lambda$ . It is known [20, §1.15.1] that the powers  $A^\theta$ ,  $\theta \in (-\infty, \infty)$  for a positive operator  $A$  exist. The operator  $A(h)$ ,  $h \in Q \subset \mathbb{C}$  is said to be  $\varphi$ -positive (or positive) in  $E$  uniformly with respect to  $h \in Q$  if  $D(A(h))$  is independent of  $h$ ,  $D(A(h))$  is dense in  $E$  and  $\left\| (A(h) + \lambda)^{-1} \right\| \leq M(1 + |\lambda|)^{-1}$  for all  $\lambda \in S_\varphi$ ,  $0 \leq \varphi < \pi$ , where  $M$  does not depend on  $h$  and  $\lambda$ . Let  $E(A^\theta)$  denote the space  $D(A^\theta)$  with the norm

$$\|u\|_{E(A^\theta)} = \left( \|u\|^p + \|A^\theta u\|^p \right)^{\frac{1}{p}}, 1 \leq p < \infty, 0 < \theta < \infty.$$

A set  $W \subset B(E_1, E_2)$  is called  $R$ -bounded (see e.g. [23]) if there is a constant  $C > 0$  such that for all  $T_1, T_2, \dots, T_m \in W$  and  $u_1, u_2, \dots, u_m \in E_1, m \in \mathbb{N}$ ,

$$\int_0^1 \left\| \sum_{j=1}^m r_j(y) T_j u_j \right\|_{E_2} dy \leq C \int_0^1 \left\| \sum_{j=1}^m r_j(y) u_j \right\|_{E_1} dy,$$

where  $\{r_j\}$  is an arbitrary sequence of independent symmetric  $\{-1, 1\}$ -valued random variables on  $[0, 1]$ . The smallest  $C$  for which the above estimate holds is called an  $R$ -bound of the collection  $W$  and is denoted by  $R(W)$ . A set of operators  $G_h \subset B(E_1, E_2)$  depending on parameter  $h \in Q \subset \mathbb{C}$  is called uniformly  $R$ -bounded in  $h$  if there is a constant  $C$  independent of  $h \in Q$  such that

$$\int_\Omega \left\| \sum_{j=1}^m r_j(y) T_j(h) u_j \right\|_{E_2} dy \leq C \int_\Omega \left\| \sum_{j=1}^m r_j(y) u_j \right\|_{E_1} dy$$

for all  $T_1(h), T_2(h), \dots, T_m(h) \in G_h$  and  $u_1, u_2, \dots, u_m \in E_1, m \in \mathbb{N}$ . It implies that  $\sup_{h \in Q} R(G_h) \leq C$ .

The operator  $A$  is said to be  $R$ -positive in a Banach space  $E$  if the set  $\left\{ \lambda (A + \lambda)^{-1} : \lambda \in S_\varphi \right\}$  is  $R$ -bounded. A positive operator  $A(h)$  is said to be uniformly  $R$ -positive in a Banach space  $E$  if there exists  $\varphi \in [0, \pi)$  such that the set

$$\left\{ \lambda (A(h) + \lambda)^{-1} : \lambda \in S_\varphi \right\}$$

is uniformly  $R$ -bounded. Let  $S(\mathbb{R}^n; E)$  denote the  $E$ -valued Schwartz class, i.e., the space of all  $E$ -valued rapidly decreasing smooth functions on  $\mathbb{R}^n$  equipped with its usual topology generated by seminorms. For  $E = \mathbb{C}$  this space will be denoted by  $S = S(\mathbb{R}^n)$ . Here,  $S'(E) = S'(\mathbb{R}^n; E)$  denotes the space of linear continuous mappings from  $S$  into  $E$  and is called  $E$ -valued Schwartz distributions. For any  $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_n)$ ,  $\alpha_i \in [0, \infty)$  the function  $(i\xi)^\alpha$  will be defined as:

$$(i\xi)^\alpha = \begin{cases} (i\xi_1)^{\alpha_1}, \dots, (i\xi_n)^{\alpha_n}, \xi_1 \xi_2, \dots, \xi_n \neq 0 \\ 0, \xi_1, \xi_2, \dots, \xi_n = 0, \end{cases}$$

where

$$(i\xi_k)^{\alpha_k} = \exp[\alpha_k (\ln |\xi_k| + i\pi \operatorname{sgn} \xi_k / 2)], k = 1, 2, \dots, n.$$

The Liouville derivatives  $D^\alpha u$  of an  $E$ -valued function  $u$  are defined similarly to the case of scalar functions [13].

$C(\Omega; E)$  and  $C^{(m)}(\Omega; E)$  will denote the spaces of  $E$ -valued bounded uniformly strongly continuous and  $m$  times continuously differentiable functions on  $\Omega$ , respectively. Let  $F$  and  $F^{-1}$  denote the Fourier and inverse Fourier transforms defined as

$$Fu = (2\pi)^{-\frac{n}{2}} \int_{\mathbb{R}^n} [\exp(x, \xi)] u(x) dx, F^{-1}u = (2\pi)^{-\frac{n}{2}} \int_{\mathbb{R}^n} [\exp - (x, \xi)] u(\xi) d\xi,$$

where

$$x = (x_1, x_2, \dots, x_n), \xi = (\xi_1, \xi_2, \dots, \xi_n) \in \mathbb{R}^n, (x, \xi) = \sum_{k=1}^n x_k \xi_k.$$

Through this section, the Fourier transformation of a function  $u$  will be denoted by  $\hat{u}$ . It is known that

$$F(D_x^\alpha u) = (i\xi_1)^{\alpha_1}, \dots, (i\xi_n)^{\alpha_n} \hat{u}, D_\xi^\alpha (F(u)) = F [(-ix_n)^{\alpha_1}, \dots, (-ix_1)^{\alpha_n} u]$$

for all  $u \in S'(\mathbb{R}^n; E)$ . Let  $E_1$  and  $E_2$  be two Banach spaces.  $B(E_1, E_2)$  denotes the space of bounded linear operators from  $E_1$  to  $E_2$ . A function  $\Psi \in C(\mathbb{R}^n; B(E_1, E_2))$  is called a Fourier multiplier from  $L_p(\mathbb{R}^n; E_1)$  to  $L_p(\mathbb{R}^n; E_2)$  if the map

$$u \rightarrow \Lambda u = F^{-1} \Psi(\xi) Fu, u \in S(\mathbb{R}^n; E_1)$$

is well defined and extends to a bounded linear operator

$$\Lambda : L_p(\mathbb{R}^n; E_1) \rightarrow L_p(\mathbb{R}^n; E_2).$$

The set of all Fourier multipliers from  $L_p(\mathbb{R}^n; E_1)$  to  $L_p(\mathbb{R}^n; E_2)$  will be denoted by  $M_p^p(E_1, E_2)$ . For  $E_1 = E_2 = E$  it is denoted by  $M_p^p(E)$ . Let  $\Phi_h = \{\Psi_h \in M_p^p(E_1, E_2), h \in Q\}$  denote a collection of multipliers depending on the parameter  $h$ . We say that  $W_h$  is a uniform collection of multipliers if there exists a positive constant  $M$  independent of  $h \in Q$  such that

$$\|F^{-1} \Psi_h Fu\|_{L_p(\mathbb{R}^n; E_2)} \leq M \|u\|_{L_p(\mathbb{R}^n; E_1)}$$

for all  $h \in Q$  and  $u \in S(\mathbb{R}^n; E_1)$ .

Let  $E_0$  and  $E$  be two Banach spaces and  $E_0$  be continuously and densely embedded into  $E$ . Let  $s \in \mathbb{R}$  and  $\xi = (\xi_1, \xi_2, \dots, \xi_n) \in \mathbb{R}^n$ . Consider the following Liouville-Lions space

$$H_p^s(\mathbb{R}^n; E_0, E) = \left\{ u \in S'(\mathbb{R}^n; E_0), F^{-1} \left( 1 + |\xi|^2 \right)^{\frac{s}{2}} Fu \in L_p(\mathbb{R}^n; E), \right. \\ \left. \|u\|_{H_p^s(\mathbb{R}^n; E_0, E)} = \|u\|_{L_p(\mathbb{R}^n; E_0)} + \left\| F^{-1} \left( 1 + |\xi|^2 \right)^{\frac{s}{2}} Fu \right\|_{L_p(\mathbb{R}^n; E)} < \infty \right\}.$$

Let  $\varepsilon = (\varepsilon_1, \varepsilon_2, \dots, \varepsilon_n)$  and  $\varepsilon_k$  be positive parameters. We define the following parameterized norm in  $H_p^s(\mathbb{R}^n; E_0, E)$ ,

$$\|u\|_{H_p^{s, \varepsilon}(\mathbb{R}^n; E_0, E)} = \|u\|_{L_p(\mathbb{R}^n; E_0)} + \left\| F^{-1} \left[ 1 + \left( \sum_{k=1}^n \varepsilon_k^{\frac{2}{s}} \xi_k^2 \right)^{\frac{1}{2}} \right]^s Fu \right\|_{L_p(\mathbb{R}^n; E)} < \infty.$$

Sometimes we use one and the same symbol  $C$  without distinction in order to denote positive constants which may differ from each other even in a single context. When we want to specify the dependence of such a constant on a parameter, say  $\alpha$ , we write  $C_\alpha$ .

By using the techniques of [9, Theorem 3.7] and reasoning as in [19, Theorem A<sub>0</sub>], we obtain the following proposition.

PROPOSITION A<sub>0</sub>. *Let  $E_1$  and  $E_2$  be two UMD spaces and*

$$\Psi_h \in C^n(\mathbb{R}^n \setminus \{0\}; B(E_1, E_2)).$$

*Suppose there is a positive constant  $K$  such that*

$$\sup_{h \in Q} R \left( \left\{ |\xi|^{|\beta|} D^\beta \Psi_h(\xi) : \xi \in \mathbb{R}^n \setminus \{0\}, \beta_i \in \{0, 1\} \right\} \right) \leq K,$$

*for*

$$\beta = (\beta_1, \beta_2, \dots, \beta_n), \quad |\beta| = \sum_{k=1}^n \beta_k.$$

*Then  $\Psi_h$  is a uniform collection of multipliers from  $L_p(\mathbb{R}^n; E_1)$  to  $L_p(\mathbb{R}^n; E_2)$  for  $p \in (1, \infty)$ .*

*Proof.* Some steps (Lemma 3.1, Proposition 3.2) of proof [9, Theorem 3.7] trivially work for the parameter dependent case. Other steps (Theorem 3.3, Lemma 3.5) can be easily shown by replacing

$$\left\{ |\xi|^{|\beta|} D^\beta \Psi(\xi) : \xi \in \mathbb{R}^n \setminus \{0\} \right\}$$

with

$$\Sigma_h = \left\{ |\xi|^{|\beta|} D^\beta \Psi_h(\xi) : \xi \in \mathbb{R}^n \setminus \{0\} \right\}$$

and by using the uniform  $R$ -boundedness of the set  $\Sigma_h$ . However, the parameter dependent analog of Proposition 3.4 in [9] is not straightforward. Let  $M_h, M_{h,N} \in L_1^{loc}(\mathbb{R}^n, \mathcal{B}(E_1, E_2))$  be Fourier multipliers from  $L_p(\mathbb{R}^n; E_1)$  to  $L_p(\mathbb{R}^n; E_2)$ . Let  $M_{h,N}$  converge to  $M_h$  in  $L_1^{loc}(\mathbb{R}^n, \mathcal{B}(E_1, E_2))$  and  $T_{h,N} = F^{-1}M_{h,N}F$  be uniformly bounded. Then the operator  $T_h = F^{-1}M_hF$  is uniformly bounded, so we obtain the assertion of Proposition A<sub>0</sub>.

The embedding theorems in vector valued spaces play a key role in the theory of DOEs. For estimating lower order derivatives in terms of interpolation spaces we use following embedding theorems from [17].

**THEOREM A<sub>1</sub>.** *Suppose  $E$  is an UMD space,  $0 < \varepsilon_k \leq \varepsilon_0 < \infty, 1 < p \leq q < \infty$  and  $A$  is an  $R$ -positive operator in  $E$ . Then for  $s \in (0, \infty)$  with  $\varkappa = |\alpha| + n\left(\frac{1}{p} - \frac{1}{q}\right) \leq s, 0 \leq \mu \leq 1 - \varkappa$  the embedding*

$$D^\alpha H_p^s(\mathbb{R}^n; E(A), E) \subset L_q(\mathbb{R}^n; E(A^{1-\varkappa-\mu}))$$

is continuous and there exists a constant  $C_\mu > 0$ , depending only on  $\mu$  such that

$$\varepsilon(\alpha) \|D^\alpha u\|_{L_q(\mathbb{R}^n; E(A^{1-\varkappa-\mu}))} \leq C_\mu \left[ h^\mu \|u\|_{H_{p,\varepsilon}^s(\mathbb{R}^n; E(A), E)} + h^{-(1-\mu)} \|u\|_{L_p(\mathbb{R}^n; E)} \right]$$

for all  $u \in H_p^s(\mathbb{R}^n; E(A), E)$  and  $0 < h \leq h_0 < \infty$ .

### 2. FDOE with parameters in Banach spaces

Consider the principal part of the problem (1.1),

$$(L_\varepsilon + \lambda)u = P_\varepsilon(D)u + Au + \lambda u = f(x), \quad x \in \mathbb{R}^n, \tag{2.1}$$

where  $P_\varepsilon(D)$  is the fractional differential operator defined by

$$P_\varepsilon(D)u = F^{-1}P_\varepsilon(\xi)\hat{u}(\xi) = (2\pi)^{-\frac{n}{2}} \int_{\mathbb{R}^n} e^{i(x,\xi)} P_\varepsilon(\xi)\hat{u}(\xi) d\xi. \tag{2.2}$$

**CONDITION 2.1.** Assume  $P_\varepsilon(\xi) \in S^m$  for some positive number  $m$ , i.e.,

$$\left| D_\xi^\alpha P_\varepsilon(\xi) \right| \leq C_\alpha \left[ 1 + \left( \sum_{k=1}^n \varepsilon_k^{\frac{2}{m-|\alpha|}} \xi_k^2 \right)^{\frac{1}{2}} \right]^{m-|\alpha|}$$

for all  $\xi \in \mathbb{R}^n, \alpha_k > 0, \alpha = (\alpha_1, \alpha_2, \dots, \alpha_n)$  with  $|\alpha| \leq m$  and  $\varepsilon_k \in (0, \varepsilon_0]$ . Suppose  $P_\varepsilon(\xi) \in S_{\varphi_1}$  for all  $\xi \in \mathbb{R}^n, 0 \leq \varphi_1 < \pi$  and there is a constant  $\gamma > 0$  such that

$$|P_\varepsilon(\xi)| \geq \gamma \sum_{k=1}^n \varepsilon_k |\xi_k|^m.$$

Let

$$X = L_p(\mathbb{R}^n; E), Y = H_p^m(\mathbb{R}^n; E(A), E).$$

In this section we prove the following:

**THEOREM 2.1.** *Assume the Condition 2.1 holds. Suppose  $E$  is an UMD space,  $p \in (1, \infty)$  and  $A$  is an  $R$ -positive operator in  $E$  with respect to  $\varphi \in (0, \pi]$ . Then for  $f \in X$ ,  $\lambda \in S_{\varphi_2}$ ,  $0 \leq \varphi_1 < \pi - \varphi_2$  and  $\varphi_1 + \varphi_2 \leq \varphi$  there is a unique solution  $u$  of the equation (2.1) belonging to  $Y$  and the following coercive uniform estimate holds*

$$\sum_{|\alpha| \leq m} \varepsilon(\alpha) |\lambda|^{1 - \frac{|\alpha|}{m}} \|D^\alpha u\|_X + \|Au\|_X \leq C \|f\|_X. \tag{2.3}$$

*Proof.* By applying the Fourier transform to equation (2.1) we obtain

$$[P_\varepsilon(\xi) + A + \lambda] \hat{u}(\xi) = \hat{f}(\xi). \tag{2.4}$$

By construction  $\lambda + P_\varepsilon(\xi) \in S_\varphi$ , for all  $t_k \in (0, t_0]$ ,  $\xi \in \mathbb{R}^n$  and the operator  $A + \lambda + P_\varepsilon(\xi)$  is invertible in  $E$ . So, from (2.4) we obtain that the solution of equation (2.1) can be represented in the form

$$u(x) = F^{-1} [A + \lambda + P_\varepsilon(\xi)]^{-1} \hat{f}. \tag{2.5}$$

By definition of the positive operator  $A$ , the inverse of  $A^{-1}$  is bounded in  $E$ . Then the operator  $A$  is a closed linear operator (as an inverse of bounded linear operator  $A^{-1}$ ). By the differential properties of the Fourier transform and by using (2.5) we have

$$\begin{aligned} \|Au\|_X &= \left\| F^{-1} A [A + \lambda + P_\varepsilon(\xi)]^{-1} \hat{f} \right\|_X, \\ \|D^\alpha u\|_X &= \left\| F^{-1} \xi^\alpha [A + \lambda + P_\varepsilon(\xi)]^{-1} \hat{f} \right\|_X, \end{aligned}$$

where  $X = L_p(\mathbb{R}^n; E)$ . Hence, it suffices to show that operator-functions

$$\begin{aligned} \sigma(\varepsilon, \lambda, \xi) &= A [A + \lambda + P_\varepsilon(\xi)]^{-1}, \\ \sigma_\alpha(\varepsilon, \lambda, \xi) &= \varepsilon(\alpha) |\lambda|^{1 - \frac{|\alpha|}{m}} \xi^\alpha [A + \lambda + P_\varepsilon(\xi)]^{-1} \end{aligned}$$

are collections of multipliers in  $X$  uniformly with respect to  $\varepsilon_k \in (0, \varepsilon_0]$  and  $\lambda \in S_{\varphi_2}$ . By virtue of [5, Lemma 2.3], for  $\lambda \in S_{\varphi_1}$  and  $\nu \in S_{\varphi_2}$  with  $\varphi_1 + \varphi_2 < \pi$  there is a positive constant  $C$  such that

$$|\lambda + \nu| \geq C(|\lambda| + |\nu|). \tag{2.6}$$

By using the positivity properties of operator  $A$ , we get that

$$B(\lambda, \varepsilon) = [A + \lambda + P_\varepsilon(\xi)]^{-1}$$

is bounded for all  $\xi \in \mathbb{R}^n$ ,  $\lambda \in S_{\varphi_1}$ ,  $\varepsilon_k \in (0, \varepsilon_0]$  and

$$\|B(\lambda, \varepsilon)\| \leq C(1 + |\lambda + P_\varepsilon(\xi)|)^{-1}.$$

By using Condition 2.1 and estimate (2.6), we obtain that

$$\|B(\lambda, \varepsilon)\| \leq C(1 + |\lambda| + |P_\varepsilon(\xi)|)^{-1} \leq C_2 \left[ 1 + |\lambda| + \sum_{k=1}^n \varepsilon_k |\xi_k|^m \right]^{-1}. \tag{2.7}$$

Then by resolvent properties of positive operators and uniform estimate (2.7) we obtain

$$\begin{aligned} \|\sigma(\varepsilon, \lambda, \xi)\| &\leq \left\| I + (\lambda + P_\varepsilon(\xi)) [A + \lambda + P_\varepsilon(\xi)]^{-1} \right\| \\ &\leq 1 + (|\lambda| + |P_\varepsilon(\xi)|) (1 + |\lambda| + |P_\varepsilon(\xi)|)^{-1} \leq C_3, \end{aligned}$$

where  $I$  is an identity operator in  $E$ . Moreover, by using the well known inequality

$$y_1^{\beta_1} y_2^{\beta_2} \dots y_n^{\beta_n} \leq C \left( 1 + \sum_{k=1}^n y_k^m \right)$$

for  $|\beta| \leq m$ ,  $y_k > 0$  and  $\beta = (\beta_1, \beta_2, \dots, \beta_n)$  for all  $u \in E$ , we have

$$\begin{aligned} \|\sigma_\alpha(\varepsilon, \lambda, \xi) u\|_E &\leq \varepsilon(\alpha) |\lambda|^{1 - \frac{|\alpha|}{m}} |\xi^\alpha| \|B(\lambda, t) u\|_E \\ &\leq |\lambda| \prod_{k=1}^n \left( \varepsilon_k^{\frac{1}{m}} |\lambda|^{\frac{1}{m}} |\xi_k| \right)^{\alpha_k} \|B(\lambda, \varepsilon) u\|_E \\ &\leq C_\alpha \left( |\lambda| + \sum_{k=1}^n \varepsilon_k |\xi_k|^m \right) \|B(\lambda, \varepsilon) u\|_E. \end{aligned}$$

In view of estimate (2.7) and by Condition 2.1 we get from the above inequality

$$\|\sigma_\alpha(\varepsilon, \lambda, \xi) u\|_E \leq C_\alpha \|u\|_E.$$

So, we obtain that the operator functions  $\sigma(\varepsilon, \lambda, \xi)$  and  $\sigma_\alpha(\varepsilon, \lambda, \xi)$  are uniformly bounded, i.e.,

$$\|\sigma(\varepsilon, \lambda, \xi)\|_{B(E)} \leq C, \|\sigma_\alpha(\varepsilon, \lambda, \xi)\|_{B(E)} \leq C_\alpha. \tag{2.8}$$

Due to  $R$ -positivity of  $A$ , by (2.8) and by Kahane’s contraction principle [6, Lemma 3.5], we obtain that the set  $\{\sigma(\varepsilon, \lambda, \xi); \xi \in \mathbb{R}^n \setminus \{0\}\}$  is uniformly  $R$ -bounded, i.e.,

$$\sup_{\varepsilon, \lambda} R\{\sigma(\varepsilon, \lambda, \xi); \xi \in \mathbb{R}^n \setminus \{0\}\} \leq M_0.$$

In a similar way for  $\beta = (\beta_1, \beta_2, \dots, \beta_n)$ ,  $\beta_i \in \{0, 1\}$ , we obtain

$$R\left(\left\{|\xi|^{|\beta|} D_\xi^\beta \sigma(\varepsilon, \lambda, \xi) : \xi \in \mathbb{R}^n \setminus \{0\}\right\}\right) \leq M. \tag{2.9}$$

Consider the following sets

$$\begin{aligned} \sigma^\beta(\varepsilon, \lambda, \xi) &= \left\{ |\xi|^{|\beta|} D_\xi^\beta \sigma(\varepsilon, \lambda, \xi) : \xi \in \mathbb{R}^n \setminus \{0\} \right\}, \\ \sigma_\alpha^\beta(\varepsilon, \lambda, \xi) &= \left\{ |\xi|^{|\beta|} D_\xi^\beta \sigma_\alpha(\varepsilon, \lambda, \xi) : \xi \in \mathbb{R}^n \setminus \{0\} \right\}, \\ \beta &= (\beta_1, \beta_2, \dots, \beta_n), \beta_i \in \{0, 1\}. \end{aligned}$$

In view of the  $R$ -positivity properties of operator  $A$ , due to Kahane’s contraction, addition and product properties of the collection of  $R$ -bounded operators (see e.g. [7, 23]),



by (2.9) for all  $\{\xi^{(j)}\} \in \mathbb{R}^n$ ,  $\{\sigma_\alpha^\beta(\varepsilon, \lambda, \xi^{(j)})\}$ ,  $j = 1, 2, \dots, \mu$  and  $u_1, u_2, \dots, u_\mu \in E$  and independent symmetric  $\{-1, 1\}$ -valued random variables  $r_j(y)$ ,  $\mu \in \mathbb{N}$  we obtain the following uniform estimate

$$\begin{aligned} & \int_{\Omega} \left\| \sum_{j=1}^{\mu} r_j(y) \sigma_\alpha^\beta(\varepsilon, \lambda, \xi^{(j)}) u_j \right\|_E dy \\ & \leq C \int_{\Omega} \left\| \sum_{j=1}^{\mu} \sigma^\beta(\varepsilon, \lambda, \xi^{(j)}) r_j(y) u_j \right\|_E dy \leq C \int_{\Omega} \left\| \sum_{j=1}^{\mu} r_j(y) u_j \right\|_E dy, \end{aligned}$$

i.e.,

$$R\left(\left\{\xi^\beta D_\xi^\beta \sigma_\alpha(\varepsilon, \lambda, \xi) : \xi \in \mathbb{R}^n \setminus \{0\}\right\}\right) \leq M_\beta.$$

Hence, we infer that the operator-valued functions  $\sigma(\varepsilon, \lambda, \xi)$  and  $\sigma_\alpha(\varepsilon, \lambda, \xi)$  are uniform  $R$ -bounded multipliers and it's  $R$ -bounds are independent of  $\varepsilon$  and  $\lambda$ . By virtue of Proposition A<sub>0</sub>, the operator-valued functions  $\sigma(\varepsilon, \lambda, \xi)$  and  $\sigma_\alpha(\varepsilon, \lambda, \xi)$  are uniform collections of Fourier multipliers in  $X$ . So, we obtain that for all  $f \in X$  there is a unique solution of equation (2.1) and estimate (2.3) holds.

Let  $O_\varepsilon$  denote the operator in  $X$  generated by problem (2.1) for  $\lambda = 0$ , i.e.,

$$D(O_\varepsilon) \subset H_p^m(\mathbb{R}^n; E(A), E), \quad O_\varepsilon u = P_\varepsilon(D)u + Au.$$

Theorem 2.1 and the definition of the space  $H_p^m(\mathbb{R}^n; E(A), E)$  imply the following result:

RESULT 2.1. Assume all conditions of Theorem 2.1 are satisfied. Then there are positive constants  $C_1$  and  $C_2$  so that

$$C_1 \|O_\varepsilon u\|_X \leq \|u\|_{H_{p,\varepsilon}^m(\mathbb{R}^n; E(A), E)} \leq C_2 \|O_\varepsilon u\|_X$$

for  $u \in Y$ . Indeed, if we put  $\lambda = 1$  in (2.3), by Theorem 2.1 we get

$$\sum_{|\alpha| \leq m} \varepsilon(\alpha) \|D^\alpha u\|_X + \|Au\|_X \leq C \|O_\varepsilon u\|_X \tag{2.10}$$

for  $u \in Y$ . Due to the closedness of  $A$  and by the differential properties of the Fourier transform, we have

$$\|Au\|_X = \|F^{-1}A\hat{u}\|_X, \quad \|D^\alpha u\|_X = \|F^{-1}\xi^\alpha \hat{u}\|_X.$$

So, in view of estimate (2.10) and by definition of  $Y$ , we obtain

$$\|u\|_{H_{p,\varepsilon}^m(\mathbb{R}^n; E(A), E)} \leq C_2 \|O_\varepsilon u\|_X.$$

The first inequality is equivalent to the following estimate

$$\begin{aligned} & \|F^{-1}A\hat{u}\|_X + \|F^{-1}P_\varepsilon(\xi)\hat{u}\|_X \\ & \leq C \left\{ \|F^{-1}A\hat{u}\|_X + \left\| F^{-1} \left[ 1 + \left( \sum_{k=1}^n \varepsilon_k^{\frac{2}{m}} \xi_k^2 \right)^{\frac{1}{2}} \right]^m \hat{u} \right\|_X \right\}. \end{aligned}$$

So, it suffices to show that the operator functions

$$A \left\{ A + \left[ 1 + \left( \sum_{k=1}^n \varepsilon_k^{\frac{2}{m}} \xi_k^2 \right)^{\frac{1}{2}} \right]^m \right\}^{-1}, \varepsilon(\alpha) \xi^\alpha \left[ 1 + \left( \sum_{k=1}^n \varepsilon_k^{\frac{2}{m}} \xi_k^2 \right)^{\frac{1}{2}} \right]^{-m}$$

are uniform Fourier multipliers in  $X$ . This fact is proved in a similar way as in the proof of Theorem 2.1.

From Theorem 2.1, we have:

RESULT 2.2. Assume all conditions of Theorem 2.1 hold. Then, for all  $\lambda \in S_\varphi$  the resolvent of operator  $O_\varepsilon$  exists and the following sharp uniform estimate holds

$$\sum_{|\alpha| \leq m} \varepsilon(\alpha) \left\| D^\alpha (O_\varepsilon + \lambda)^{-1} \right\|_{B(X)} + \left\| A(O_\varepsilon + \lambda)^{-1} \right\|_{B(X)} \leq C. \tag{2.11}$$

Indeed, we infer from Theorem 2.1 that the operator  $O_\varepsilon + \lambda$  has a bounded inverse from  $X$  to  $Y$ . So, the solution  $u$  of the equation (2.1) can be expressed as  $u(x) = (O_\varepsilon + \lambda)^{-1} f$  for all  $f \in X$ . Then estimate (2.3) implies the estimate (2.11).

RESULT 2.3. Theorem 2.1 particularly implies that the operator  $O_\varepsilon$  is positive in  $X$ . Then the operators  $O_\varepsilon^\sigma$  are generators of analytic semigroups in  $X$  for  $\sigma \leq \frac{1}{2}$  (see e.g. [20, §1.14.5]).

Now consider the problem (1.1). By using Theorem 2.1 and the perturbation theory of linear operators, we have:

THEOREM 2.2. Assume all conditions of Theorem 2.1 are satisfied. Suppose  $A_\alpha(x) A^{-\left(1 - \frac{|\alpha|}{m} - \mu\right)} \in L_\infty(\mathbb{R}^n; B(E))$  for  $\mu \in \left(0, 1 - \frac{|\alpha|}{m}\right)$ . Then for  $f \in X$ ,  $\lambda \in S_{\varphi_2}$ ,  $0 \leq \varphi_2 < \pi - \varphi_1$ ,  $\varphi_1 + \varphi_2 \leq \varphi$  and for sufficiently large  $|\lambda|$  there is a unique solution  $u$  of the equation (1.1) belonging to  $Y$  and the following coercive uniform estimate holds

$$\sum_{|\alpha| \leq m} \varepsilon(\alpha) |\lambda|^{1 - \frac{|\alpha|}{m}} \|D^\alpha u\|_X + \|Au\|_X \leq C \|f\|_X. \tag{2.12}$$

*Proof.* It is clear that  $Q_\varepsilon = O_\varepsilon + L_\varepsilon$ , where  $O_\varepsilon$  is the operator in  $X$  generated by problem (2.1) for  $\lambda = 0$  and

$$L_\varepsilon u = \sum_{|\alpha| < m} \varepsilon(\alpha) A_\alpha(x) D^\alpha u \text{ for } u \in Y.$$

In view of the condition on  $A_\alpha(x)$  and by Theorem A<sub>1</sub> for  $u \in Y$  we have

$$C_\mu \sum_{|\alpha| < m} \varepsilon(\alpha) \left\| A^{1 - \frac{|\alpha|}{m} - \mu} D^\alpha u \right\|_X \leq C_\mu \left[ h^\mu \|u\|_{H_{p,\varepsilon}^m(\mathbb{R}^n, E(A), E)} + h^{-(1-\mu)} \|u\|_X \right]. \tag{2.13}$$

$$\|L_\varepsilon u\|_X \leq \sum_{|\alpha| < m} \varepsilon(\alpha) \|A_\alpha(x) D^\alpha u\|_X.$$

Then from estimates (2.12), (2.13) and for  $u \in Y$  we obtain

$$\|L_\varepsilon u\|_X \leq C \left[ h^\mu \|O_\varepsilon u\|_X + h^{-(1-\mu)} \|u\|_X \right]. \tag{2.14}$$

Since  $\|u\|_X = \frac{1}{\lambda} \|(O_\varepsilon + \lambda)u + L_\varepsilon u\|_X$  for  $\lambda \in S_{\varphi_2}$ , we get

$$\begin{aligned} \|u\|_X &\leq \frac{1}{|\lambda|} [\|(O_\varepsilon + \lambda)u\|_X + \|O_\varepsilon u\|_X] \\ &\leq \frac{1}{|\lambda|} \|(O_\varepsilon + \lambda)u\|_X + \frac{1}{|\lambda|} \left[ \sum_{|\alpha| < m} \varepsilon(\alpha) \|D^\alpha u\|_X + \|Au\|_X \right]. \end{aligned} \tag{2.15}$$

From estimates (2.13) – (2.15) for  $u \in Y$ , we obtain

$$\|L_\varepsilon u\|_X \leq Ch^\mu \|(O_\varepsilon + \lambda)u\|_X + C_1 |\lambda|^{-1} h^{-(1-\mu)} \|(O_\varepsilon + \lambda)u\|_X. \tag{2.16}$$

Then choosing  $h$  and  $\lambda$  such that  $Ch^\mu < 1$ ,  $C_1 |\lambda|^{-1} h^{-(1-\mu)} < 1$  from (2.16), we obtain that

$$\left\| L_\varepsilon (O_\varepsilon + \lambda)^{-1} \right\|_{B(X)} < 1. \tag{2.17}$$

From Theorem 2.1 and (2.17) we get that the operator  $(Q_\varepsilon + \lambda)$  has a bounded inverse in  $X$ . Moreover, it is clear that

$$(Q_\varepsilon + \lambda)^{-1} = \left[ I + L_\varepsilon (O_\varepsilon + \lambda)^{-1} \right] (O_\varepsilon + \lambda), \tag{2.18}$$

where  $I$  is an identity operator in  $X$ . Using relation (2.18), estimates (2.3), (2.17) and perturbation theory of linear operators, we obtain that the operator  $Q_\varepsilon + \lambda$  has a bounded inverse from  $X$  into  $Y$  and the estimate (2.12) holds.

### 3. The Cauchy problem for parabolic FDOE with parameter

In this section, we shall consider the following Cauchy problem for the parabolic FDOE

$$\frac{\partial u}{\partial t} + P_\varepsilon(D)u + Au = f(t, x), \quad u(0, x) = 0, \tag{3.1}$$

where  $P_\varepsilon(D)$  is the fractional differential operator defined by (2.2) and  $A$  is a linear operator in  $E$ ,  $\varepsilon = (\varepsilon_1, \varepsilon_2, \dots, \varepsilon_n)$ ,  $\varepsilon_k$  are positive parameters.

By applying Theorem 2.1 we establish the maximal regularity of the problem (3.1) in  $E$ -valued mixed  $L_{\mathbf{p}}$  spaces, where  $\mathbf{p} = (p_1, p)$ . Let  $O_\varepsilon$  denote the operator generated by problem (2.1). For this aim we need the following result:

**THEOREM 3.1.** *Suppose Condition 2.1 holds,  $E$  is an UMD space and operator  $A$  is  $R$ -positive in  $E$  with  $0 \leq \varphi < \pi - \varphi_1$ . Then operator  $O_\varepsilon$  is uniformly  $R$ -positive in  $L_p(\mathbb{R}^n; E)$ .*

*Proof.* From Result 2.3 we obtain that the operator  $O_\varepsilon$  is positive in  $X = L_p(\mathbb{R}^n; E)$ . We have to prove the  $R$ -boundedness of the set

$$\sigma(\varepsilon, \lambda, \xi) = \left\{ \lambda (O_\varepsilon + \lambda)^{-1} : \lambda \in S_\varphi \right\}.$$

From the proof of Theorem 2.1, we have

$$\lambda (O_\varepsilon + \lambda)^{-1} f = F^{-1} \Phi(\varepsilon, \xi, \lambda) \hat{f}, f \in X,$$

where

$$\Phi(\varepsilon, \xi, \lambda) = \lambda (A + P_\varepsilon(\xi) + \lambda)^{-1}.$$

By reasoning as in the proof of Theorem 2.1, we obtain the following uniform estimate

$$\|\Phi(\varepsilon, \xi, \lambda)\|_{B(E)} \leq |\lambda| \left\| (A + P_\varepsilon(\xi) + \lambda)^{-1} \right\|_{B(E)} \leq C.$$

By definition of  $R$ -boundedness, it suffices to show that the operator function  $\Phi(\varepsilon, \xi, \lambda)$  (which depends on variable  $\lambda$  and parameters  $\xi, \varepsilon$ ) is a multiplier in  $X$  uniformly with respect to  $\xi$  and  $\varepsilon$ . Indeed, by reasoning as in Theorem 2.1 we can easily show that  $\Phi(\varepsilon, \xi, \lambda)$  is a uniform multiplier in  $L_p(\mathbb{R}; E)$ . Then, by the definition of a  $R$ -bounded set we have

$$\begin{aligned} \int_0^1 \left\| \sum_{j=1}^m r_j(y) \lambda_j (O_\varepsilon + \lambda_j)^{-1} f_j \right\|_X dy &= \int_0^1 \left\| \sum_{j=1}^m r_j(y) F^{-1} \Phi(\varepsilon, \xi, \lambda_j) \hat{f}_j \right\|_X dy \\ &= \int_0^1 \left\| F^{-1} \sum_{j=1}^m r_j(y) \Phi(\varepsilon, \xi, \lambda_j) \hat{f}_j \right\|_X dy \leq C \int_0^1 \left\| \sum_{j=1}^m r_j(y) f_j \right\|_X dy \end{aligned}$$

for all  $\xi \in \mathbb{R}^n, \lambda_1, \lambda_2, \dots, \lambda_m \in S_\varphi, f_1, f_2, \dots, f_m \in X, m \in N$ , where  $\{r_j\}$  is a sequence of independent symmetric  $\{-1, 1\}$ -valued random variables on  $[0, 1]$ . Hence, the set  $\Phi(\varepsilon, \xi, \lambda)$  is uniformly  $R$ -bounded.

Let  $E$  be a Banach space. For  $\mathbf{p} = (p, p_1), Z = L_{\mathbf{p}}(\mathbb{R}_+^{n+1}; E)$  will denote the space of all  $\mathbf{p}$ -summable  $E$ -valued functions with mixed norm, i.e., the space of all measurable  $E$ -valued functions  $f$  defined on  $\mathbb{R}_+^{n+1}$  for which

$$\|f\|_{L_{\mathbf{p}}(\mathbb{R}_+^{n+1}; E)} = \left( \int_{\mathbb{R}^n} \left( \int_0^\infty \|f(t, x)\|_E^p dx \right)^{\frac{p_1}{p}} dt \right)^{\frac{1}{p_1}} < \infty.$$

Let  $E$  be a Banach space and  $A$  be a positive operator in  $E$ . Suppose,  $l$  is a positive integer.  $W_p^l(0, \infty; E(A), E)$  denotes the space of all functions  $u \in L_p(0, \infty; E(A))$  possessing the generalized derivatives  $u^{(l)} \in L_p(0, \infty; E)$  with the norm

$$\|u\|_{W_p^l(0, \infty; E(A), E)} = \|Au\|_{L_p(0, \infty; E)} + \|u^{(l)}\|_{L_p(0, \infty; E)}.$$

Let  $m$  be a positive number.  $W_{\mathbf{p}}^{1,m}(\mathbb{R}_+^{n+1}; E(A), E)$  denotes the space of all functions  $u \in L_{\mathbf{p}}(\mathbb{R}_+^{n+1}; E(A))$  possessing the generalized derivative  $D_t u = \frac{\partial u}{\partial t} \in Z$  with respect to  $y$  and fractional derivatives  $D_x^\alpha u \in Z$  with respect to  $x$  for  $|\alpha| \leq m$  with the norm

$$\|u\|_{W_{\mathbf{p}}^{1,m}(\mathbb{R}_+^{n+1}; E(A), E)} = \|Au\|_Z + \left\| \frac{du}{dt} \right\|_Z + \sum_{|\alpha| \leq m} \|D_x^\alpha u\|_Z,$$

where  $u = u(t, x)$ .

Now, we are ready to state the main result of this section.

**THEOREM 3.2.** *Assume the conditions of Theorem 2.1 hold for  $\varphi \in (\frac{\pi}{2}, \pi)$ . Then for  $f \in Z$  problem (3.1) has a unique solution*

$$u \in W_{\mathbf{p}}^{1,m}(\mathbb{R}_+^{n+1}; E(A), E)$$

satisfying the following uniform coercive estimate

$$\left\| \frac{du}{dt} \right\|_Z + \|P_\varepsilon(D)u\|_Z + \|Au\|_Z \leq C \|f\|_Z.$$

*Proof.* By definition of  $X = L_p(\mathbb{R}^n; E)$  and mixed space  $L_{\mathbf{p}}(\mathbb{R}_+^{n+1}; E)$ ,  $\mathbf{p} = (p, p_1)$ , we have

$$\begin{aligned} \|u\|_{L_{p_1}(0,\infty;X)} &= \left( \int_0^\infty \|u(y)\|_X^{p_1} dt \right)^{\frac{1}{p_1}} = \left( \int_0^\infty \|u(y)\|_{L_p(\mathbb{R}^n;E)}^{p_1} dt \right)^{\frac{1}{p_1}} \\ &= \left( \int_{\mathbb{R}^n} \left( \int_0^\infty \|u(y,x)\|_E^p dt \right)^{\frac{p_1}{p}} dx \right)^{\frac{1}{p_1}} = \|u\|_Z. \end{aligned}$$

Moreover, by definition of the space  $W_p^m(0, \infty; E(A), E)$  and by Result 2.1, we obtain

$$\begin{aligned} \|u\|_{W_{p_1}^1(0,\infty;D(O_\varepsilon),X)} &= \|O_\varepsilon u\|_{L_p(0,\infty;X)} + \|u'\|_{L_p(0,\infty;X)} \\ &= \|Au\|_Z + \|D_t u\|_Z + \sum_{|\alpha| \leq m} \|D_x^\alpha u\|_Z \\ &= \|u\|_{W_{\mathbf{p}}^{1,m}(\mathbb{R}_+^{n+1};E(A),E)}. \end{aligned} \tag{3.2}$$

Hence, we deduced from the above equalities that,

$$\begin{aligned} Z &= L_{\mathbf{p}}(\mathbb{R}_+^{n+1}; E) = L_{p_1}(0, \infty; X), W_{\mathbf{p}}^{1,m}(\mathbb{R}_+^{n+1}; E(A), E) \\ &= W_{p_1}^1(0, \infty; D(O_\varepsilon), X). \end{aligned}$$

Therefore, the problem (3.1) can be expressed as the following Cauchy problem for the abstract parabolic equation

$$\frac{du}{dt} + O_\varepsilon u(t) = f(t), \quad u(0) = 0, \quad t \in (0, \infty). \tag{3.3}$$

By virtue of [1, Theorem 4.5.2], the condition  $E \in UMD$  implies  $X \in UMD$  for  $p \in (1, \infty)$ . Then due to the  $R$ -positivity of  $O_\varepsilon$  and by virtue of [23, Theorem 4.2], we obtain that for  $f \in L_{p_1}(0, \infty; X)$  equation (3.3) has a unique solution  $u \in W_{p_1}^1(0, \infty; D(O_\varepsilon), X)$  satisfying the following uniform estimate

$$\left\| \frac{du}{dt} \right\|_{L_{p_1}(0,\infty;X)} + \|O_\varepsilon u\|_{L_{p_1}(0,\infty;X)} \leq C \|f\|_{L_{p_1}(0,\infty;X)}.$$

From the Theorem 2.1, relation (3.2) and from the above estimate we get the assertion.

### 4. BVP for anisotropic FDE

In this section, the maximal regularity properties of the anisotropic FDE is studied.

Let  $\tilde{\Omega} = \Omega \times \mathbb{R}^n$ , where  $\Omega \subset \mathbb{R}^{\mu}$  is an open connected set with compact  $C^{2l}$ -boundary  $\partial\Omega$ . Consider the BVP for the FDE

$$P_{\varepsilon}(D)u + \sum_{|\alpha| \leq 2l} (b_{\alpha} D_y^{\alpha} + \lambda)u = f(x, y), \quad y \in \Omega, \tag{4.1}$$

$$B_j u = \sum_{|\beta| \leq l_j} b_{j\beta}(y) D_y^{\beta} u(x, y) = 0, \quad x \in \mathbb{R}^n, \tag{4.2}$$

$$y \in \partial\Omega, \quad j = 1, 2, \dots, l,$$

where  $u = u(x, y)$ ,  $P_{\varepsilon}(D)$  is the fractional differential operator defined by (2.1),

$$D_j = -i \frac{\partial}{\partial y_j}, \quad y = (y_1, \dots, y_{\mu}), \quad b_{\alpha} = b_{\alpha}(y),$$

$\alpha = (\alpha_1, \alpha_2, \dots, \alpha_{\mu})$ ,  $\beta = (\beta_1, \beta_2, \dots, \beta_{\mu})$  are nonnegative integer numbers, and  $\varepsilon = (\varepsilon_1, \varepsilon_2, \dots, \varepsilon_n)$  and  $\varepsilon_k$  are positive parameters. For  $\tilde{\Omega} = \mathbb{R}^n \times \Omega$ ,  $\mathbf{p} = (p_1, p)$  here,  $L_{\mathbf{p}}(\tilde{\Omega})$  will denote the space of all  $\mathbf{p}$ -summable scalar-valued functions with mixed norm i.e., the space of all measurable functions  $f$  defined on  $\tilde{\Omega}$ , for which

$$\|f\|_{L_{\mathbf{p}}(\tilde{\Omega})} = \left( \int_{\mathbb{R}^n} \left( \int_{\Omega} |f(x, y)|^{p_1} dx \right)^{\frac{p}{p_1}} dy \right)^{\frac{1}{p}} < \infty.$$

Analogously,  $W_{\mathbf{p}}^{m, 2l}(\tilde{\Omega})$  denotes the anisotropic fractional Sobolev space with corresponding mixed norm, i.e.,  $W_{\mathbf{p}}^{m, 2l}(\tilde{\Omega})$  denotes the space of all functions  $u \in L_{\mathbf{p}}(\tilde{\Omega})$  possessing the fractional derivatives  $D_x^{\alpha} u \in L_{\mathbf{p}}(\tilde{\Omega})$  with respect to  $x$  for  $|\alpha| \leq m$  and generalized derivative  $\frac{\partial^{2l} u}{\partial y_k^{2l}} \in L_{\mathbf{p}}(\tilde{\Omega})$  with respect to  $y$  with the norm

$$\|u\|_{W_{\mathbf{p}}^{m, 2l}(\tilde{\Omega})} = \sum_{|\alpha| \leq m} \|D_x^{\alpha} u\|_{L_{\mathbf{p}}(\tilde{\Omega})} + \sum_{k=1}^{\mu} \left\| \frac{\partial^{2l} u}{\partial y_k^{2l}} \right\|_{L_{\mathbf{p}}(\tilde{\Omega})}.$$

Let  $Q$  denote the operator generated by problem (4.1) – (4.2), i.e.,

$$D(Q) = W_{\mathbf{p}}^{m, 2l}(\tilde{\Omega}, B_j) = \left\{ u : u \in W_{\mathbf{p}}^{m, 2l}(\tilde{\Omega}), B_j u = 0, j = 1, 2, \dots, l \right\},$$

$$Qu = P_{\varepsilon}(D)u + \sum_{|\alpha| \leq 2l} b_{\alpha} D_y^{\alpha} u.$$

Let  $\xi' = (\xi_1, \xi_2, \dots, \xi_{\mu-1}) \in \mathbb{R}^{\mu-1}$ ,  $\alpha' = (\alpha_1, \alpha_2, \dots, \alpha_{\mu-1}) \in Z^\mu$  and

$$A(y_0, \xi', D_y) = \sum_{|\alpha'|+j \leq 2l} a_{\alpha'}(y_0) \xi_1^{\alpha_1} \xi_2^{\alpha_2} \dots \xi_{\mu-1}^{\alpha_{\mu-1}} D_\mu^j \text{ for } y_0 \in \bar{G}$$

$$B_j(y_0, \xi', D_y) = \sum_{|\beta'|+j \leq l_j} b_{j\beta'}(y_0) \xi_1^{\beta_1} \xi_2^{\beta_2} \dots \xi_{\mu-1}^{\beta_{\mu-1}} D_\mu^j \text{ for } y_0 \in \partial G.$$

CONDITION 4.1. Let the following conditions be satisfied:

(1)  $b_\alpha \in C(\bar{\Omega})$  for each  $|\alpha| = 2l$  and  $b_\alpha \in L_\infty(\Omega) + L_{r_k}(\Omega)$  for each  $|\alpha| = k < 2l$

with  $r_k \geq p_1$ ,  $p_1 \in (1, \infty)$  and  $2l - k > \frac{l}{r_k}$ ;

(2)  $b_{j\beta} \in C^{2l-l_j}(\partial\Omega)$  for each  $j, \beta, l_j < 2l, p \in (1, \infty), \lambda \in S_\varphi, \varphi \in [0, \pi]$ ;

(3) for  $y \in \bar{\Omega}, \xi \in R^\mu, \sigma \in S_{\varphi_0}, \varphi_0 \in (0, \frac{\pi}{2}), |\xi| + |\sigma| \neq 0$  let  $\sigma + \sum_{|\alpha|=2l} b_\alpha(y) \xi^\alpha \neq$

0;

(4) for each  $y_0 \in \partial\Omega$  local BVP in local coordinates corresponding to  $y_0$ ,

$$\lambda + A(y_0, \xi', D_y) \vartheta(y) = 0,$$

$$B_j(y_0, \xi', D_y) \vartheta(0) = h_j, j = 1, 2, \dots, l$$

has a unique solution  $\vartheta \in C_0(0, \infty)$  for all  $h = (h_1, h_2, \dots, h_l) \in \mathbb{C}^l$  and for  $\xi' \in \mathbb{R}^{\mu-1}$ .

Suppose  $v = (v_1, v_2, \dots, v_n)$  are nonnegative real numbers. In this section, we present the following result:

THEOREM 4.1. Assume Condition 2.1 and Condition 4.1 are satisfied. Then for  $f \in L_p(\tilde{\Omega}), \lambda \in S_\varphi, \varphi \in (0, \pi]$ , problem (4.1) – (4.2) has a unique solution  $u \in W_p^{m, 2l}(\tilde{\Omega})$  and the following coercive uniform estimate holds

$$\sum_{|v| \leq m} \prod_{k=1}^n \varepsilon_k^{\frac{v_k}{m}} |\lambda|^{1 - \frac{|v|}{m}} \|D_x^v u\|_{L_p(\tilde{\Omega})} + \sum_{|\alpha| \leq 2l} \|D_y^\alpha u\|_{L_p(\tilde{\Omega})} \leq C \|f\|_{L_p(\tilde{\Omega})}.$$

Proof. Let  $E = L_{p_1}(\Omega)$ . It is known [4] that  $L_{p_1}(\Omega)$  is an UMD space for  $p_1 \in (1, \infty)$ . Consider the operator  $A$  defined by

$$D(A) = W_{p_1}^{2l}(\Omega; B_j u = 0), Au = \sum_{|\alpha| \leq 2l} b_\alpha(x) D^\alpha u(y).$$

Therefore, the problem (4.1) – (4.2) can be rewritten in the form of (2.1), where  $u(x) = u(x, \cdot), f(x) = f(x, \cdot)$  are functions with values in  $E = L_{p_1}(\Omega)$ . From [6, Theorem 8.2] we get that the following problem

$$\eta u(y) + \sum_{|\alpha| \leq 2l} b_\alpha(y) D^\alpha u(y) = f(y), \tag{4.3}$$

$$B_j u = \sum_{|\beta| \leq l_j} b_{j\beta}(y) D^\beta u(y) = 0, j = 1, 2, \dots, l$$

has a unique solution for  $f \in L_{p_1}(\Omega)$  and  $\arg \eta \in S(\varphi_1), |\eta| \rightarrow \infty$ . Moreover, the operator  $A$  generated by (4.3) is  $R$ -positive in  $L_{p_1}$ . Then from Theorem 2.1 we obtain the assertion.

### 5. The system of FDE of infinite many order

Consider the following system of FDEs

$$P_\varepsilon(D)u_i + \sum_{j=1}^N (a_{ij} + \lambda)u_j(x) = f_i(x), \quad x \in \mathbb{R}^n, \quad (5.1)$$

$$i = 1, 2, \dots, N, \quad N \in [1, \infty],$$

where  $P_\varepsilon(D)$  is the fractional differential operator defined by (2.2),  $\varepsilon = (\varepsilon_1, \varepsilon_2, \dots, \varepsilon_n)$  and  $\varepsilon_k$  are positive parameters. Let  $a_{ij}$  be real numbers and

$$l_q(A) = \left\{ u \in l_q, \|u\|_{l_q(A)} = \|Au\|_{l_q} \right.$$

$$= \left. \left( \sum_{i=1}^N |(Au)_i|^q \right)^{\frac{1}{q}} = \left( \sum_{i=1}^N \left| \sum_{j=1}^N a_{ij}u_j \right|^q \right)^{\frac{1}{q}} < \infty \right\},$$

$$u = \{u_j\}, \quad Au = \left\{ \sum_{j=1}^N a_{ij}u_j \right\}, \quad i, j = 1, 2, \dots, N.$$

CONDITION 5.1. Let

$$a_{ij} = a_{ji}, \quad \sum_{i,j=1}^N a_{ij}\xi_i\xi_j \geq C_0|\xi|^2 \quad \text{for } \xi \neq 0.$$

Let

$$f(x) = \{f_i(x)\}_1^N, \quad u = \{u_i(x)\}_1^N.$$

**THEOREM 5.1.** Assume that the Condition 2.1 and Condition 5.1 are satisfied. Then, for  $f(x) \in L_p(\mathbb{R}^n; l_q)$ ,  $|\arg \lambda| \leq \varphi$ ,  $\varphi \in (0, \pi]$  and for sufficiently large  $|\lambda|$ , problem (5.1) has a unique solution  $u \in H_p^m(\mathbb{R}^n, l_q(A), l_q)$  and the following uniform coercive estimate holds

$$\sum_{|\alpha| \leq m} \varepsilon(\alpha) |\lambda|^{1 - \frac{|\alpha|}{m}} \left[ \int_{\mathbb{R}^n} \left( \sum_{j=1}^N |D^\alpha u_j(x)|^q \right)^{\frac{2}{q}} dx \right]^{\frac{1}{p}}$$

$$+ \left[ \int_{\mathbb{R}^n} \left( \sum_{i=1}^N \left| \sum_{j=1}^N a_{ij}u_j \right|^q \right)^{\frac{2}{q}} dx \right]^{\frac{1}{p}} \leq C \left[ \int_{\mathbb{R}^n} \left( \sum_{i=1}^N |f_i(x)|^q \right)^{\frac{2}{q}} dx \right]^{\frac{1}{p}}.$$

*Proof.* Let  $E = l_q$  and  $A$  be a matrix such that  $A = [a_{ij}]$ ,  $i, j = 1, 2, \dots, N$ . It is easy to see that

$$B(\lambda) = \lambda(A + \lambda)^{-1} = \frac{\lambda}{D(\lambda)} [A_{ji}(\lambda)], \quad i, j = 1, 2, \dots, N,$$



where  $D(\lambda) = \det(A - \lambda I)$ ,  $A_{ji}(\lambda)$  are entries of the corresponding adjoint matrix of  $A - \lambda I$ . Since the matrix  $A$  is symmetric and positive definite, it generates a positive operator in  $l_q$  for  $q \in (1, \infty)$ . For all  $u_1, u_2, \dots, u_\mu \in l_q$ ,  $\lambda_1, \lambda_2, \dots, \lambda_\mu \in \mathbb{C}$  and independent symmetric  $\{-1, 1\}$ -valued random variables  $r_k(y)$ ,  $k = 1, 2, \dots, \mu$ ,  $\mu \in \mathbb{N}$  we have

$$\begin{aligned} & \int_{\Omega} \left\| \sum_{k=1}^{\mu} r_k(y) B(\lambda_k) u_k \right\|_{l_q}^q dy \\ & \leq C \left\{ \int_{\Omega} \sum_{j=1}^N \left| \sum_{k=1}^{\mu} \sum_{j=1}^N \frac{\lambda_k}{D(\lambda_k)} A_{ji}(\lambda_k) r_k(y) u_{kj} \right|^q dy \right. \\ & \left. \leq \sup_{k,i} \sum_{j=1}^N \left| \frac{\lambda_k}{D(\lambda_k)} A_{ji}(\lambda_k) \right|^q \int_{\Omega} \left| \sum_{k=1}^{\mu} r_k(y) u_{kj} \right|^q dy. \right. \end{aligned} \tag{5.2}$$

Since  $A$  is symmetric and positive definite, we have

$$\sup_{k,i} \sum_{j=1}^N \left| \frac{\lambda_k}{D(\lambda_k)} A_{ji}(\lambda_k) \right|^q \leq C.$$

From (5.2) and (5.3) we get

$$\int_{\Omega} \left\| \sum_{k=1}^{\mu} r_k(y) B(\lambda_k) u_k \right\|_{l_q}^q dy \leq C \int_{\Omega} \left\| \sum_{k=1}^{\mu} r_k(y) u_k \right\|_{l_q}^q dy,$$

i.e., the operator  $A$  is  $R$ -positive in  $l_q$ . From Theorem 2.1, we obtain that problem (5.1) has a unique solution  $u \in H_p^m(\mathbb{R}^n; l_q(A), l_q)$  for  $f \in L_p(\mathbb{R}^n; l_q)$  and the following estimate holds

$$\sum_{|\alpha| \leq m} \varepsilon(\alpha) |\lambda|^{1 - \frac{|\alpha|}{m}} \|D^\alpha u\|_{L_p(\mathbb{R}^n; l_q)} + \|Au\|_{L_p(\mathbb{R}^n; l_q)} \leq M \|f\|_{L_p(\mathbb{R}^n; l_q)}.$$

From the above estimate we obtain the assertion.

*Acknowledgements.* The author would like to express a gratitude to Dr. Erchan Aptoula for his useful advices in English in preparing of this paper.

REFERENCES

[1] AMANN H., *Linear and quasi-linear equations*,1, Birkhauser, Basel 1995.  
 [2] AGARWAL R., BOHNER M., SHAKHMUROV V. B., *Maximal regular boundary value problems in Banach-valued weighted spaces*, Bound. Value Probl., (1)(2005), 9–42.  
 [3] ASHYRALYEV A., SHARIFOV Y. A., *Existence and uniqueness of solutions for the system of non-linear fractional differential equations with nonlocal and integral boundary conditions*, Abstr. Appl. Anal., ID 594802, (2012).

- [4] BURKHOLDER D. L., *A geometrical conditions that implies the existence certain singular integral of Banach space-valued functions*, Proc. conf. Harmonic analysis in honor of Antonu Zigmund, Chicago, 1981, Wads Worth, Belmont, (1983), 270–286.
- [5] DORE C. AND YAKUBOV S., *Semigroup estimates and non coercive boundary value problems*, Semigroup Forum 60 (2000), 93–121.
- [6] DUBINSKII YU. A., *The Cauchy problem and pseudodifferential operators in the complex domain*, Uspekhi Mat. Nauk, 45:2(272) (1990), 115–142.
- [7] DIESTEL J., JARCHOW H., TONGE A., *Absolutely summing operators*, Cambridge Univ. Press, Cambridge, 1995.
- [8] DENK R., KRAINER T., *R-Boundedness, pseudodifferential operators, and maximal regularity for some classes of partial differential operators*, Manuscripta Math. 124(3) (2007), 319–342.
- [9] HALLER R., HECK H., NOLL A., *Mikhlin's theorem for operator-valued Fourier multipliers in  $n$  variables*, Math. Nachr. 244 (2002), 110–130.
- [10] HERMANDER L., *The analysis of linear partial differential operators*, V. 3, 4, Springer-Verlag, New York, 1985.
- [11] FAVINI A., SHAKHMUROV V. B., YAKOV YAKUBOV, *Regular boundary value problems for complete second order elliptic differential-operator equations in UMD Banach spaces*, Semigroup form, 79 (1)(2009), 22–54.
- [12] KREIN S. G., *Linear differential equations in Banach space*, American Mathematical Society, Providence, 1971.
- [13] LIZORKIN P. I., *Generalized Liouville differentiation and functional spaces  $L_p^s(E_n)$* , Embedding theorems, Mathematics of the USSR-Sbornik (3)60 (1963), 325–353.
- [14] NAGASE M., *The  $L_p$ -boundedness of pseudodifferential operators with non-regular symbols*, Comm. in Partial Differential Equations, 2 (10), (1977), 1045–1061.
- [15] PORTAL P., ŠTRKALJ Ž., *Pseudodifferential operators on Bochner spaces and an application*, Math. Z. 253 (2006), 805–819.
- [16] SHAKHMUROV V. B., *Coercive boundary value problems for regular degenerate differential-operator equations*, J. Math. Anal. Appl., 292 (2) (2004), 605–620.
- [17] SHAKHMUROV V. B., *Embedding and separable differential operators in Sobolev-Lions type spaces*, Mathematical Notes, (84) 6, (2008), 906–926.
- [18] SHAKHMUROV V. B., *Maximal regular abstract elliptic equations and applications*, Siberian Math. Journal, (51) 5, (2010), 935–948.
- [19] SHAKHMUROV V. B., *Embedding and maximal regular differential operators in Banach-valued weighted spaces*, Acta Math. Sinica, 22(5) (2006), 1493–1508.
- [20] TRIEBEL H., *Interpolation theory, Function spaces, Differential operators*, North-Holland, Amsterdam, 1978.
- [21] TAYLOR M., *Pseudo-differential operators*, Princeton University press, Princeton, NJ, 1981.
- [22] TREVES F., *Introduction to pseudo-differential and Fourier integral operators*, Plenum press, New York and London, 1980.
- [23] WEIS L., *Operator-valued Fourier multiplier theorems and maximal  $L_p$  regularity*, Math. Ann. 319(2001), 735–758.
- [24] YAKUBOV S., YAKUBOV YA., *Differential-operator Equations. Ordinary and Partial Differential Equations*, Chapman and Hall /CRC, Boca Raton, 2000.

(Received May 9, 2018)

Veli Shakhmurov  
 Istanbul Okan University  
 Department of Mechanical Engineering  
 Akfirat, Tuzla 34959 Istanbul, Turkey  
 e-mail: veli.sahmurov@okan.edu.tr