

## BESSEL PROPERTY AND BASICITY OF THE SYSTEM OF ROOT VECTOR–FUNCTIONS OF DIRAC OPERATOR WITH SUMMABLE COEFFICIENT

VALI M. KURBANOV AND AFSANA M. ABDULLAYEVA

(Communicated by G. Teschl)

*Abstract.* In the paper we study one-dimensional Dirac operator

$$Dy = By' + P(x)y, \quad y = (y_1, y_2)^T,$$

where  $B = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ ,  $P(x) = \text{diag}(p(x), q(x))$ ,  $p(x)$  and  $q(x)$  are complex valued functions from the class  $L_1(G)$ ,  $G = (0, 2\pi)$ .

Necessary and sufficient conditions of Bessel property and unconditional basicity (the Riesz basicity) of the system of root-functions of the operator  $D$  in  $L_2^2(G)$  are set up. A theorem on equivalent basicity for these systems in  $L_2^2(G)$  is proved.

### 1. Introduction and statement of results

In the paper we study Bessel property and basicity of the system of root vector-functions of one-dimensional Dirac operator with summable coefficient (potential). The root vector-functions are understood in generalized representation, i.e. without regard to boundary conditions (see [8]). Under this sense of root functions, necessary and sufficient conditions of unconditional basicity in  $L_2$  of the system conditions of unconditional basicity in  $L_2$  of the system of root functions of the operator  $Lu = -u'' + q(x)u$  were first set up by V. A. Il'in in [8]. Further, in the papers [2, 11, 13, 14, 15, 21, 28] these and other issues were studied for an ordinary differential operator of arbitrary order, and the criteria of Bessel property, Riesz property and unconditional basicity were set up. Criterion of Bessel property and unconditional basicity for the Dirac operator with the potential  $P(x) \in L_2(G)$  were set up in the paper [16]. The papers [3, 17, 18, 19, 20] were devoted to componentwise uniform equiconvergence on a compact, uniform convergence, the Riesz property of the system of root vector functions of the Dirac operator.

Rich references [1, 4, 5, 7, 12, 23, 24, 25, 26, 27, 29] deal with properties of basicity and other spectral properties of root vector-functions of the Dirac operator (with boundary conditions). In the paper [29] the Riesz basicity is set up in the case when the potential belongs to the class  $L_2$  and boundary conditions are separated. The

---

*Mathematics subject classification* (2010): 34L10, 34L20, 34B05, 34B24.

*Keywords and phrases:* Dirac operator, unconditional basicity, root vector-functions, Bessel property.

case with regular boundary conditions and with potential from the class  $L_2$  was studied in the paper [5]. The Riesz basicity from subspaces, and in the case of strongly regular boundary conditions the Riesz basicity is proved. The Dirac operator with a potential from  $L_p, p \geq 1$  was studied in the papers [26], [27] and the Riesz basicity was proved for the case of strongly regular boundary conditions while in the case of regular (but not strongly regular) boundary conditions, the Riesz basicity from subspaces is proved.

Let  $L_p^2(G), p \geq 1, G = (0, 2\pi)$  be the space of two-component vector-functions  $f(x) = (f_1(x), f_2(x))^T$  with the norm

$$\|f\|_p = \|f\|_{p,2} = \left\{ \int_G |f(x)|^p dx \right\}^{\frac{1}{p}} = \left\{ \int_G \left( \sum_{i=1}^n |f_i(x)|^2 \right)^{\frac{p}{2}} dx \right\}^{\frac{1}{p}}.$$

In the case  $p = \infty$  the norm is defined by the equality  $\|f\|_\infty = \|f\|_{\infty,2} = \sup_{x \in \bar{G}} \text{vrai} |f(x)|$ .

For  $f(x) \in L_p^2(G), g(x) \in L_q^2(G), p^{-1} + q^{-1} = 1, 1 \leq p \leq \infty$ , there exist  $(f, g) = \int_G \sum_{j=1}^2 f_j(x) \overline{g_j(x)} dx$ .

Let us consider one-dimensional Dirac operator

$$Dy = By' + P(x)y, \quad y(x) = (y_1(x), y_2(x))^T, \quad x \in G = (0, 2\pi),$$

where  $B = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, P(x) = \text{diag}(p(x), q(x)), p(x)$  and  $q(x)$  are complex-valued functions from the class  $L_1(G)$ .

Following [8] we will understand the root vector-functions of the operator  $D$  without regard to the form of boundary conditions, more exactly under eigen vector-function of the Dirac operator  $D$  responding to the eigen value  $\lambda$ , we will understand any identically nonzero two-component vector-function  $y^0(x)$  that is absolutely continuous on  $\bar{G}$  and almost everywhere in  $G$  satisfies the equation  $Dy^0 = \lambda y^0$ . Under the associated vector-function of order  $\ell, \ell \geq 1$  responding to the same  $\lambda$  and associated vector-function  $y^0(x)$ , we will understand any two-component vector-function  $y^\ell(x)$  that is absolutely continuous  $\bar{G}$  and almost everywhere in  $G$  satisfies the equation  $Dy^\ell = \lambda y^\ell + y^{\ell-1}$ .

Let  $\{u_k(x)\}_{k=1}^\infty$  be an arbitrary system comprised from the root vector-functions of the operator  $D, \{\lambda_k\}_{k=1}^\infty$  be the appropriate system of eigen values. Furthermore, the vector-function  $u_k(x)$  is included into the system  $\{u_k(x)\}_{k=1}^\infty$  together with appropriate associated vector-functions of less order.

DEFINITION 1.1. The system  $\{\varphi_k(x)\}_{k=1}^\infty \subset L_2^2(G)$  is said to be Bessel if there exists a constant  $M$  such that for any vector-function  $f(x) \in L_2^2(G)$  the equality

$$\sum_{k=1}^\infty |(f, \varphi_k)|^2 \leq M \|f\|_{2,2}^2$$

is fulfilled.

DEFINITION 1.2. The system  $\{\varphi_k(x)\}_{k=1}^\infty \subset L_2^2(G)$  is quadratically close to the system  $\{\psi_k(x)\}_{k=1}^\infty \subset L_2^2(G)$  if  $\sum_{k=1}^\infty \|\varphi_k - \psi_k\|_{2,2}^2 < \infty$ .

DEFINITION 1.3. Two sequences of elements in Hilbert space  $H$  are equivalent if there exists an operator  $H$  that is linear bounded and boundedly invertible in  $H$  and takes one of these sequences to another one.

In the paper the following theorems are proved.

THEOREM 1.4. (Criterion of Bessel property) *Let  $P(x) \in L_1(G)$ , the lengths of the chains of root vector-functions be uniformly bounded and there exist a constant  $C_0$  such that*

$$|Im\lambda_k| \leq C_0, \quad k = 1, 2, \dots \tag{1.1}$$

*Then for the Bessel property of the system  $\{u_k(x) \|u_k\|_2^{-1}\}_{k=1}^\infty$  in  $L_2^2(G)$  it is necessary and sufficient the existence of a constant  $M_1$  such that*

$$\sum_{|Re\lambda_k - \tau| < 1} 1 \leq M_1, \tag{1.2}$$

where  $\tau$  is an arbitrary real number.

REMARK 1.5. In sufficient part of theorem 1.4 the condition of uniform boundedness of the length of chains of root vector-functions is fulfilled in all cases, because it follows from inequality (1.2).

Let  $D^*$  be an operator formally associated to the operator  $D$ :

$$D^* = B \frac{d}{dx} + P^*(x).$$

Let  $\{\vartheta_k(x)\}_{k=1}^\infty$  be a system biorthogonally associated to  $\{u_k(x)\}_{k=1}^\infty$  in  $L_2^2(G)$  and consists of the root vector-functions of the operator  $D^*$ .

THEOREM 1.6. (On unconditional basicity) *Let  $P(x) \in L_1(G)$ , the lengths of the chains of root vector-functions be uniformly bounded, one of the systems  $\{u_k(x)\}_{k=1}^\infty$  and  $\{\vartheta_k(x)\}_{k=1}^\infty$  be complete in  $L_2^2(G)$  and condition (1.1) be fulfilled. Then necessary and sufficient condition for unconditional basicity in  $L_2^2(G)$  of each of these systems is the existence of constants  $M_1$  and  $M_2$  that ensure validity of inequality (1.2) and*

$$\|u_k\|_2 \|\vartheta_k\|_2 \leq M_2, \quad k = 1, 2, \dots \tag{1.3}$$

*Note that under the conditions of theorem 1.5 satisfaction of inequalities (1.2) and (1.3) is the necessary and sufficient condition for Riesz basicity of each of the systems  $\{u_k(x) \|u_k\|_2^{-1}\}_{k=1}^\infty$  and  $\{\vartheta_k(x) \|\vartheta_k\|_2^{-1}\}_{k=1}^\infty$  in  $L_2^2(G)$ .*

**THEOREM 1.7.** (On basicity) *Let  $P(x) \in L_1(G)$ , the lengths of the chains of root vector-functions be uniformly bounded, condition (1.1)–(1.3) be fulfilled and the system  $\left\{u_k(x) \|u_k\|_2^{-1}\right\}_{k=1}^\infty$  be quadratically close to some basis  $\{\psi_k(x)\}_{k=1}^\infty$  of the space  $L_2^2(G)$ . Then the systems  $\left\{u_k(x) \|u_k\|_2^{-1}\right\}_{k=1}^\infty$  and  $\{\vartheta_k(x) \|u_k\|_2\}_{k=1}^\infty$  are the bases in  $L_2^2(G)$ , and they are equivalent to the basis  $\{\psi_k(x)\}_{k=1}^\infty$  and its biorthogonally associated system, respectively.*

**REMARK 1.8.** Theorems 1.4, 1.5, and 1.6 remain valid also in the case when  $P(x) \in L_1(G)$  is an arbitrary but not necessarily a diagonal matrix-function.

### 2. Some auxiliary statements

Cite some necessary statements that will be used when proving theorems 1.4–1.6.

**STATEMENT 2.1.** (see [16]) *If  $p(x)$  and  $q(x)$  belong to the class  $L_1^{loc}(G)$  and the points  $x-t, x, x+t$  are in the domain  $G$ , then the following formulas are valid for the root vector-function  $u_k(x)$ :*

$$u_k(x \pm t) = (\cos \lambda_k t \cdot I \mp \sin \lambda_k t \cdot B) u_k(x) + \int_x^{x \pm t} (\sin \lambda_k (t - |\xi - x|) I \pm \cos \lambda_k (t - |\xi - x|) B) [P(\xi) u_k(\xi) - \theta_k u_{k-1}(\xi)] d\xi; \tag{2.1}$$

$$u_k(x-t) + u_k(x+t) = 2u_k(x) \cos \lambda_k t + \int_{x-t}^{x+t} (\sin \lambda_k (t - |\xi - x|) I + \operatorname{sgn}(\xi - x) \cos \lambda_k (t - |x - \xi|) B) [P(\xi) u_k(\xi) - \theta_k u_{k-1}(\xi)] d\xi, \tag{2.2}$$

where  $I$  is a unit operator in  $E^2$ ;  $\theta_k = 0$ , if  $u_k(x)$  is an eigen vector-function;  $\theta_k = 1$ , if  $u_k(x)$  is an associated vector-function, and in the last case it is assumed that  $\lambda_k = \lambda_{k-1}$ .

**STATEMENT 2.2.** (see [16]) *Let the functions  $p(x)$  and  $q(x)$  belong to the class  $L_1(G)$ . Then there exist the constants  $G_i(n_k, G)$ ,  $i = 1, 2$ , independent of  $\lambda_k$ , such that the following estimates are valid*

$$\|\theta_k u_{k-1}\|_\infty \leq C_1(n_k, G) (1 + |\operatorname{Im} \lambda_k|) \|u_k\|_\infty, \tag{2.3}$$

$$\|u_k\|_\infty \leq C_2(n_k, G) (1 + |\operatorname{Im} \lambda_k|)^{\frac{1}{r}} \|u_k\|_r, \tag{2.4}$$

where  $n_k$  is the order of the associated vector-function  $u_k$ ,  $r \geq 1$ .

### 3. Proof of the Bessel property criterion

In this section of the paper we will prove theorem 1.4 on Bessel property of root vector-function of Dirac’s operator.

*Necessity.* Let us fix an arbitrary number  $\tau \in (-\infty, +\infty)$  and introduce the indices set  $J_\tau = \{k : |\operatorname{Re}\lambda_k - \tau| \leq 1, |\operatorname{Im}\lambda_k| \leq C_0\}$ , where  $C_0$  is a constant from condition (1.1). Choose a number  $n_0 \geq 1$  so that  $R = \left(n_0(1 + C_0)^{\frac{3}{2}}\right)^{-1} \leq \frac{\operatorname{mes}G}{4} = \frac{\pi}{2}$  and for any set  $E \subset \bar{G}$ ,  $\operatorname{mes}E \leq 2R$ , the following inequality be fulfilled:

$$\omega(R) = \sup_{E \subset \bar{G}} \left\{ \|P\|_{1,E} \right\} \leq L^{-1}, \text{ where } \|P\|_{1,E} = \int_E \{|p(x)| + |q(x)|\} dx,$$

$L$  is a positive number and choice of its value will be defined later.

Let  $k \in J_\tau$ ,  $x \in [0, \pi]$ . Write the mean value formula (2.2) for the points  $x, x + t, x + 2t$ , where  $t \in [0, R]$ :

$$\begin{aligned} u_k(x) &= 2u_k(x + t) \cos \lambda_k t - u_k(x + 2t) + \int_x^{x+2t} (\sin \lambda_k(t - |x + t - \xi|)I \\ &\quad + \operatorname{sgn}(\xi - x - t) \cos \lambda_k(t - |x + t - \xi|)B) [P(\xi)u_k(\xi) - \theta_k u_{k-1}(\xi)] d\xi. \end{aligned}$$

Adding and subtracting  $2u_k(x + t) \cos \tau t$  in the right hand side of this equality, and using the operation  $R^{-1} \int_0^R dt$  we get

$$\begin{aligned} u_k(x) &= \frac{2}{R} \int_0^R u_k(x + t) \cos \tau t dt - \frac{1}{R} \int_0^R u_k(x + 2t) dt \\ &\quad + \frac{4}{R} \int_0^R u_k(x + t) \sin \frac{\lambda_k + \tau}{2} t \sin \frac{\tau - \lambda_k}{2} t dt \\ &\quad + \frac{1}{R} \int_0^R \int_x^{x+2t} (\sin \lambda_k(t - |x + t - \xi|)I \\ &\quad + \operatorname{sgn}(\xi - x - t) \cos \lambda_k(t - |x + t - \xi|)B) [P(\xi)u_k(\xi) - \theta_k u_{k-1}(\xi)] d\xi dt. \end{aligned}$$

Having used formula (2.1) in the third addend, we have

$$\begin{aligned} u_k(x) &= R^{-1} \int_0^{2\pi} u_k(t)w(t)dt + 4R^{-1} \int_0^R (\cos \lambda_k t I - \sin \lambda_k t B) \\ &\quad \times \sin \frac{\lambda_k + \tau}{2} t \sin \frac{\tau - \lambda_k}{2} t dt u_k(x) + 4R^{-1} \cdot \int_0^R \int_x^{x+t} (\sin \lambda_k(t - |x + t - \xi|)I \\ &\quad + \cos \lambda_k(t - |\xi - x|) B) \cdot [P(\xi)u_k(\xi) - \theta_k u_{k-1}(\xi)] d\xi \sin \frac{\tau + \lambda_k}{2} t \sin \frac{\tau - \lambda_k}{2} t dt \\ &\quad + R^{-1} \int_0^R \int_x^{x+2t} (\sin \lambda_k(t - |x + t - \xi|)I \\ &\quad + \cos \lambda_k(t - |x + t - \xi|)B) [P(\xi)u_k(\xi) - \theta_k u_{k-1}(\xi)] d\xi dt \\ &= R^{-1} \int_0^{2\pi} u_k(t)w(t) dt + I_1 + I_2 + I_3 \end{aligned} \tag{3.1}$$

where  $w(t) = 2 \cos \tau(x - t) - \frac{1}{2}$  for  $x \leq t \leq x + R$ ,  $w(t) = -\frac{1}{2}$  for  $x + R < t \leq x + 2R$  and  $w(t) = 0$  for  $t \notin [x, x + 2R]$ .

Taking  $k \in J_\tau$  into account and using the inequalities  $|\sin z| \leq 2$ ,  $|\cos z| \leq 2$  for  $|Imz| \leq 1$  and  $|\sin z| \leq 2|z|$  for  $|Imz| \leq 1$  we find

$$\begin{aligned} |I_1| &\leq 16R|\tau - \lambda_k| |u_k(x)| \leq 16R(1 + |Im\lambda_k|) |u_k(x)| \\ &\leq 16R(1 + C_0) |u_k(x)| \leq 16n_0^{-1} |u_k(x)|; \\ |I_2| &\leq 64 \left( \omega(R) \|u_k(x)\|_\infty + \frac{R}{2} \|\theta_k u_{k-1}\|_\infty \right); \\ |I_3| &\leq 8(\omega(R) \|u_k(x)\|_\infty + R \|\theta_k u_{k-1}\|_\infty). \end{aligned}$$

Taking these estimations into account in equality (3.1), we get the inequality

$$|u_k(x)| \leq R^{-1} \left| \int_0^{2\pi} u_k(t)w(t)dt \right| + 16n_0^{-1} |u_k(x)| + 72\omega(R) \|u_k\|_\infty + 40R \|\theta_k u_{k-1}\|_\infty.$$

Requiring  $n_0 \geq 32$ , hence we find that for  $x \in [0, \pi]$  it is valid

$$|u_k(x)| \leq 2R^{-1} \left| \int_0^{2\pi} u_k(t)w(t)dt \right| + 144\omega(R) \|u_k\|_\infty + 80R \|\theta_k u_{k-1}\|_\infty. \tag{3.2}$$

In the same way, inequality (3.2) is proved in the case  $x \in [\pi, 2\pi]$ . In this case the function  $w(t)$  is defined in the following way:  $w(t) = -\frac{1}{2}$  for  $x - 2R \leq t < x - R$ ,  $w(t) = 2\cos v(x - t) - \frac{1}{2}$  for  $x - R \leq t \leq x$ ,  $w(t) = 0$  for  $t \in [x - 2R, x]$ . Therefore, inequality (3.2) is fulfilled for any  $x \in [0, 2\pi]$ . Having used estimations (2.3) and (2.4), allowing for  $1 + |Im\lambda_k| \leq 1 + C_0$ , from inequality (3.2) we have

$$\begin{aligned} |u_k(x)| &\leq 2R^{-1} \left| \int_0^{2\pi} u_k(t)w(t)dt \right| + \left[ 144C_2(n_k, G)(1 + C_0)^{\frac{1}{2}} \omega(R) \right. \\ &\quad \left. + 80R\theta_k C_1(n_k, G)c_2(n_k, G)(1 + C_0)^{\frac{3}{2}} \right] \|u_k\|_2. \end{aligned}$$

By uniform boundedness of the length of the chain of the root vectors  $C_2(n_k, G) \leq \gamma_1 = const$ ,  $C_1(n_k, G)C_2(n_k, G) \leq \gamma_2 = const$ . Consequently,

$$|u_k(x)| \leq 2R^{-1} \left| \int_0^{2\pi} u_k(t)w(t)dt \right| + \left[ 144\gamma_1(1 + C_0)^{\frac{1}{2}}L^{-1} + 80\theta_k\gamma_2n_0^{-1} \right] \|u_k\|_2$$

Hence, by inequalities  $|\sum_{i=1}^m a_i|^2 \leq m\sum_{i=1}^m |a_i|^2$  we have

$$\begin{aligned} |u_k(x)|^2 \|u_k\|_2^{-2} &\leq \frac{16}{R^2} \left\{ \left| \int_0^{2\pi} u_k^1(t)w(t)dt \right|^2 \|u_k\|_2^{-2} + \left| \int_0^{2\pi} u_k^2(t)w(t)dt \right|^2 \|u_k\|_2^{-2} \right\} \\ &\quad + 4 \left\{ \left( 144\gamma_1(1 + C_0)^{\frac{1}{2}}L^{-1} \right)^2 + \left( 80\theta_k\gamma_2n_0^{-1} \right)^2 \right\}, \end{aligned}$$

where  $u_k(t) = (u_k^1(t), u_k^2(t))^T$ .

Having applied the Bessel inequality, we get that for any finite set  $J \subset J_\tau$  the following inequality is fulfilled

$$\sum_{k \in J} |u_k(x)|^2 \|u_k\|_2^{-2} \leq \frac{32M}{R^2} \|w\|_2^2 + 4 \left\{ \left( 144\gamma_1(1 + C_0)^{\frac{1}{2}}L^{-1} \right)^2 + (80\theta_k\gamma_2n_0^{-1})^2 \right\} \sum_{k \in J} 1. \tag{3.3}$$

Taking into account  $\|w\|_2^2 = O(R)$  and choosing the number  $n_0$  so large that

$$4 \left\{ \left( 144\gamma_1(1 + C_0)^{\frac{1}{2}}L^{-1} \right)^2 + (80\gamma_2n_0^{-1})^2 \right\} \leq \frac{1}{4\pi}$$

be fulfilled, from (3.3) we find

$$\sum_{k \in J} 1 \leq \text{const}R^{-1} = \text{const}n_0(1 + C_0)^{\frac{3}{2}} = \text{const}.$$

By arbitrariness of the set  $J \subset J_\tau$ , hence we get inequality (1.2). The part of necessity of theorem 1.1 is proved.

*Sufficiency.* Writing the shift formula (2.1) for  $u_k(x+t)$  for  $x = 0$  and scalarly multiplying by the vector-function  $f(t) = (f_1(t), f_2(t))^T \in L_2^2(0, 2\pi)$ , we conclude that for the Bessel inequality to be fulfilled for the system  $\varphi_k(t) = u_k(t) \|u_k\|_2^{-1}$ ,  $k = 1, 2, \dots$ , it suffices to set up validity of the following inequalities:

$$\sum_{k=1}^{\infty} \left| \int_0^{2\pi} \overline{f_i(t)} \cos \lambda_k t dt \right|^2 |\varphi_k^i(0)|^2 \leq C \|f\|_2^2, \quad i = 1, 2; \tag{3.4}$$

$$\sum_{k=1}^{\infty} \left| \int_0^{2\pi} \overline{f_i(t)} \sin \lambda_k t dt \right|^2 |\varphi_k^{3-i}(0)|^2 \leq C \|f\|_2^2, \quad i = 1, 2; \tag{3.5}$$

$$\sum_{k=1}^{\infty} \left| \int_0^{2\pi} \overline{f_1(t)} \int_0^t p(\xi) \varphi_k^1(\xi) \sin \lambda_k(t - \xi) d\xi dt \right|^2 \leq C \|f\|_2^2; \tag{3.6}$$

$$\sum_{k=1}^{\infty} \left| \int_0^{2\pi} \overline{f_1(t)} \int_0^t q(\xi) \varphi_k^2(\xi) \cos \lambda_k(t - \xi) d\xi dt \right|^2 \leq C \|f\|_2^2; \tag{3.7}$$

$$\sum_{k=1}^{\infty} \left| \int_0^{2\pi} \overline{f_2(t)} \int_0^t p(\xi) \varphi_k^1(\xi) \cos \lambda_k(t - \xi) d\xi dt \right|^2 \leq C \|f\|_2^2; \tag{3.8}$$

$$\sum_{k=1}^{\infty} \left| \int_0^{2\pi} \overline{f_2(t)} \int_0^t q(\xi) \varphi_k^2(\xi) \sin \lambda_k(t - \xi) d\xi dt \right|^2 \leq C \|f\|_2^2; \tag{3.9}$$

$$\sum_{k=1}^{\infty} \left| \theta_k \int_0^{2\pi} \overline{f_i(t)} \int_0^t \frac{u_{k-1}^i(\xi)}{\|u_k\|_2} \sin \lambda_k(t - \xi) d\xi dt \right|^2 \leq C \|f\|_2^2, \quad i = 1, 2; \tag{3.10}$$

$$\sum_{k=1}^{\infty} \left| \theta_k \int_0^{2\pi} \overline{f_i(t)} \int_0^t \frac{u_{k-1}^{3-i}(\xi)}{\|u_k\|_2} \cos \lambda_k(t - \xi) d\xi dt \right|^2 \leq C \|f\|_2^2, \quad i = 1, 2; \tag{3.11}$$

where  $\varphi_k^i(\xi) = u_k^i(\xi) \|u_k\|_2^{-1}$ .

Prove estimation (3.4). As by estimation (2.4) and conditions (1.1), (1.2)

$$\begin{aligned} |\varphi_k^i(0)| &= |u_k^i(0)| \|u_k\|_2^{-1} \leq \|u_k\|_\infty \|u_k\|_2^{-1} \\ &\leq C_2(n_k, G) (1 + C_0)^{\frac{1}{2}} \|u_k\|_2 \|u_k\|_2^{-1} = C_2(n_k, G) (1 + C_0)^{\frac{1}{2}} \leq const, \end{aligned}$$

as by (1.2) the quantity  $C_2(n_k, G)$  is bounded, then for validity of (3.4) it is sufficient the following inequality to be fulfilled

$$\sum_{k=1}^{\infty} \left| \int_0^{2\pi} \overline{f_1(t)} \cos \lambda_k t dt \right|^q \leq C \|f\|_2^2. \tag{3.12}$$

This inequality was set up in the paper [8] provided  $Re\lambda_k \geq 0$ , (1.1) and (1.2) for  $\tau \geq 1$ . Hence inequality (3.12) follows for  $Re\lambda_k \in (-\infty, +\infty)$ ,  $|Im\lambda_k| \leq C_0$ , as by the theorem condition (1.2) is fulfilled for any  $\tau \in (-\infty, +\infty)$ . Inequality (3.5) is set up just in the same way.

Now let us be convinced in validity of inequalities (3.6)–(3.9). They are proved in the same way. Therefore we prove only the inequality (3.6). Denote

$$g_i(t, \xi) = \begin{cases} f_i(t + \xi) & \text{for } 0 \leq t \leq 2\pi - \xi \\ 0, & \text{for } 2\pi - \xi < t \leq 2\pi, \end{cases} \quad i = 1, 2,$$

where  $\xi \in [0, 2\pi]$ . Then by estimation (2.4) and conditions (1.1), (1.2), we have

$$\begin{aligned} I_k &= \left| \int_0^{2\pi} \overline{f_1(t)} \int_0^t p(\xi) \varphi_k^1(\xi) \sin \lambda_k(t - \xi) d\xi dt \right|^2 \\ &= \int_0^{2\pi} \overline{f_1(t)} \int_0^t p(\xi) \varphi_k^1(\xi) \sin \lambda_k(t - \xi) d\xi dt \\ &\quad \times \int_0^{2\pi} f_1(t) \int_0^t \overline{p(\xi) \varphi_k^1(\xi) \sin \lambda_k(t - \xi)} d\xi dt \\ &= \int_0^{2\pi} p(\xi) \varphi_k^1(\xi) \int_0^{2\pi} g_1(t, \xi) \sin \lambda_k t dt d\xi \\ &\quad \times \int_0^{2\pi} \overline{p(\tau) \varphi_k^1(\tau)} \int_0^{2\pi} g_1(r, \tau) \overline{\sin \lambda_k r} dr d\tau \\ &= \int_0^{2\pi} \int_0^{2\pi} p(\xi) \overline{p(\tau)} \varphi_k^1(\xi) \overline{\varphi_k^1(\tau)} \int_0^{2\pi} \overline{g_1(t, \xi)} \\ &\quad \times \sin \lambda_k t dt \cdot \int_0^{2\pi} g_1(r, \tau) \overline{\sin \lambda_k r} dr d\xi d\tau \\ &\leq C_2^2(n_k, G) (1 + C_0) \int_0^{2\pi} \int_0^{2\pi} |p(\xi)| |p(\tau)| \\ &\quad \times \left| \int_0^{2\pi} \overline{g_1(t, \xi)} \sin \lambda_k t dt \right| \left| \int_0^{2\pi} g_1(r, \tau) \overline{\sin \lambda_k r} dr \right| d\xi d\tau \\ &\leq const \int_0^{2\pi} \int_0^{2\pi} |p(\xi)| |p(\tau)| \left| \int_0^{2\pi} \overline{g_1(t, \xi)} \sin \lambda_k t dt \right| \left| \int_0^{2\pi} g_1(r, \tau) \overline{\sin \lambda_k r} dr \right| d\xi d\tau. \end{aligned}$$



Hence for an arbitrary natural number  $N$  we have

$$\begin{aligned} \sum_{k=1}^N I_k &\leq \text{const} \int_0^{2\pi} \int_0^{2\pi} |p(\xi)| |p(\tau)| \\ &\quad \times \left( \sum_{k=1}^N \left| \int_0^{2\pi} g_1(t, \xi) \sin \lambda_k t dt \right| \left| \int_0^{2\pi} g_1(r, \tau) \sin \lambda_k r dr \right| \right) d\xi d\tau \\ &\leq \text{const} \int_0^{2\pi} \int_0^{2\pi} |p(\xi)| |p(\tau)| \|g_1(\cdot, \xi)\|_2 \|g_1(\cdot, \tau)\|_2 d\xi d\tau. \end{aligned}$$

Taking into account that  $\|g_1(\cdot, \xi)\|_2 \leq \|f_1\|_2$  is fulfilled for any fixed  $\xi \in [0, 2\pi]$ , we get  $\sum_{k=1}^N I_k \leq \text{const} \|p\|_1^2 \|f_1\|_2^2 \leq \text{const} \|f\|_2^2$ .

Hence, the validity of inequality (3.5) follows from arbitrariness of the number  $N$ . Prove inequality (3.10). By estimations (2.3), (2.4) and conditions (1.1), (1.2)

$$\begin{aligned} \theta_k |u_{k-1}^i(\xi)| \|u_k\|_2^{-1} &\leq \theta_k C_1(n_k, G) C_2(n_k, G) (1 + C_0)^{\frac{3}{2}} \\ \|u_k\|_2 \|u_k\|_2^{-1} &= \theta_k C_1(n_k, G) C_2(n_k, G) \leq C = \text{const}. \end{aligned}$$

After changing the order of integration, the left hand side of (3.10) with regard to the last inequality is majorized from above with the series

$$C \sum_{k=1}^{\infty} \int_0^{2\pi} \left| \int_0^{2\pi} g_i(t, \xi) \sin \lambda_k t dt \right|^2 d\xi.$$

This series converges and its some does not exceed the value  $C \|f\|_2^2$ . The validity of inequality (3.10) is proved. Inequality (3.11) is proved just in the same way. Theorem 1.4 is completely proved.  $\square$

### 4. Proof of theorems 1.2 and 1.3

*Proof of theorem 1.5. Necessity.* Let the systems  $\{u_k(x)\}_{k=1}^{\infty}$  and  $\{\vartheta_k(x)\}_{k=1}^{\infty}$  form an unconditional basis in  $L_2^2(G)$ . According to Lorch's theorem [22] the systems,  $\{u_k(x) \|u_k\|_2^{-1}\}_{k=1}^{\infty}$  and  $\{\vartheta_k(x) \|\vartheta_k\|_2^{-1}\}_{k=1}^{\infty}$  are the Riesz basis in  $L_2^2(G)$ . Then by N. K. Barri's theorem [6, pp. 347–375] the series

$$\sum_{k=1}^{\infty} |(f, u_k)|^2 \|u_k\|_2^{-2}, \quad \sum_{k=1}^{\infty} |(g, \vartheta_k)|^2 \|\vartheta_k\|_2^{-2}$$

converge for any  $f(x), g(x)$  from the class  $L_2^2(G)$ . Since the Bessel property of the systems  $\{u_k(x) \|u_k\|_2^{-1}\}_{k=1}^{\infty}$  and  $\{\vartheta_k(x) \|\vartheta_k\|_2^{-1}\}_{k=1}^{\infty}$  follows from convergence of these series (see [9], pp. 433–435), then necessity of condition (1.2) follows from theorem 1.4. The necessity of condition (1.3) is known well not only for unconditional but for ordinary basicity of the arbitrary system  $\{u_k(x)\}_{k=1}^{\infty}$  as well (see [6], p. 370).

*Sufficiency.* Let conditions (1.1)–(1.3) be fulfilled, and the system  $\{u_k(x)\}$  be complete in  $L^2_2(G)$ . By theorem 1.4, the systems  $\{u_k(x) \|u_k\|_2^{-1}\}_{k=1}^\infty$  and  $\{\vartheta_k(x) \|\vartheta_k\|_2^{-1}\}_{k=1}^\infty$  are Bessel in  $L^2_2(G)$ .

As conditions (1.3) are fulfilled, then for arbitrary  $g(x) \in L^2_2(G)$

$$\sum_{k=1}^\infty |(g, \vartheta_k \|u_k\|_2)|^2 \leq M_2 \sum_{k=1}^\infty |(g, \vartheta_k \|\vartheta_k\|_2^{-1})|^2 \leq \text{const} \|g\|_2^2.$$

Therefore, the systems  $\{u_k(x) \|u_k\|_2^{-1}\}_{k=1}^\infty$  and  $\{\vartheta_k(x) \|u_k\|_2\}_{k=1}^\infty$  are Bessel in  $L^2_2(G)$ .

Prove the unconditional basicity of the system  $\{u_k(x)\}_{k=1}^\infty$ . Let  $P_n f = \sum_{k=1}^n (f, \vartheta_k) u_k(x)$ ,  $n = 1, 2, \dots$ . Then

$$\begin{aligned} \|P_n f\|_2 &= \sup_{\|g\|_2=1} |(P_n f, g)| = \sup_{\|g\|_2=1} \left| \sum_{k=1}^n (f, \vartheta_k) (u_k, g) \right| \\ &\leq \sup_{\|g\|_2=1} \sum_{k=1}^n |(f, \vartheta_k \|u_k\|_2)| \left| (g, u_k \|u_k\|_2^{-1}) \right| \leq \text{const} \|f\|_2. \end{aligned}$$

Consequently, the sequence  $\{P_n f\}$  is uniformly bounded. Then by the theorem on bases (see [10, p. 11]), the system  $\{u_k(x)\}_{k=1}^\infty$  is a basis in  $L^2_2(G)$ . It will be an unconditional basis as well by the Bessel property of the systems  $\{u_k(x) \|u_k\|_2^{-1}\}_{k=1}^\infty$  and  $\{\vartheta_k(x) \|u_k\|_2\}_{k=1}^\infty$  in  $L^2_2(G)$ . The system  $\{\vartheta_k(x)\}_{k=1}^\infty$  is also an unconditional basis in  $L^2_2(G)$ , because it is a system biorthogonally associated to  $\{u_k(x)\}_{k=1}^\infty$ . Theorem 1.5 is proved.  $\square$

*Proof of theorem 1.6.* As the system  $\{\vartheta_k(x)\}_{k=1}^\infty$  consists of root vector-functions of the operator  $D^* = B \frac{d}{dx} + P^*(x)$ , then by theorem 1.4 conditions (1.1) and (1.2) provide Bessel property of the system  $\{\vartheta_k(x) \|\vartheta_k\|_2^{-1}\}_{k=1}^\infty$  in  $L^2_2(G)$ , i.e.

$$\sum_{k=1}^\infty |(f, \vartheta_k \|\vartheta_k\|_2^{-1})|^2 \leq M \|f\|_2^2 \tag{4.1}$$

for any vector-function  $f(x) \in L^2_2(G)$ .

From inequality (4.1), condition (1.3) and quadratical closeness of the system  $\{u_k \|u_k\|_2^{-1}\}_{k=1}^\infty$  and  $\{\psi_k\}_{k=1}^\infty$  it follows that the series

$$\sum_{k=1}^\infty \tilde{f}_k \|u_k\|_2 \|\vartheta_k\|_2 \left( u_k(x) \|u_k\|_2^{-1} - \psi_k(x) \right)$$

converges in  $L^2_2(G)$  for any  $f(x) \in L^2_2(G)$ , where  $\tilde{f}_k = (f, \vartheta_k \|\vartheta_k\|_2^{-1})$ . Let us denote the sum of this series by  $Kf$ , where  $K$  is some linear operator acting in  $L^2_2(G)$ . This

follows from fundamental property of the sequence

$$K_n f = \sum_{k=1}^n \tilde{f}_k \|u_k\|_2 \|\vartheta_k\|_2 \left( u_k(x) \|u_k\|_2^{-1} - \psi_k(x) \right)$$

in the space  $L_2^2(G)$ .

Obviously, the relation  $\|Kf - K_n f\|_2 = o(1) \|f\|_2$  is fulfilled for any  $f(x) \in L_2^2(G)$ . Consequently, the sequence of finite-dimensional operators  $K_n$  converges to the operator  $K$ , i.e.  $\|K - K_n\|_{L_2^2 \rightarrow L_2^2} \rightarrow 0$   $n \rightarrow \infty$ . Hence, it follows the compactness of the operator  $K$  in  $L_2^2(G)$ .

It is clear that  $Ku_k \|u_k\|_2^{-1} = u_k \|u_k\|_2^{-1} - \psi_k$ , i.e.  $(I - K)u_k \|u_k\|_2^{-1} = \psi_k$ ,  $k = 1, 2, \dots$

Show that the operator  $I - K$  is continuously invertible. As  $K$  is a compact operator, then from the Fredholm alternative it follows that if the operator  $I - K$  is irreversible, then there exists a nonzero element  $g \in L_2^2(G)$  such that  $(I - K)^* g = 0$ . This element  $g$  satisfies the relation

$$(g, \psi_k) = (g, (I - K)u_k \|u_k\|_2^{-1}) = ((I - K)^* g, u_k \|u_k\|_2^{-1}) = 0, \quad k = 1, 2, \dots$$

Hence, by the basicity of the system  $\{\psi_k(x)\}_{k=1}^\infty$  in  $L_2^2(G)$  it follows that  $g \equiv 0$ . The obtained contradiction shows that the operator  $I - K$  is reversible. Consequently, the system  $\{u_k(x) \|u_k\|_2^{-1}\}_{k=1}^\infty$  is a basis in  $L_2^2(G)$  equivalent to the basis  $\{\psi_k(x)\}_{k=1}^\infty$ .

If by  $\{z_k(x)\}_{k=1}^\infty$  we denote a system biorthogonal to  $\{\psi_k(x)\}_{k=1}^\infty$ , then  $\vartheta_k(x) \|u_k\|_2 = (I - K)^* z_k(x)$ . Hence it follows that the system  $\{\vartheta_k(x) \|u_k\|_2\}_{k=1}^\infty$  is a basis equivalent to the basis  $\{z_k(x)\}_{k=1}^\infty$  in  $L_2^2(G)$ .

Theorem 1.6 is proved.  $\square$

## REFERENCES

- [1] T. SH. ABDULLAEV, I. M. NABIEV, *An algorithm for reconstructing the Dirac operator with a spectral parameter in the boundary condition*, Com. Math. and Math. Phys. **56** (2016), (2), 256–257.
- [2] V. D. BUDAEV, *Criteria for the Bessel property and Riesz basis property of systems of root functions of differential operator I*, Differ. Uravn. **27**, (1991), no. 12, 2033–2044.
- [3] L. Z. BUKSAYEVA, *Necessary conditions of Riesz property of root vector-functions of Dirac discontinuous operator with summable coefficient*, Pros. IMM of NAS **42** (2016), no. 1, 106–115.
- [4] P. DJAKOV, B. MITYAGIN, *Criteria for existence of Riesz basis consisting of root functions of Hill and 1D Dirac operators*, J. Funct. Anal. **263** (2012), 2300–2332.
- [5] P. DJAKOV, B. MITYAGIN, *Unconditional convergence of spectral decompositions of 1D Dirac operators with regular boundary conditions*, Indiana Univ. Math. J. **61** (2012), no. 1, 359–398.
- [6] I. T. HOCHBERG, M. G. KREIN, *Introduction to theory of linear non self-adjoint operators in Hilbert space*, M. Nauka, 1965 (in Russian).
- [7] S. HASSI, L. L. ORIDOROQA, *Completeness and Riesz basis property of systems of eigen-functions and associated functions of Dirac-type operators with boundary conditions depending of the spectral paramet*, Math. Notes **79** (2016), (3) 589–593.
- [8] V. A. IL'IN, *Unconditional basis property on a closed interval of systems of eigen-and associated functions of a second order differential operator*, Dokl. Akad. Nauk SSSR **273** (1983), no. 5, 1048–1053.

- [9] S. KACZMARZ, H. STEINHAUS, *Theory of orthogonal series*, 1958 (in Russian).
- [10] B. S. KASHIN, A. A. SAAKYAN, *Orthogonal series*, Trans. of Math. Monographs, AMS **75** (1989).
- [11] N. B. KERIMOV, *Unconditional basis property of a system of eigen- and associated functions of a fourth-order differential operator*, Dokl. Akad. Nauk SSSR **286** (1986), no. 4, 803–808.
- [12] V. V. KORNEV, A. P. KHROMOV, *Dirac system with undifferentiable potential and antiperiodic boundary conditions*, Iz. Saratov Univ. Nov. Ser. Math. Mech. Inform. **13** (2013), 28–35.
- [13] L. V. KRITSKOV, A. M. SERSENBİ, *Basicity in  $L_p$  of root functions for differential equations with involution*, Electron. J. Differ. Equ. **2015** 278 (2015), 1–9.
- [14] V. M. KURBANOV, *On the distribution of eigenvalues and a criterion for the Bessel property of root functions of a differential operator I*, Differ. Uravn. **41** (2005), no. 4, 464–478.
- [15] V. M. KURBANOV, *On the distribution of eigenvalues and a criterion for the Bessel property of root functions of a differential operator II*, Differ. Uravn. **41** (2005), no. 5, 623–631.
- [16] V. M. KURBANOV, *The Bessel property and the unconditional basis property of systems, of root vector functions of the Dirac operator*, Differ. Equ. **32** (1996), no. 12, 1601–1610.
- [17] V. M. KURBANOV, A. I. ISMAILOVA, *Componentwise uniform eqiconvergence of expansions in the root vector functions of the Dirac operator with the trigonometric expansion*, Differ. Equ. **48** (2012), no. 5, 655–669.
- [18] V. M. KURBANOV, A. I. ISMAILOVA, *Absolute and uniform convergence of expansions in the root vector functions of the Dirac operator*, Dokl. Math. **86** (2012), no. 2, 663–666.
- [19] V. M. KURBANOV, A. I. ISMAILOVA, *Riesz inequality for systems of root vector functions of the Dirac operator*, Differ. Equ. **48** (2012), no. 3, 336–342.
- [20] V. M. KURBANOV, A. I. ISMAILOVA, *Two-sided estimates for root vector functions of the Dirac operator*, Differ. Equ. **48** (2012), no. 4, 494–505.
- [21] I. S. LOMOV, *The Bessel inequality, the Riesz theorem, and the unconditional Bazis property for root vectors of ordinary differential operators*, Vestnik Moskov. Univ., ser 1, Mat. Mekh., (1992), no. 5, 42–52.
- [22] E. R. LORCH, *Bicontinuous linear transformations in certain vector spaces*, Bull Amer. Math. Soc. **45**, (1939), no. 8, 564–569.
- [23] A. LUNYOV, M. M. MALAMUD, *On the Riesz basis property of the root vector system for Dirac-type  $2 \times 2$  systems*, Docl. Math. **90**, (2) (2014), 556–561.
- [24] A. A. LUNYOV, M. M. MALAMUD, *On the Riesz basis property of the root vector system for Dirac-type  $2 \times 2$  systems*, J. Math. Anal. and Apl. **441**, (1) (2016), 57–103.
- [25] R. MAMEDOV, O. AKCHAY, *Inverse eigenvalue problem for a class of Dirac operators with discontinuous coefficient*, Bound Value Probl. (2014) 2014; 110, doi:10.1186/1687-2770-2014-110.
- [26] A. M. SAVCHUK, SHKALIKOV, *The Dirac operator with complex-valued Summable potential*, Math. Notes **96**, (2014), no. 5, 3–36.
- [27] A. M. SAVCHUK, SADOVNICHAYA, *Riesz basis property with parentheses for Dirac system with integrable potential*, Sovrem. Math. Fundam. Napravl. **58** (2015), 128–152.
- [28] A. M. SERSENBİ, *Criteria for the Riesz basis property of systems of eigen and associated functions of higher-order differential operators on an interval*, Dokl. Akad. Nauk **419** (2008), no. 5, 601–603.
- [29] I. TROOSHIN, M. YAMAMOTA, *Riesz basis of root vectors of a nonsymmetric system of first-order ordinary differential operators and application to inverse eigenvalue problems*, Appl. Anal. **80** (2001), no. 1–2, 19–51.

(Received August 12, 2017)

Vali M. Kurbanov  
 Institute of Mathematics and Mechanics  
 of NAS of Azerbaijan  
 Baku, Azerbaijan  
 e-mail: q.valii@yahoo.com

Afsana M. Abdullayeva  
 Azerbaijan State Pedagogical University  
 Baku, Azerbaijan  
 e-mail: ef.abdullayeva@inbox.ru