

A FUNCTIONAL DECOMPOSITION OF FINITE BANDWIDTH REPRODUCING KERNEL HILBERT SPACES

GREGORY T. ADAMS AND NATHAN A. WAGNER

(Communicated by V. Bolotnikov)

Abstract. In this work, we consider “finite bandwidth” reproducing kernel Hilbert spaces which have orthonormal bases consisting of certain polynomials. We provide general conditions based on a matrix recursion that guarantee such spaces contain a functional multiple of the Hardy space. In a particular case, we obtain an explicit functional decomposition of these spaces that greatly generalizes a previous result in the tridiagonal case due to Adams and McGuire. We also prove that multiplication by z is a bounded operator on these spaces and that they contain the polynomials.

1. The problem

If $K(z, w)$ is a function defined on an open disc about the origin which is analytic in z and coanalytic in w , then K has a power series representation $K(z, w) = \sum_{j=0}^{\infty} \sum_{k=0}^{\infty} a_{j,k} z^j \bar{w}^k$. In the case that $A = (a_{j,k})$ is a bounded matrix, it is an easy exercise to check that A is positive semi-definite on ℓ^2 if and only if the function K is, and in this case by the Moore-Aronszajn Theorem the function K is the kernel for a reproducing kernel Hilbert space $H(K)$ (see [4]). In this case, the space $H(K)$ consists of analytic functions on a domain containing a disk about the origin in \mathbb{C} . Recall the well-known fact that if $\{f_n\}$ is an orthonormal basis for the reproducing kernel Hilbert space (RKHS) of functions $H(K)$ associated with K , then $K(z, w) = \sum_{n=0}^{\infty} f_n(z) \overline{f_n(w)}$ [7]. Conversely, if A can be factored as $A = LL^*$ where L has no kernel, then the columns of L give the Taylor coefficients of an orthonormal basis for $H(K)$ [1]. In fact, $H(K)$ can be identified with the range space of L in a very natural way [1]. This range space identification will lie at the heart of most of our computations.

The Cholesky algorithm always allows for a factorization of a positive semi-definite matrix $A = LL^*$ with L lower triangular. If A has finite bandwidth $2J + 1$, then L is lower triangular with $J + 1$ non-trivial diagonals and we speak of a “bandwidth- $2J + 1$ ” kernel K . In particular, we say an analytic kernel K is of finite bandwidth- $2J + 1$ if there exists an orthonormal basis of polynomials for $H(K)$ of the form

$$\{f_n(z) = (b_{0,n} + b_{1,n}z + \dots + b_{J,n}z^J)z^n\}.$$

Mathematics subject classification (2020): 46E22, 47B32.

Keywords and phrases: Reproducing kernel Hilbert spaces, multiplication operators, Douglas’ Range Inclusion Lemma, Hardy space.

The simplest case where the space $H(K)$ has bandwidth 1 was extensively studied by Shields in [8] in the context of multiplication operators. Such spaces are referred to as *diagonal spaces* and have orthonormal bases consisting of monomials.

In the context of bandwidth- $2J + 1$ analytic kernels, the *natural domain* of $H(K)$ is given by $\text{Dom}(K) = \{z \in \mathbb{C} : \sum_{n=0}^{\infty} |f_n(z)|^2 < \infty\}$. Adams and McGuire established that the natural domain for $H(K)$ is a disk about the origin with up to J additional points [2]. They explored the $J = 1$ case and gave an interesting family of kernels K where $H(K)$ is a nontrivial extension of a diagonal space [3]. In this paper, we show how to generalize their results to higher bandwidths.

Now we can state the problem of interest. Throughout this work, z_1, z_2, \dots, z_J will be distinct points on the unit circle \mathbb{T} and w_1, w_2, \dots, w_J will be the corresponding conjugates. The sequence of complex numbers a_0, a_1, \dots will be a sequence converging to 1 so that $1 - a_j$ is nonvanishing. Define

$$\phi(z) = \prod_{j=1}^J (1 - w_j z) = \sum_{k=0}^J \beta_k z^k,$$

and $f_n(z) = z^n \phi(a_n z)$. We will follow the notational convention that $\beta_j = 0$ if $j < 0$ or $j > J$. Then

$$K(z, w) = \sum_{n=0}^{\infty} f_n(z) \overline{f_n(w)}$$

is a bandwidth- $2J + 1$ kernel for a RKHS $H(K)$ with orthonormal basis $\{f_0, f_1, \dots\}$.

Theorems 3.4 and 3.9 show that in the case where $\lim_{n \rightarrow \infty} n(1 - a_n) = p$ and $p > 1/2$, $H(K)$ has natural domain $\mathcal{D} = \mathbb{D} \cup \{z_1, z_2, \dots, z_J\}$ and decomposes as

$$H(K) = \phi(z)H^2(\mathbb{D}) + \mathbb{C}K(z, z_1) + \mathbb{C}K(z, z_2) + \dots + \mathbb{C}K(z, z_J).$$

Moreover, in this case, multiplication by z is a bounded operator and the polynomials are contained in $H(K)$.

These results generalize those in [3] and [9] to higher bandwidth and more general weight sequences. This leads to a very nice functional characterization of certain finite bandwidth spaces. The primary innovation in this work is the use of matrix recursion to bound the norm of infinite dimensional matrices, a program which was started in [9]. Key also is the role played by the combinatorial Theorems 4.2 and 4.3.

2. Preliminaries

The first result shows that the restrictions of the functions in $H(K)$ to the disc \mathbb{D} are in the Hardy space.

PROPOSITION 2.1. $H(K) \subset H^2(\mathbb{D})$.

Proof. If $f \in H(K)$, then there exists an ℓ^2 sequence $\{\alpha_n\}$ such that $f = \sum_{n=0}^\infty \alpha_n f_n$. Thus, treating any variables with negative subscripts as 0:

$$\begin{aligned} f(z) &= \sum_{n=0}^\infty \alpha_n f_n(z) \\ &= \sum_{n=0}^\infty \alpha_n \left(\sum_{k=0}^J \beta_k a_n^k z^{n+k} \right) \\ &= \sum_{n=0}^\infty \left(\sum_{k=0}^J \alpha_{n-k} \beta_k a_{n-k}^k \right) z^n \\ &= \sum_{n=0}^\infty \widehat{\alpha}_n z^n. \end{aligned}$$

By the Cauchy-Schwarz inequality, $|\widehat{\alpha}_n|^2 \leq c^2 \sum_{k=0}^J |\alpha_{n-k}|^2$, where c is a constant that depends only on the zeros z_1, z_2, \dots, z_J and the sequence $\{a_n\}$ (which of course is bounded). In fact, we can take

$$c^2 = (J + 1) \max_{0 \leq k \leq J} |\beta_k|^2 \max_{0 \leq k \leq J} \|\{a_n\}\|_{\ell^\infty}^{2k}.$$

Thus, $\sum_{n=0}^\infty |\widehat{\alpha}_n|^2 \leq (J + 1)c^2 \sum_{n=0}^\infty |\alpha_n|^2$ and f is in $H^2(\mathbb{D})$. \square

Given the basis $f_n(z) = \phi(a_n z) z^n$ and the fact that $a_n \rightarrow 1$ it is reasonable to ask when functions of the form $\phi(z)f(z)$ for $f \in H^2(\mathbb{D})$ are in $H(K)$. The rate of convergence of a_n to 1 is crucial in assessing when this is the case. Douglas' Range Inclusion Lemma (see [6]) will provide the major tool to answer this question.

To this end, let L be the matrix whose n th column consists of the Taylor coefficients of $f_n(z)$ and let \widehat{L} be the matrix whose n th column consists of the Taylor coefficients of $z^n \phi(z)$. By Douglas' Lemma, $\phi(z)H^2(\mathbb{D}) \subset H(K)$ if and only if there is a bounded matrix $C = (c_{j,k})_{j,k \geq 0}$ such that $\widehat{L} = LC$. Solving this equation for C is complicated and will involve a recursion. First note that L and \widehat{L} are both lower triangular which implies that C is as well. So one must solve

$$\begin{pmatrix} \beta_0 & 0 & 0 & \cdots \\ \beta_1 & \beta_0 & 0 & \cdots \\ \beta_2 & \beta_1 & \beta_0 & \ddots \\ \vdots & \vdots & \vdots & \ddots \ddots \\ \beta_J & \beta_{J-1} & \beta_{J-2} & \ddots \\ 0 & \beta_J & \beta_{J-1} & \ddots \\ 0 & 0 & \beta_J & \ddots \\ \vdots & \vdots & \ddots & \ddots \end{pmatrix} = \begin{pmatrix} \beta_0 & 0 & 0 & \cdots \\ \beta_1 a_0 & \beta_0 & 0 & \cdots \\ \beta_2 a_0^2 & \beta_1 a_1 & \beta_0 & \ddots \\ \vdots & \vdots & \vdots & \ddots \ddots \\ \beta_J a_0^J & \beta_{J-1} a_1^{J-1} & \beta_{J-2} a_2^{J-2} & \ddots \\ 0 & \beta_J a_1^J & \beta_{J-1} a_2^{J-1} & \ddots \\ 0 & 0 & \beta_J a_2^J & \ddots \\ \vdots & \vdots & \ddots & \ddots \end{pmatrix} \begin{pmatrix} c_{0,0} & 0 & 0 & \cdots \\ c_{1,0} & c_{1,1} & 0 & \ddots \\ c_{2,0} & c_{2,1} & c_{2,2} & \ddots \\ c_{3,0} & c_{3,1} & c_{3,2} & \ddots \\ c_{4,0} & c_{4,1} & c_{4,2} & \ddots \\ c_{5,0} & c_{5,1} & c_{5,2} & \ddots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

for C .

Considering the n th column of matrix C and using the fact that $\beta_0 = 1$ for all n , leads to the recursion:

$$\begin{aligned}
 c_{n,n} &= 1 \quad \text{for all } n, \\
 c_{n+k,n} &= \beta_k - \sum_{i=1}^k \beta_i a_{n+k-i}^i c_{n+k-i,n} \quad \text{if } 1 \leq k \leq J, \quad * \\
 c_{n+k,n} &= - \sum_{i=1}^J \beta_i a_{n+k-i}^i c_{n+k-i,n} \quad \text{if } k > J. \quad **
 \end{aligned}$$

This recursion is profitably viewed as a vector recursion. For $n \geq 0$ and $j \geq n + J$, let $\vec{v}_{j,n} = (c_{j-J+1,n}, c_{j-J+2,n}, \dots, c_{j,n})^T$. The J by J matrix

$$M_n = \begin{pmatrix} 0 & 1 & 0 & \dots & 0 & 0 \\ 0 & 0 & 1 & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \dots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & 0 & 1 \\ -\beta_J a_{n-J+1}^J & -\beta_{J-1} a_{n-J+2}^{J-1} & -\beta_{J-2} a_{n-J+3}^{J-2} & \dots & -\beta_2 a_{n-1}^2 & -\beta_1 a_n \end{pmatrix}$$

encodes the map which takes $(c_1, c_2, \dots, c_J)^T$ to $(c_2, c_3, \dots, c_J, -\sum_{i=1}^J \beta_i a_{n-i+1}^i c_{J+1-i})^T$. This allows equation ** to be expressed by the recursion: $\vec{v}_{n+k,n} = M_{n+k} \vec{v}_{n+k-1,n}$ for $k > J$. Tracing the recursion backwards, one obtains

$$\vec{v}_{n+k,n} = M_{n+k} M_{n+k-1} \dots M_{n+J+1} \vec{v}_{n+J,n} \quad \text{for } k > J.$$

The recursion matrix M_n and its pointwise limit

$$M_\infty = \begin{pmatrix} 0 & 1 & 0 & \dots & 0 & 0 \\ 0 & 0 & 1 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & 1 & 0 \\ 0 & 0 & 0 & \dots & 0 & 1 \\ -\beta_J & -\beta_{J-1} & -\beta_{J-2} & \dots & -\beta_2 & -\beta_1 \end{pmatrix}$$

will play dominant roles in what follows. Note that $\vec{v}_j = (z_j^{J-1}, z_j^{J-2}, \dots, z_j, 1)^T$ is an eigenvector for M_∞ with eigenvalue w_j for $j = 1, \dots, J$. It is well-known that $\{\vec{v}_j : j = 1, 2, \dots, J\}$ forms a basis for \mathbb{C}^J , and it turns out that in the proceeding section it will be useful to describe the action of M_n in terms of a basis of these eigenvectors.

To determine when C is bounded, we will estimate the norms of such matrix products for large k . The following result due to Adams and McGuire in [3] will then provide the desired condition:

THEOREM 2.2. (Adams-McGuire) *If $p > 0$, then the matrix*

$$M = \begin{pmatrix} 0 & 0 & 0 & 0 & \dots \\ \frac{p}{2} & 0 & 0 & 0 & \dots \\ \frac{p}{2}(\frac{2}{3})^p & \frac{p}{3} & 0 & 0 & \dots \\ \frac{p}{2}(\frac{2}{4})^p & \frac{p}{3}(\frac{3}{4})^p & \frac{p}{3} & 0 & \dots \\ \frac{p}{2}(\frac{2}{5})^p & \frac{p}{3}(\frac{3}{5})^p & \frac{p}{4}(\frac{4}{5})^p & \frac{p}{5} & \dots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$

is bounded if and only if $p > \frac{1}{2}$.

The following result gives sufficient conditions on the decay of the norms of products of the matrices M_n and the norms of the “starting vectors” in order for the containment $\phi(z)H^2(\mathbb{D}) \subset H(K)$ to hold.

THEOREM 2.3. *If M_n is the recursion matrix defined above and for some $p > 1/2$, $\mu \in \mathbb{Z}^+$, $N \geq J$, and $D_1 > 0$, we have the estimate*

$$\|M_{n+\mu-1}M_{n+\mu-2}\cdots M_n\| \leq (1 - p\mu/n)$$

for all $n \geq N$, and

$$\|\vec{v}_{n+J,n}\| \leq D_1 \frac{p}{n+J}$$

for all n , then $\phi(z)H^2(\mathbb{D}) \subset H(K)$.

Proof. First notice that it suffices to prove that the matrix C defined above is the matrix of a bounded operator on ℓ^2 . Let $D_2 = \sup_n \|M_n\|$. Note it is clear that $D_2 < \infty$ as the entries in M_n are uniformly bounded in n .

Given $n, k \in \mathbb{Z}^+$ with $k \geq N + J$, let m be the largest integer such that $k - m\mu \geq N + J$. Then $m \geq 0$, and from the recursion

$$\begin{aligned} |c_{n+k,n}| &\leq \|\vec{v}_{n+k,n}\| \\ &= \|M_{n+k}M_{n+k-1}\cdots M_{n+k-m\mu+1}\vec{v}_{n+k-m\mu,n}\| \\ &\leq \|M_{n+k}M_{n+k-1}\cdots M_{n+k-m\mu+1}\| \|\vec{v}_{n+k-m\mu,n}\| \\ &\leq \prod_{j=1}^m (1 - p\mu/(n+k+1-j\mu)) \|\vec{v}_{n+k-m\mu,n}\|. \end{aligned}$$

For $0 < \varepsilon < 1$, $\log(1 - \varepsilon) < -\varepsilon$. Without loss of generality we may assume $N > p\mu$, which affords

$$\begin{aligned} \log \prod_{j=1}^m (1 - p\mu/(n+k+1-j\mu)) &< \sum_{j=1}^m (-p\mu/(n+k+1-j\mu)) \\ &< \sum_{j=0}^{m-1} (-p\mu/(n+N+J+1+(j+1)\mu)) \\ &\leq \int_0^m \left(-\frac{p\mu}{N'+\mu x} \right) dx \\ &= -p \log(N'+\mu x) \Big|_0^m \\ &= \log \left(\left[\frac{N'}{N'+m\mu} \right]^p \right) \end{aligned}$$

where $N' = n + N + J + \mu + 1$. Therefore,

$$\begin{aligned} |c_{n+k,n}| &\leq \left[\frac{N'}{N'+m\mu} \right]^p \|\vec{v}_{n+k-m\mu,n}\| \\ &= \left[\frac{N'}{N'+m\mu} \right]^p \|M_{n+k-m\mu} M_{n+k-m\mu-1,n} \cdots M_{n+J+1} \vec{v}_{n+J,n}\| \\ &\leq \left[\frac{N'}{N'+m\mu} \right]^p D_2^{N+\mu} \|\vec{v}_{n+J,n}\| \\ &\leq D_2^{N+\mu} D_1 \frac{p}{n+J} \left[\frac{N'}{N'+m\mu} \right]^p. \end{aligned}$$

Recalling that the Schur or Hadamard product of a bounded matrix with another matrix with entries bounded away from 0 and ∞ is bounded (see Lemma 2.1 in [3]), a simple application of the preceding theorem demonstrates that C is bounded. \square

3. Finite bandwidth reproducing kernels

In this section, we obtain an explicit decomposition for these spaces in analogy with [3] in the case $p > 1/2$ and $\lim_{n \rightarrow \infty} n(1 - a_n) = p$. In doing so we substantially extend their results to arbitrary bandwidths and more general weight sequences.

The following two lemmas have routine proofs and are needed for the purposes of computation.

LEMMA 3.1. *If A_1, A_2, \dots, A_k are $n \times n$ matrices with complex entries bounded in modulus by c then*

$$\|A_1 \dots A_k\| \leq n^k c^k.$$

LEMMA 3.2. *If z_1, z_2, \dots, z_J are points on the unit circle \mathbb{T} , then $(1, 1, \dots, 1) \in \mathbb{C}^J$ is a limit point of the set $\{(z_1^\mu, z_2^\mu, \dots, z_J^\mu) : \mu \in \mathbb{Z}^+\}$.*

Proof. Repeatedly apply the compactness of \mathbb{T} . \square

We now proceed to the statement and proof of the main lemma.

LEMMA 3.3. *Let M_n denote the recursion matrix defined above, $\{a_n\}$ a sequence satisfying $\lim_{n \rightarrow \infty} n(1 - a_n) = p$ where $p > 1/2$, and X the change of basis matrix whose j th column is the eigenvector \vec{v}_j of the limiting matrix M_∞ . If $\widehat{M}_n = X^{-1}M_nX$, then for all $\varepsilon > 0$, there exist positive integers μ and N such that for all $n > N$*

$$\|\widehat{M}_{n+\mu-1} \dots \widehat{M}_n\| \leq 1 - \frac{(\mu p - \varepsilon)}{n}.$$

Proof. Let μ be a large positive integer to be chosen later and fix k with $0 \leq k < \mu - 1$. We will choose N later based on an appropriate choice of μ . Linearize M_{n+k} by writing $M_{n+k} = M_\infty + (p/n)B + R_{n,k}$, where B is the J by J matrix whose first $J - 1$ rows are zero and whose last row is

$$(J\beta_J (J - 1)\beta_{J-1} (J - 2)\beta_{J-2} \dots 2\beta_2 \beta_1)$$

and $R_{n,k}$ is the J by J matrix whose first $J - 1$ rows are zero and whose J th row is

$$\left(\left(1 - a_{n-J+k+1}^J - \frac{pJ}{n} \right) \beta_J \dots \left(1 - a_{n-1+k}^2 - \frac{2p}{n} \right) \beta_2 \left(1 - a_{n+k} - \frac{p}{n} \right) \beta_1 \right).$$

Since $R_{n,k}$ can be bounded entrywise by $\frac{E(n)}{n}$, where $E(n)$ is some function satisfying $\lim_{n \rightarrow \infty} E(n) = 0$, it follows by Lemma 3.1 that $\|R_{n,k}\| \leq \frac{JE(n)}{n}$. We compute

$$\begin{aligned} \widehat{M}_{n+\mu-1} \dots \widehat{M}_n &= X^{-1} \prod_{k=0}^{\mu-1} (M_\infty + \frac{pB}{n} + R_{n,k})X \\ &= X^{-1} \left(M_\infty^\mu + \sum_{k=0}^{\mu-1} M_\infty^k \frac{pB}{n} M_\infty^{\mu-1-k} + R \right) X, \end{aligned}$$

where R is the sum of all products in the expansion involving the matrices $R_{n,k}$. (There are $3^\mu - \mu - 1$ such terms). Thus, $\|X^{-1}RX\| < \frac{C_1 E(n)}{n}$ where C_1 is a constant that depends only on J and μ .

The crucial norm estimate will come from

$$X^{-1} \left(M_\infty^\mu + \sum_{k=0}^{\mu-1} M_\infty^k \frac{B}{n} M_\infty^{\mu-1-k} \right) X,$$

so we turn to a computation of this norm. A straightforward Gaussian elimination shows that the vector $\vec{v}_0 = (0, 0, \dots, 0, 1)$ can be expressed in terms of the eigenvectors for M_∞ as $\sum_{j=1}^J -w_j / \phi'(z_j) \vec{v}_j$.

To compute the norm of $X^{-1} \left(M_\infty^\mu + \sum_{k=0}^{\mu-1} M_\infty^k \frac{B}{n} M_\infty^{\mu-1-k} \right) X$, consider the action of $\sum_{k=0}^{\mu-1} M_\infty^k \frac{B}{n} M_\infty^{\mu-1-k}$ on \vec{v}_h for $h \in \{1, 2, \dots, J\}$. Note that $\phi(z) = 1 + \sum_{k=1}^J \beta_k z^k =$

$\prod_{j=1}^J (1 - w_j z)$ and notice that

$$\phi'(z_h) = -w_h \prod_{j:j \neq h} (1 - w_j z_h) = \sum_{k=1}^J k \beta_k z_h^{k-1}.$$

Now, z_j is on the unit circle, so $(1 - w_j z_h) = w_j(z_j - z_h)$.

Thus,

$$\phi'(z_h) = \left(-\prod_{j=1}^J w_j\right) \prod_{j:j \neq h} (z_j - z_h).$$

Therefore,

$$\begin{aligned} B\vec{v}_h &= \phi'(z_h)\vec{v}_0 \\ &= \phi'(z_h) \sum_{j=1}^J -w_j/\phi'(z_j)\vec{v}_j \\ &= -w_h\vec{v}_h - \sum_{j:j \neq h} w_j \frac{\phi'(z_h)}{\phi'(z_j)}\vec{v}_j. \end{aligned}$$

Thus,

$$\begin{aligned} \sum_{k=0}^{\mu-1} M_\infty^k \frac{pB}{n} M_\infty^{\mu-1-k} \vec{v}_h &= \sum_{k=0}^{\mu-1} w_h^{\mu-1-k} M_\infty^k \frac{pB}{n} \vec{v}_h \\ &= -\frac{p}{n} w_h^{\mu-1} \sum_{k=0}^{\mu-1} w_h^{-k} M_\infty^k \left(w_h \vec{v}_h + \sum_{j:j \neq h} w_j \frac{\phi'(z_h)}{\phi'(z_j)} \vec{v}_j \right) \\ &= -\frac{p}{n} w_h^{\mu-1} \sum_{k=0}^{\mu-1} w_h^{-k} \left(w_h^{k+1} \vec{v}_h + \sum_{j:j \neq h} w_j^{k+1} \frac{\phi'(z_h)}{\phi'(z_j)} \vec{v}_j \right) \\ &= -\frac{\mu p}{n} w_h^\mu \vec{v}_h + \sum_{j:j \neq h} -\frac{p}{n} \frac{w_j}{w_h^{1-\mu}} \left(\frac{1 - (w_j/w_h)^\mu}{1 - w_j/w_h} \right) \frac{\phi'(z_h)}{\phi'(z_j)} \vec{v}_j. \end{aligned}$$

By Lemma 3.2, for each $\varepsilon > 0$, there is a $\mu \in \mathbb{N}$ such that each of the modulus of each of coefficients of v_j for $j \neq h$ above is less than $\frac{\varepsilon}{2Jn}$.

Since $M_\infty^\mu \vec{v}_h = w_h^\mu \vec{v}_h$, it follows that the norm of $X^{-1} \left(M_\infty^\mu + \sum_{k=0}^{\mu-1} M_\infty^k \frac{B}{n} M_\infty^{\mu-1-k} \right) X$ is bounded above by the norm of the matrix

$$P = \begin{pmatrix} \left(1 - \frac{\mu p}{n}\right) & \frac{\varepsilon}{2Jn} & \frac{\varepsilon}{2Jn} & \frac{\varepsilon}{2Jn} & \cdots & \frac{\varepsilon}{2Jn} \\ \frac{\varepsilon}{2Jn} & \left(1 - \frac{\mu p}{n}\right) & \frac{\varepsilon}{2Jn} & \frac{\varepsilon}{2Jn} & \cdots & \frac{\varepsilon}{2Jn} \\ \frac{\varepsilon}{2Jn} & \frac{\varepsilon}{2Jn} & \left(1 - \frac{\mu p}{n}\right) & \frac{\varepsilon}{2Jn} & \cdots & \frac{\varepsilon}{2Jn} \\ \frac{\varepsilon}{2Jn} & \frac{\varepsilon}{2Jn} & \frac{\varepsilon}{2Jn} & \left(1 - \frac{\mu p}{n}\right) & \cdots & \frac{\varepsilon}{2Jn} \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ \frac{\varepsilon}{2Jn} & \frac{\varepsilon}{2Jn} & \frac{\varepsilon}{2Jn} & \frac{\varepsilon}{2Jn} & \cdots & \left(1 - \frac{\mu p}{n}\right) \end{pmatrix}.$$

But from the triangle inequality we have the estimate

$$\|P\| \leq \left(1 - \frac{\mu p}{n}\right) + \frac{\varepsilon}{2n}.$$

Putting all of our calculations together and choosing N large enough so that for $n > N$, $E(n) < \frac{\varepsilon}{2C_1}$, we deduce that, for all $n > N$:

$$\left\|\widehat{M}_{n+\mu-1} \dots \widehat{M}_n\right\| \leq 1 - \frac{\mu p}{n} + \frac{\varepsilon}{2n} + \frac{\varepsilon}{2n} = 1 - \frac{(\mu p - \varepsilon)}{n}. \quad \square$$

Now we are ready to prove the containment result.

THEOREM 3.4. *If $H(K)$ denotes the reproducing kernel Hilbert space with orthonormal basis*

$$f_n(z) = \phi(a_n z) z^n$$

satisfying $p > 1/2$ and $\lim_{n \rightarrow \infty} n(1 - a_n) = p$, then $\phi(z)H^2(\mathbb{D}) \subset H(K)$.

Proof. This is a simple application of Theorem 2.3 and Lemma 3.3. First, choose $\varepsilon > 0$ sufficiently small so that $p - \varepsilon > 1/2$. By Lemma 3.3, there exist positive integers μ and N such that for all $n > N$

$$\left\|\widehat{M}_{n+\mu-1} \dots \widehat{M}_n\right\| \leq 1 - \frac{(\mu p - \varepsilon)}{n} = 1 - \frac{\mu p'}{n},$$

where $p' = p - \frac{\varepsilon}{\mu} > 1/2$. Note

$$\begin{aligned} \left\|M_{n+\mu-1} M_{n+\mu-2} \dots M_n\right\| &= \left\|X \widehat{M}_{n+\mu-1} \widehat{M}_{n+\mu-2} \dots \widehat{M}_n X^{-1}\right\| \\ &\leq \left\|\widehat{M}_{n+k} \widehat{M}_{n+k-1} \dots \widehat{M}_{n+k-m\mu+1}\right\| \|X\| \|X^{-1}\| \\ &\leq \|X\| \|X^{-1}\| \left(1 - \frac{\mu p'}{n}\right). \end{aligned}$$

The extra constant is harmless in regards to the proof of Theorem 2.3.

It only remains to check the growth rate on the starting vectors $\vec{v}_{n+J,n}$, using our previous notation. We claim that for each $1 \leq j \leq J$, there exists a bounded sequence of complex numbers $\{\alpha_{n,j}\}_n$, such that for all $n \in \mathbb{N}$, $c_{n+j,n} = (1 - a_n)\alpha_{n,j}$. Note that this implies there exists a positive real constant M such that $\|\vec{v}_{n+J,n}\| \leq M|1 - a_n|$, which in turn implies the starting vectors satisfy the growth rate of Theorem 2.3.

We prove the claim by induction on j . For the base case, note that $c_{n+1,n} = \beta_1 - a_n \beta_1 c_{n,n} = \beta_1(1 - a_n)$. Then notice that

$$\begin{aligned} c_{n+j,n} &= \beta_j(1 - a_n^j) - \sum_{i=1}^{j-1} \beta_i a_{n+j-i}^i c_{n+j-i,n} \\ &= \beta_j(1 + a_n + a_n^2 + \dots + a_n^{j-1})(1 - a_n) - \sum_{i=1}^{j-1} \beta_i a_{n+j-i}^i (1 - a_n) \alpha_{n,j-i}. \end{aligned}$$

By induction, the claim holds.

As the hypotheses of Theorem 2.3 are evidently satisfied, the containment follows. \square

EXAMPLE 3.5. This example shows that if $a_n \rightarrow 1$ more rapidly then $a_n = 1 - p/n$, then the containment of the previous result does not occur. Specifically, if $J = 2$, $z_1 = 1$, $z_2 = -1$, and $a_n = 1 - \frac{1}{(n+2)^2}$, then $(1 - z)(1 + z)H^2(\mathbb{D}) \subseteq H(K)$ if and only if there is a bounded matrix C satisfying $\hat{L} = LC$, where

$$\begin{pmatrix} 1 & 0 & 0 & \cdots \\ 0 & 1 & 0 & \cdots \\ -1 & 0 & 1 & \cdots \\ 0 & -1 & 0 & \ddots \\ 0 & 0 & -1 & \ddots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & \cdots \\ 0 & 1 & 0 & \cdots \\ -\frac{9}{16} & 0 & 1 & \cdots \\ 0 & -\frac{64}{81} & 0 & \ddots \\ 0 & 0 & -\frac{225}{256} & \ddots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix} \begin{pmatrix} c_{0,0} & 0 & 0 & \cdots \\ c_{1,0} & c_{1,1} & 0 & \cdots \\ c_{2,0} & c_{2,1} & c_{2,2} & \ddots \\ c_{3,0} & c_{3,1} & c_{3,2} & \ddots \\ c_{4,0} & c_{4,1} & c_{4,2} & \ddots \\ c_{5,0} & c_{5,1} & c_{5,2} & \ddots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

The entries of C are completely determined by this equation and it is straightforward to show that $\lim c_{n,0} \neq 0$ and thus that C is not bounded. The same argument works for $a_n = 1 - \frac{1}{(n+2)^p}$ with $p > 1$.

Before tackling the second half of the decomposition, a few different results will be required. First, to ensure this decomposition actually makes sense we need to establish that the natural domain of $H(K)$, which we denote by \mathcal{D} , of $H(K)$ consists of the unit disc \mathbb{D} plus the J “extra” points on the boundary z_1, z_2, \dots, z_J .

PROPOSITION 3.6. *If \mathcal{D} denotes the natural domain of the space $H(K)$, then*

$$\mathcal{D} = \mathbb{D} \cup \{z_1, z_2, \dots, z_J\}$$

Proof. It suffices to verify that for $1 \leq j \leq J$ we have $\sum_{n=0}^\infty |f_n(z_j)|^2 < \infty$. But this is clear, as $\sum_{n=0}^\infty |f_n(z_j)|^2 \lesssim \sum_{n=0}^\infty |1 - a_n|^2$ which is comparable to $\sum_{n=0}^\infty \frac{p^2}{n^2} < \infty$. \square

Next, we proceed to state two technical propositions that we will need in the forthcoming proof. The proofs are postponed to the next section. The second theorem relies on results from the theory of symmetrical polynomials.

PROPOSITION 3.7. *The matrix A defined by*

$$A = \begin{pmatrix} K(z_1, z_1) & K(z_2, z_1) & \cdots & K(z_J, z_1) \\ K(z_1, z_2) & K(z_2, z_2) & \cdots & K(z_J, z_2) \\ \vdots & \vdots & \cdots & \vdots \\ K(z_1, z_J) & K(z_2, z_J) & \cdots & K(z_J, z_J) \end{pmatrix}$$

is invertible.

PROPOSITION 3.8. For $j \in \{1, 2, \dots, J\}$ define

$$\mu_j = \prod_{k \neq j} (w_j - w_k).$$

If, for $n \in \mathbb{Z}$

$$Q_n(x) = \sum_{j=1}^J \frac{w_j^n}{\mu_j} \phi(x/w_j) w_j^n,$$

then $Q_0(x), Q_1(x), \dots$ satisfy the recursion:

$$\sum_{i=0}^n \beta_i Q_{n-i}(x) = \beta_{n+1} (x^{n+1} - 1).$$

THEOREM 3.9. For every $f \in H(K)$, there exists a $g \in H^2(\mathbb{D})$ and constants $b_1, b_2, \dots, b_J \in \mathbb{C}$, such that

$$f(z) = \phi(z)g(z) + b_1 K(z, z_1) + \dots + b_J K(z, z_J).$$

Proof. Given $f \in H(K)$, first choose b_1, b_2, \dots, b_J so that

$$f(z) - b_1 K(z, z_1) - b_2 K(z, z_2) - \dots - b_J K(z, z_J)$$

vanishes at $z = z_1, \dots, z_J$. Note this is always possible in light of Proposition 3.7. Thus, assume, without loss of generality, that $f \in H(K)$ satisfies $f(z_1) = f(z_2) = \dots = f(z_J) = 0$ for $j = 1, 2, \dots, J$. Our goal now becomes to demonstrate the existence of a $g \in H^2(\mathbb{D})$ so $f = \phi g$.

As $f \in H(K)$, there exists $\{\alpha_n\} \in \ell^2$ such that

$$f(z) = \sum_{n=0}^{\infty} \alpha_n f_n(z).$$

We shall refer to such a sequence $\{\alpha_n\}$ as permissible. We will produce a sequence $\{g_n\} \in \ell^2$ such that

$$f(z) = \phi(z) \left(\sum_{n=0}^{\infty} g_n z^n \right).$$

Expanding both expressions for f and equating gives:

$$\sum_{n=0}^{\infty} \sum_{k=0}^J \alpha_n a_n^k \beta_k z^k z^n = \sum_{n=0}^{\infty} \sum_{k=0}^J g_n \beta_k z^k z^n$$

Equating like powers of z above leads to the equation

$$\sum_{k=0}^J \alpha_{n-k} \beta_k a_{n-k}^k - g_{n-k} \beta_k = 0 \quad \text{for } n = 0, 1, 2, \dots$$

where any quantities with negative subscripts are treated as zero. Since $\beta_0 = 1$, this relationship can be expressed as the recursion:

$$* \quad g_n = \alpha_n + \left(\sum_{j=n-J}^{n-1} \alpha_j \beta_{n-j} a_j^{n-j} - g_j \beta_{n-j} \right).$$

Recursion * shows that one may express g_j as a linear combination,

$$g_n = \sum_{k=0}^n c_{n,k} \alpha_k,$$

for some constants $c_{n,k}$.

Applying * and equating like coefficients leads to

$$c_{n,n} = 1,$$

$$c_{n,k} = \beta_{n-k} a_k^{n-k} - \sum_{i=1}^{n-k} \beta_i c_{n-i,k} \quad n - J \leq k \leq n - 1,$$

and for $0 \leq k \leq n - J - 1$,

$$c_{n,k} = - \sum_{i=1}^J \beta_i c_{n-i,k}.$$

This suggests that one let $\{p_n : n \in \mathbb{Z}_+\}$ be the sequence of polynomials defined by the linear recursion:

$$\begin{aligned} p_0(x) &= 1, \\ p_1(x) &= -\beta_1(1-x), \\ &\vdots \\ p_n(x) &= \beta_n x^n - \sum_{i=1}^n \beta_i p_{n-i}(x) \\ &\vdots \\ p_J(x) &= \beta_J x^J - \sum_{i=1}^J \beta_i p_{J-i}(x) \end{aligned}$$

and thereafter, if $n \geq J + 1$,

$$** \quad p_n(x) = - \sum_{i=1}^J \beta_i p_{n-i}(x).$$

Then

$$c_{n+k,k} = p_n(a_k) \quad \text{if } n \geq 0.$$

To prove this claim, notice that it follows directly for all $k \geq 0$ if $n = 0, 1, \dots, J$ using induction. The cases $n > J$ then follow from the recursion by induction.

Thus the map $\{\alpha_n\} \mapsto \{g_n\}$ is encoded by the following matrix B_p (that is, $\{g_n\}_{n=0}^\infty = B_p\{\alpha_n\}_{n=0}^\infty$) where

$$B_p = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & \dots \\ p_1(a_0) & 1 & 0 & 0 & 0 & 0 & \dots \\ p_2(a_0) & p_1(a_1) & 1 & 0 & 0 & 0 & \ddots \\ p_3(a_0) & p_2(a_1) & p_1(a_2) & 1 & 0 & 0 & \ddots \\ p_4(a_0) & p_3(a_1) & p_2(a_2) & p_1(a_3) & 1 & 0 & \ddots \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots \end{pmatrix}.$$

If the matrix B_p were bounded as an operator, then the desired result would follow immediately. However, the columns of B_p are not in ℓ^2 . We will use the assumption that $f(z_j) = 0$ for $j = 1, 2, \dots, J$, to find an equivalent encoding of the map $\{\alpha_n\} \mapsto \{g_n\}$ which is bounded.

To find this alternate encoding of B_p , begin by considering the vector

$$\vec{v}_n = (p_n(a_0) \ p_{n-1}(a_1) \ \dots \ p_2(a_{n-2}) \ p_1(a_{n-1}) \ 1 \ 0 \ \dots)$$

which equals the n 'th row of B_p . Let z_j be a root of ϕ . The fact that $f(z_j) = 0$ is equivalent to the equation $\sum_{n=0}^\infty \alpha_n \phi(a_n z_j) z_j^n = 0$ which in turn means that the vector

$$\vec{w}_j = (\phi(a_0 z_j) \ \phi(a_1 z_j) z_j \ \phi(a_2 z_j) z_j^2 \ \phi(a_3 z_j) z_j^3 \ \dots) \quad \text{for } j \in \{1, 2, \dots, J\}$$

is orthogonal to any permissible $\vec{\alpha} = (\alpha_n)_{n=0}^\infty$.

Let $q_{j,n}(x) = \phi(x z_j) z_j^{-n}$ for $n \in \mathbb{Z}$. Then the polynomial sequence $\{q_{j,n} : n \in \mathbb{Z}\}$ satisfies condition ** satisfied by $\{p_n : n \in \mathbb{Z}_+\}$. (This follows directly from the fact that z_j is a root of ϕ .) Moreover, the vector

$$\vec{u}_j = (q_{j,n}(a_0) \ q_{j,n-1}(a_1) \ \dots \ q_{j,1}(a_{n-1}) \ q_{j,0}(a_n) \ q_{j,-1}(a_{n+1}) \ \dots)$$

equals $w_j^n \vec{w}_j$ and thus is orthogonal to all permissible sequences.

Therefore, the n th row \vec{v}_n of B_p can be replaced by \vec{v}_n less any linear combination of the vectors $\vec{u}_1, \vec{u}_2, \dots, \vec{u}_J$ without changing the action on permissible vectors. Proposition 3.8 shows that subtracting $\vec{v}'_n = (Q_{n-1}(a_0), Q_{n-2}(a_1), Q_{n-3}(a_2), \dots)$ from \vec{v}_n zeroes out the first n entries. Thus, an equivalent encoding of B_p is given by the matrix

$$C = \begin{pmatrix} 1 - Q_{-1}(a_0) & -Q_{-2}(a_1) & -Q_{-3}(a_2) & -Q_{-4}(a_3) & \dots \\ 0 & 1 - Q_{-1}(a_1) & -Q_{-2}(a_2) & -Q_{-3}(a_3) & \dots \\ 0 & 0 & 1 - Q_{-1}(a_2) & -Q_{-2}(a_3) & \ddots \\ 0 & 0 & 0 & 1 - Q_{-1}(a_3) & \ddots \\ \dots & \dots & \dots & \ddots & \ddots \end{pmatrix}.$$

Since w_1, w_2, \dots, w_J are discrete points on the unit circle, it is a straightforward exercise to show that there exists a constant c , independent of m and n , such that $|Q_n(a_m)| \leq c(1 - a_m)$.

Thus the map $\{\alpha_j\} \mapsto \{g_j\}$ is bounded if the matrix \widehat{C} is bounded where

$$\widehat{C} = \begin{pmatrix} 1 - a_0 & 1 - a_1 & 1 - a_2 & \dots \\ 0 & 1 - a_1 & 1 - a_2 & \ddots \\ 0 & 0 & 1 - a_3 & \ddots \\ \dots & \dots & \ddots & \ddots \end{pmatrix}.$$

But this matrix is known to be bounded since the entries behave asymptotically like $\frac{p}{n}$ (see Theorem 2.2 in [3]), establishing the result. \square

REMARK 3.10. Note that the preceding result is independent of p (it holds for all $p > 0$). Compare this to Theorem 3.4.

REMARK 3.11. Note that the proof of the preceding theorem demonstrates that if we had taken a_j s with a slower convergence rate, we would not have obtained a bounded matrix for \widehat{C} . In particular, suppose that $a_j = 1 - \left(\frac{1}{j+2}\right)^p$ where $p < 1/2$. Then we would obtain

$$\widehat{C} = \begin{pmatrix} \frac{1}{2^p} & \frac{1}{3^p} & \frac{1}{4^p} & \dots \\ 0 & \frac{1}{3^p} & \frac{1}{4^p} & \ddots \\ 0 & 0 & \frac{1}{4^p} & \ddots \\ \dots & \dots & \ddots & \ddots \end{pmatrix}.$$

This matrix is easily seen to be unbounded (in particular the ℓ^2 norms of its columns approach ∞), which suggests (but does not prove) that we might not obtain the result of the theorem in this case. Together with Example 3.5, this helps justify the consideration of spaces with the specific growth rate given in the hypothesis of the theorem.

Theorem 3.9 admits the following corollary, completing our characterization of these spaces when $p > \frac{1}{2}$ and $\lim_{n \rightarrow \infty} n(1 - a_n) = p$:

COROLLARY 3.12. *If $p > 1/2$ and $\lim_{n \rightarrow \infty} n(1 - a_n) = p$, then*

$$H(K) = \phi(z)H^2(\mathbb{D}) + \mathbb{C}K(z, z_1) + \mathbb{C}K(z, z_2) + \dots + \mathbb{C}K(z, z_J).$$

4. Proof of combinatorial propositions

LEMMA 4.1. *If $f_n(z) = \phi(a_n z)z^n$ is the n th basis vector for $H(K)$, then for some n , the matrix*

$$B_n = \begin{pmatrix} f_n(z_1) & f_n(z_2) & \cdots & f_n(z_J) \\ f_{n+1}(z_1) & f_{n+1}(z_2) & \cdots & f_{n+1}(z_J) \\ \vdots & \vdots & \vdots & \vdots \\ f_{n+J-1}(z_1) & f_{n+J-1}(z_2) & \cdots & f_{n+J-1}(z_J) \end{pmatrix}$$

is invertible.

Proof. Define $\phi_j(z) = \prod_{k \neq j} (1 - w_k z)$ and notice that $f_n(z_j) = \phi_j(a_n z_j)z_j^n (1 - a_n)$. Notice that B_n can be written as the product $B_n = D_1 C_n D_2$ where D_1 is the diagonal matrix with entries $1 - a_n, 1 - a_{n+1}, \dots, 1 - a_{n+J-1}$ and D_2 is the diagonal matrix with entries $z_1^{n+1}, z_2^{n+1}, \dots, z_J^{n+1}$. Thus,

$$C_n = \left(\phi_j(a_{n+i} z_j) z_j^{i-1} \right)_{i,j=1}^J.$$

Notice that the component-wise limit of C_n as $n \rightarrow \infty$ is

$$C_\infty = \left(\phi_j(z_j) z_j^{i-1} \right)_{i,j=1}^J,$$

which is the matrix product of the Vandermonde matrix $V = \left(z_j^{i-1} \right)_{i,j=1}^J$ with the diagonal matrix D_3 with entries $\phi_1(z_1), \phi_2(z_2), \dots, \phi_J(z_J)$. Since these matrices are invertible, so too is C_∞ . Since the invertible matrices form an open set in \mathbb{C}^{J^2} , C_n must be invertible for some n . \square

Proof of Proposition 3.7. Suppose that $A\vec{v} = \vec{0}$ for some $\vec{v} \in \mathbb{C}^J$. Then

$$0 = \langle A\vec{v}, \vec{v} \rangle = \left\| \sum_{k=1}^J v_k K(z, z_k) \right\|^2$$

But, this implies that $\sum_{k=1}^J v_k K(z, z_k) = 0$.

Use the preceding lemma to find J elements g_1, g_2, \dots, g_J of $H(K)$ with the property that $g_j(z_k) = 0$, if $k \neq j$ and $g_j(z_j) = 1$. Thus,

$$\vec{v}_j = \sum_{k=1}^J \langle g_j(z), v_k K(z, z_k) \rangle = \langle g_j(z), \sum_{k=1}^J v_k K(z, z_k) \rangle = \langle g_j(z), 0 \rangle = 0.$$

In other words, A has trivial kernel, so must be invertible. \square

The following two theorems from combinatorics provide the necessary tools to prove Proposition 3.8. Theorem 4.2 appears in [5] while Theorem 4.3 is a well-known result in combinatorics.

THEOREM 4.2. (See [5] Theorem 2.2.) For each integer $m \geq 0$,

$$\sum_{j=1}^J x_j^m / \mu_j = h_{m-J+1}(x_1, x_2, \dots, x_J),$$

where h_k is the k 'th homogeneous symmetric polynomial, which is defined to be zero for $k < 0$.

THEOREM 4.3. For each integer $m > 0$,

$$\sum_{i=0}^m \beta_i h_{m-i}(x_1, x_2, \dots, x_J) = 0.$$

Theorem 4.3 is a well-known result in the field of symmetric polynomials and we omit its proof. Now we are in a position to prove Proposition 3.8:

Proof of Proposition 3.8. First assume $0 \leq n < J$, and write

$$\sum_{i=0}^n \beta_i Q_{n-i}(x) = \sum_{k=0}^J a_k x^k.$$

Then

$$\begin{aligned} \sum_{i=0}^n \beta_i Q_{n-i}(x) &= \sum_{i=0}^n \beta_i \sum_{j=1}^J \frac{w_j^J}{\mu_j} \phi(x/w_j) w_j^{n-i} \\ &= \sum_{i=0}^n \beta_i \sum_{j=1}^J \sum_{k=0}^J \frac{w_j^J}{\mu_j} \beta_k \left(\frac{x}{w_j}\right)^k w_j^{n-i} \\ &= \sum_{k=0}^J \beta_k x^k \sum_{i=0}^n \beta_i \sum_{j=1}^J \frac{w_j^{J+n-i-k}}{\mu_j} \\ &= \sum_{k=0}^J \beta_k x^k \sum_{i=0}^n \beta_i h_{n-k-i+1}(w_1, \dots, w_J). \end{aligned}$$

Thus,

$$a_0 = \beta_0 \sum_{i=0}^n \beta_i h_{n-i+1}(w_1, \dots, w_J).$$

Now $\beta_0 = 1$ and from Theorem 2, $\sum_{i=0}^{n+1} \beta_i h_{n-i+1}(w_1, \dots, w_J) = 0$. Thus, $a_0 = -\beta_{n+1}$.

Now suppose $1 \leq k \leq n$. Then

$$\begin{aligned} a_k &= \beta_k \sum_{i=0}^n \beta_i h_{n-k-i+1}(w_1, \dots, w_J) \\ &= \beta_k \sum_{i=0}^{n-k+1} \beta_i h_{n-k-i+1}(w_1, \dots, w_J) \\ &= 0. \end{aligned}$$

For $k = n + 1$,

$$a_{n+1} = \beta_{n+1} \sum_{i=0}^n \beta_i h_{-i}(w_1, \dots, w_J) = \beta_{n+1}$$

since only the first term in the sum is non-zero.

If $n + 1 < k < J$, then $n - k - i + 1$ is always negative for $i \geq 0$ so

$$a_k = \beta_k \sum_{i=0}^n \beta_i h_{n+1-k-i}(w_1, \dots, w_J) = 0.$$

This shows that recursion * holds for $0 \leq n < J$.

Now, suppose $n \geq J$. Then,

$$\sum_{i=0}^n \beta_i Q_{n-i}(x) = \sum_{k=0}^J \beta_k \lambda^k \sum_{i=0}^n \beta_i h_{n-k-i+1}(x_1, \dots, x_J)$$

Since $n \geq J$, and $\beta_j = 0$ for $j > J$, Theorem 2 applies to show that the sum $\sum_{i=0}^n \beta_i Q_{n-i}(x)$ equals zero. \square

5. Some additional consequences

Consider next the natural question of whether $H(K)$ is closed under multiplication by the independent variable z . We have the following result:

THEOREM 5.1. *If $p > \frac{1}{2}$ and $\lim_{n \rightarrow \infty} n(1 - a_n) = p$, then z is a multiplier on $H(K)$.*

Proof. It is sufficient to show that the matrix representation of M_z with respect to the orthonormal basis $\{f_n : n \in \mathbb{N}\}$ is bounded as a matrix. Denote this matrix as $C = (c_{k,n})$. Thus

$$M_z(f_n) = \sum_{k=0}^{\infty} c_{k,n} f_k$$

with the coefficients $c_{k,n}$ yet to be determined. Expanding the sum and rearranging as powers of z shows that $c_{k,n} = 0$ for $k \leq n$ and leads to the recursion:

$$\begin{aligned} c_{n+1,n} &= 1, \\ c_{n+j+1,n} &= \beta_j a_n^j - \sum_{i=1}^j \beta_i a_{n+j+1-i}^i c_{n+j+1-i,n} \quad \text{if } 0 \leq j \leq J, \\ c_{n+J+k+1,n} &= - \sum_{i=1}^J \beta_i a_{n+J+k+1-i}^i c_{n+J+k+1-i,n} \quad \text{if } 1 \leq k. \end{aligned}$$

Notice that for $k \geq 1$, this is precisely the same recursion encoded by M_n and Theorem 3.4 applies to demonstrate the boundedness of C (as before, it is straightforward to

show the starting vectors have the appropriate decay and we omit the details, just note that the diagonal of 1s can be removed without affecting the boundedness of C). \square

Thus, in addition to establishing that the multiplier algebra of $H(K)$ contains the polynomials, we get the following nice result:

COROLLARY 5.2. *Let $H(K)$ denote the reproducing kernel Hilbert space with orthonormal basis*

$$f_n(z) = \phi(a_n z) z^n.$$

If $p > 1/2$ and $\lim_{n \rightarrow \infty} n(1 - a_n) = p$, then $H(K)$ contains the polynomials.

Proof. In light of Theorem 5.1, it suffices to show that $1 \in H(K)$. Write

$$1 = \sum_{n=0}^{\infty} c_n f_n(z) = \sum_{n=0}^{\infty} \left(\sum_{j=0}^J c_n \beta_j a_n^j z^{j+n} \right).$$

It is enough to show $\{c_n\} \in \ell^2$. Equating like powers of z leads to the recursion with starting value $c_0 = 1$ and thereafter:

$$c_j = - \sum_{i=1}^j c_{j-i} \beta_i a_{j-i}^i \quad \text{if } j \geq 1$$

where we recall that $\beta_i = 0$ if $i > J$. Once again, the vectors $\vec{v}_n = (c_{n-J+1}, c_{n-J+2}, \dots, c_n)^T$ satisfy the recursion $\vec{v}_{n+1} = M_{n+1} \vec{v}_n$ for $n = J, J + 1, \dots$ and the result follows as before. \square

Much future work could be done in this area. For instance, one could try to obtain a full characterization of the multiplier algebras of these finite bandwidth spaces.

Acknowledgements. We would like to acknowledge Paul McGuire for his careful reading of the manuscript and numerous suggestions. We would also like to acknowledge Cody Stockdale, whose honors thesis at Bucknell University in 2015 paved the way for the key matricial methods used in this paper. This paper grew out of the second author’s Honors Thesis at Bucknell University in 2017. The second author is currently supported by a NSF GRF (grant number DGE-1745038).

REFERENCES

- [1] GREGORY T. ADAMS, PAUL J. MCGUIRE, VERN I. PAULSEN, *Analytic reproducing kernels and multiplication operators*, Illinois J. Math. **36** (1992), no. 3, 404–419.
- [2] GREGORY T. ADAMS, PAUL J. MCGUIRE, *Analytic tridiagonal reproducing kernels*, J. London Math. Soc. (2) **64** (2001), no. 3, 722–738.
- [3] GREGORY T. ADAMS, PAUL J. MCGUIRE, *A class of tridiagonal reproducing kernels*, Oper. Matrices **2** (2008), no. 2, 233–247.
- [4] N. ARONSZAJN, *Theory of reproducing kernels*, Trans. Amer. Math. Soc. **68** (1950), 337–404.
- [5] WILLIAM Y. C. CHEN, JAMES D. LOUCK, *Interpolation for symmetric functions*, Adv. Math. **117** (1996), no. 1, 147–156.

- [6] R. G. DOUGLAS, *On majorization, factorization, and range inclusion of operators on Hilbert space*, Proc. Amer. Math. Soc. **17** (1966), 413–415.
- [7] VERN I. PAULSEN, *An introduction to the theory of reproducing kernel Hilbert spaces*, <http://www.math.uh.edu/~vern/rkhs.pdf> (2009).
- [8] ALLEN L. SHIELDS, *Weighted shift operators and analytic function theory*, Topics in operator theory, pp. 49–128. Math. Surveys, No. 13, Amer. Math. Soc., Providence, R. I., 1974.
- [9] CODY STOCKDALE, *Analysis of Five-Diagonal Reproducing Kernels*, Honors thesis, Bucknell University, 2015.

(Received December 9, 2020)

Gregory T. Adams
Department of Mathematics
Bucknell University
1 Dent Drive, Lewisburg, PA 17837
e-mail: adams@bucknell.edu

Nathan A. Wagner
Department of Mathematics and Statistics
Washington University in St Louis
1 Brookings Dr, St. Louis, MO 63130
e-mail: nathanawagner@wustl.edu