

NONLOCAL BOUNDARY VALUE PROBLEMS FOR (p, q) -DIFFERENCE EQUATIONS

NATTAPONG KAMSRISUK, CHANON PROMSAKON, SOTIRIS K. NTOUYAS
AND JESSADA TARIBOON

Abstract. In this paper we study existence and uniqueness of solutions for a boundary value problem for (p, q) -difference equations with nonlocal integral boundary conditions, by using classical fixed point theorems. Examples illustrating the main results are also presented.

Mathematics subject classification (2010): 05A30, 39A13, 34A12.

Keywords and phrases: (p, q) -difference equations, boundary value problems, nonlocal conditions, existence, fixed point theorems.

REFERENCES

- [1] C. R. ADAMS, *On the linear ordinary q -difference equation*, Ann. of Math. **30** (1929), 195–205.
- [2] B. AHMAD, S. K. NTOUYAS, *Boundary value problems for q -difference inclusions*, Abstr. Appl. Anal., vol. 2011, Article ID 292860, 15 pages.
- [3] B. AHMAD, S. K. NTOUYAS, J. TARIBOON, *Quantum Calculus: New Concepts, Impulsive IVPs and BVPs, Inequalities*, Trends in Abstract and Applied Analysis, vol. 4, World Scientific, Singapore, 2016.
- [4] D. W. BOYD, J. S. W. WONG, *On nonlinear contractions*, Proc. Amer. Math. Soc. **20** (1969), 458–464.
- [5] I. BURBAN, *Two-parameter deformation of the oscillator algebra and (p, q) -analog of two-dimensional conformal field theory*, J. Nonlinear Math. Phys. (2013), 384–391.
- [6] R. D. CARMICHAEL, *The general theory of linear q -difference equations*, Am. J. Math. **34** (1912), 147–168.
- [7] R. CHAKRABARTI, R. JAGANNATHAN, *A (p, q) -oscillator realization of two-parameter quantum algebras*, J. Phys. A. Math. Gen. **24** (1991), L711–0L718.
- [8] A. DOBROGOWSKA, A. ODZIEWICZ, *Second order q -difference equations solvable by factorization method*, J. Comput. Appl. Math. **193** (2006), 319–346.
- [9] M. EL-SHAHED, H. A. HASSAN, *Positive solutions of q -difference equation*, Proc. Amer. Math. Soc. **138** (2010), 1733–1738.
- [10] T. ERNST, *The history of q -calculus and a new method*, UUDM Report 2000:16, Department of Mathematics, Uppsala University, 2000, ISSN:1101-3591.
- [11] R. FERREIRA, *Nontrivial solutions for fractional q -difference boundary value problems*, Electron. J. Qual. Theory Differ. Equ., no. 70 (2010), pp. 1–10.
- [12] G. GASPER, M. RAHMAN, *Basic Hypergeometric Series*, Cambridge University Press, Cambridge, 1990.
- [13] A. GRANAS, J. DUGUNDJI, *Fixed Point Theory*, Springer-Verlag, New York, 2003.
- [14] M. N. HOUNKONNOU, J. D. BUKWELI KYEMBA, *$R(p, q)$ -calculus differentiation and integration*, SUT J. Math. **49**, no. 2, (2013), 145–167.
- [15] H. F. JACKSON, *On q -definite integral*, Quart. J. Pure and Appl. Math. **41** (1910), 193–203.
- [16] H. F. JACKSON, *q -Difference equations*, Am. J. Math. **32** (1910), 305–314.
- [17] R. JAGANNATHAN, K. SRINIVASA RAO, *Two-parameter quantum algebras, twin-basic number, and associated generalized hypergeometric series*, arXiv:math/0602613v1 [math.NT] 27 Feb 2006.

- [18] V. KAC, P. CHEUNG, *Quantum Calculus*, Springer, New York, 2002.
- [19] J. MA, J. YANG, *Existence of solutions for multi-point boundary value problem of fractional q -difference equation*, Electron. J. Qual. Theory Differ. Equ., no. 92 (2011), pp. 1–10.
- [20] T.E. MASON, *On properties of the solution of linear q -difference equations with entire function coefficients*, Am. J. Math. **37** (1915), 439–444.
- [21] M. MURSALEEN, K. J. ANSARI, A. KHAN, *On (p, q) -analogue of Bernstein Operators*, App. Math. Comp. **266** (2015), 874–882.
- [22] V. SAHAI, S. YADAV, *Representations of two parameter quantum algebras and (p, q) -special functions*, J. Math. Anal. Appl. **335** (2007), 268–279.
- [23] P. NJIONOU SADJANG, *On the fundamental theorem of (p, q) -calculus and some (p, q) -Taylor formulas*, arXiv:1309.3934v1.
- [24] J. TARIBOON, S. K. NTOUYAS, *Quantum calculus on finite intervals and applications to impulsive difference equations*, Adv. Difference Equ. (2013) 2013:282.