

GENERALIZATION OF GRACE'S THEOREM, SCHUR-SZEGÖ COMPOSITION AND COHN-EGERVÁRY THEOREM FOR BICOMPLEX POLYNOMIALS

ASHISH KUMAR* AND B. A. ZARGAR

Abstract. The aim of this paper is to extend the domain of the Grace's theorem, Schur-Szegö composition theorem and Cohn-Egerváry theorem from the set of complex numbers to the set of bicomplex numbers.

Mathematics subject classification (2020): 30C10, 30C15, 30D10.

Keywords and phrases: Bicomplex polynomials, Grace's theorem, Schur-Szegö composition.

REFERENCES

- [1] M. E. LUNA-ELIZARRARAS, M. SHAPIRO, D. C. STRUPPA, *On Clifford analysis for holomorphic mappings*, Adv. Geom., **14** (2014), 413–426.
- [2] M. MARDEN, *Geometry of Polynomials*, 2nd edition, American Mathematical Society, Providence, 1966.
- [3] A. A. POGORUI, R. M. RODRIGUEZ-DAGNINO, *On the set of zeros of bicomplex polynomials*, Complex Var. Elliptic Equ., **51**, no. 7 (2006), 725–730.
- [4] G. B. PRICE, *An Introduction to Multicomplex Spaces and Functions. Monographs and Textbooks in Pure and Applied Mathematics*, Marcel Dekker, New York, 1991.
- [5] C. SEGRE, *Le rappresentazioni reali delle forme complesse e gli enti iperalgebrici*, Math. Ann. **40** (1892), 413–467 (in Italian).