

SHARP INEQUALITIES FOR HERMITIAN TOEPLITZ DETERMINANTS FOR STRONGLY STARLIKE AND STRONGLY CONVEX FUNCTIONS

BOGUMIŁA KOWALCZYK, ADAM LECKO* AND BARBARA ŚMIAROWSKA

Abstract. Sharp upper and lower bounds are found of the second and third order Hermitian Toeplitz determinants for the classes of strongly starlike and strongly convex functions of order α ($\alpha \in [0, 1)$).

Mathematics subject classification (2010): 30C45, 30C50.

Keywords and phrases: Hermitian Toeplitz determinant, strongly starlike function, strongly convex function, Carathéodory class.

REFERENCES

- [1] J. W. ALEXANDER, *Functions which map the interior of the unit circle upon simple regions*, Ann. of Math., **17**, (1915), 12–22.
- [2] MD FIROZ ALI, D. K. THOMAS, A. VASUDEVARAO, *Toeplitz determinants whose elements are the coefficients of analytic and univalent functions*, Bull. Austr. Math. Soc., **97**, 2 (2018), 253–264.
- [3] R. M. ALI, V. SINGH, *On the fourth and fifth coefficients of strongly starlike functions*, Results Math., **29**, (1996), 197–202.
- [4] D. A. BRANNAN, W. E. KIRWAN, *On some classes of bounded univalent functions*, J. London Math. Soc., **2** (1), (1969), 431–443.
- [5] C. CARATHÉODORY, *Über den Variabilitätsbereich der Koeffizienten von Potenzreihen, die gegebene werte nicht annehmen*, Math. Ann., **64**, (1907), 95–115.
- [6] N. E. CHO, B. KOWALCZYK, O. S. KWON, A. LECKO, Y. J. SIM, *Some coefficient inequalities related to the Hankel determinant for strongly starlike functions of order alpha*, J. Math. Ineq., **11**, 2 (2017), 429–439.
- [7] N. E. CHO, B. KOWALCZYK, A. LECKO, *Sharp bounds of some coefficient functionals over the class of functions convex in the direction of the imaginary axis*, Bull. Aust. Math. Soc., **100**, (2019), 86–96.
- [8] K. CUDNA, O. S. KWON, A. LECKO, Y. J. SIM, B. ŚMIAROWSKA, *The second and third-order Hermitian Toeplitz determinants for starlike and convex functions of order α* , Boletín Soc. Mat. Mex., **26**, (2020), 361–375.
- [9] A. W. GOODMAN, *Univalent Functions*, Mariner, Tampa, Florida, 1983.
- [10] B. KOWALCZYK, O. S. KWON, A. LECKO, Y. J. SIM, B. ŚMIAROWSKA, *The third-order Hermitian Toeplitz determinant for classes of functions convex in one direction*, Bull. Malays. Math. Sci. Soc., **43**, (2020), 3143–3158.
- [11] B. KOWALCZYK, A. LECKO, Y. J. SIM, *The sharp bound of the Hankel determinant of the third kind for convex functions*, Bull. Aust. Math. Soc., **97**, (2018), 435–445.
- [12] O. S. KWON, A. LECKO, Y. J. SIM, B. ŚMIAROWSKA, *The sharp bound of the fifth coefficient of strongly starlike functions with real coefficients*, Bull. Malays. Math. Sci. Soc., **42**, 4 (2019), 1719–1735.
- [13] A. LECKO, *Some Methods in the Theory of Univalent Functions*, Oficyna Wydawnicza Politechniki Rzeszowskiej, Rzeszów, 2005.
- [14] A. LECKO, *Strongly starlike and spirallike functions*, Ann. Polon. Math., **85**, 2 (2005), 165–192.
- [15] W. MA, D. MINDA, *An internal geometric characterization of strongly starlike functions*, Ann. Univ. Mariae Curie Skłodowska Sect. A, **20**, (1991), 89–97.
- [16] W. MA, S. OWA, *Strongly starlike functions*, Panam. Math. J., **3**, 2 (1993), 49–60.

- [17] R. NEVANLINNA, *Über die konforme Abbildung von Sterngebieten*, Översikt av Finska Vetens.-Soc. Förh., Avd. A, **LXIII**, 6 (1920–1921), 1–21.
- [18] C. POMMERENKE, *Univalent functions*, Vandenhoeck & Ruprecht, Göttingen, 1975.
- [19] E. STUDY, *Vorlesungen über ausgewählte Gegenstände der Geometrie, Zweites Heft; Konforme Abbildung Einfach-Zusammenhängender Bereiche*, Druck und Verlag von B. G. Teubner, Leipzig und Berlin, 1913.
- [20] J. STANKIEWICZ, *Quelques problèmes extrémaux dans les classes des fonctions α -angulairement étoilées*, Ann. Univ. Mariae Curie-Skłodowska Sect. A, **20**, (1966), 59–75.
- [21] J. STANKIEWICZ, *On a family of starlike functions*, Ann. Univ. Mariae Curie-Skłodowska Sect. A, **22–24**, (1968–1970), 175–181.
- [22] T. SUGAWA, *A self-duality of strong starlikeness*, Kodai Math. J., **28**, (2005), 382–389.