

AN APPLICATION OF THE HAUSDORFF–YOUNG INEQUALITY

ŽIVORAD TOMOVSKI

Abstract. In this paper a generalization theorem of Zahid of a [8] has been obtained, by considering the condition $S_p(\delta)$, $p > 1$ instead of $S(\delta)$.

Mathematics subject classification (1991): 26D15, 42A20.

Key words and phrases: δ -quasi-monotone sequence, cosine trigonometric series, Fourier series, Dirichlet kernel, Abel's transformation, Holder inequality, Hausdorff–Young inequality.

REFERENCES

- [1] R. P. BOAS, *Quasi-positive sequence and trigonometric series*, Proc. Lond. Math. Soc. **14A** (1965), 38–48.
- [2] C. S. REES AND Č. V. STANOJEVIĆ, *Necessary and sufficient condition for integrability of certain cosine sums*, J. Math. Anal. Appl. **43** (1973), 579–586.
- [3] P. L. DUREN, *Theory of H^p spaces*, Academic Press, 1970.
- [4] G. A. FOMINE, *On linear method for summing Fourier series*, Mat. Sb. **66(107)** (1964), 144–152.
- [5] J. W. GARRET AND ČASLAV V. STANOJEVIĆ, *On L^1 convergence of certain cosine sums*, Proc. Amer. Math. Soc. **54** (1976), 101–105.
- [6] ———, *Necessary and sufficient conditions for L^1 convergence of trigonometric series*, Proc. Amer. Math. Soc. **60** (1976), 68–71.
- [7] S. A. TELIAKOVESKII, *Concerning a sufficient condition of Sidon for the integrability of trigonometric series*, Math. Notes (1973), 742–748.
- [8] S. A. Z. ZAHID, *Integrability of trigonometric series*, Tamkang J. Math. (1990), 295–301.