

ON THE OPERATOR EQUATION $AXB + CYD = Z$

DON HADWIN, ERIC NORDGREN AND PETER ROSENTHAL

Abstract. Suppose two n -tuples of operators A_k and B_k on a Hilbert space are given, and Φ is the mapping from the set of n -tuples of operators on the Hilbert space into the set of all operators on the space defined by $\Phi(X_1, X_2, \dots, X_n) = \sum_{k=1}^n A_k X_k B_k$. Conditions are given for Φ to be onto.

Mathematics subject classification (2000): 47B49.

Key words and phrases: Surjectivity of mappings of operator tuples.

REFERENCES

- [1] RAJENDRA BHATIA AND PETER ROSENTHAL, *How and why to solve the operator equation $AX - XB = Y$* , Bull London Math. Soc. **29** (1997), 1–21. MR1416400 (97k:47016)
- [2] M. BRESAR AND PETER SEMRL, *Elementary operators*, Proc. Royal Soc. Edinburgh Sect. A, **129** (1999), 1115–1135. MR1728537 (2001f:4701)
- [3] CHANDLER DAVIS AND PETER ROSENTHAL, *Solving linear operator equations*, Canadian J. Math. **26** (1974), 1384–1389. MR0355649 (50 #8123)
- [4] RONALD G. DOUGLAS, *On majorization, factorization, and range inclusion of operators on Hilbert space*, Proc. Amer. Math. Soc., **17** (1966), 413–415. MR0203464 (34 #3315)
- [5] PETER FILLMORE, JOSEPH STAMPLI AND JAMES WILLIAMS, *On the essential numerical range, the essential spectrum, and a problem of Halmos*, Acta Math. (Szeged) **33** (1972), 179–192. MR0322534 (48 #896)
- [6] ERIC NORDGREN AND PETER ROSENTHAL, *Boundary values of Berezin symbols*, Operator Theory: Advances and Applications, **73** (1994), 362–368. MR1320554 (96b:46036)
- [7] MARVIN ROSENBLUM, *On the operator equation $BX - XA = Q$* , Duke Math. J. **23** (1956), 263–270. MR0079235 (18, 54d)