

THE PEAK MODEL FOR FINITE RANK SUPERSINGULAR PERTURBATIONS

RYTIS JURŠENAS

Abstract. In its original form the peak model for rank one supersingular perturbations of class \mathfrak{S}_{-4} or higher of a nonnegative self-adjoint operator requires that the Gram matrix of the model should be diagonal. Here we remove the restriction on the Gram matrix. In particular we explain the origin of the Krein Q -function associated with the Gram matrix.

Mathematics subject classification (2020): 47B25, 47A06, 34B05, 35P05.

Keywords and phrases: Supersingular perturbation, exit space extension, generalized resolvent, boundary triple, linear relation, γ -field, Weyl function.

REFERENCES

- [1] S. ALBEVERIO, G. COGNOLA, M. SPREAFICO AND S. ZERBINI, *Singular perturbations with boundary conditions and the Casimir effect in the half space*, J. Math. Phys. **51** (2010), 063502.
- [2] S. S. ALBEVERIO, FASSARI AND F. RINALDI, *A remarkable spectral feature of the Schrödinger Hamiltonian of the harmonic oscillator perturbed by an attractive δ^1 -interaction centred at the origin: double degeneracy and level crossing*, J. Phys. A: Math. Theor. **46** (2013), no. 38, 385305.
- [3] S. ALBEVERIO AND P. KURASOV, *Rank One Perturbations of Not Semibounded Operators*, Integr. Equ. Oper. Theory **27**, (1997), 379–400.
- [4] S. ALBEVERIO AND P. KURASOV, *Singular Perturbations of Differential Operators*, London Mathematical Society Lecture Note Series 271, Cambridge University Press, UK, 2000.
- [5] S. ALBEVERIO, S. KUZHEL AND L. NIZHNIK, *Singularly perturbed self-adjoint operators in scales of Hilbert spaces*, Ukrainian J. Math. **59** (2007), no. 6, 787–810.
- [6] Y. ARLINSKII AND S. HASSI, *Q -functions and boundary triplets of nonnegative operators*, Recent Advances in Inverse Scattering, Schur Analysis and Stochastic Processes, 2015, pp. 89–130.
- [7] T. AZIZOV AND I. IOKHVIDOV, *Linear Operators in Spaces with an Indefinite Metric*, John Wiley & Sons. Inc., 1989.
- [8] J. BEHRNDT, V. A. DERKACH, S. HASSI AND H. DE SNOO, *A realization theorem for generalized Nevanlinna families*, Operators and Matrices **5** (2011), no. 4, 679–706.
- [9] J. BEHRNDT, S. HASSI AND H. DE SNOO, *Boundary Value Problems, Weyl Functions, and Differential Operators*, Birkhauser, 2020.
- [10] J. BEHRNDT, S. HASSI, H. DE SNOO, R. WIETSMA AND H. WINKLER, *Linear fractional transformations of Nevanlinna functions associated with a nonnegative operator*, Compl. Anal. Oper. Theory **7** (2013), no. 2, 331–362.
- [11] JUSSI BEHRNDT AND MATTHIAS LANGER, *Boundary value problems for elliptic partial differential operators on bounded domains*, J. Func. Anal. **243** (2007), 536–565.
- [12] JUSSI BEHRNDT, MATTHIAS LANGER, VLADIMIR LOTOREICHIK AND JONATHAN ROHLER, *Spectral enclosures for non-self-adjoint extensions of symmetric operators*, J. Func. Anal. **275** (2018), no. 7, 1808–1888.
- [13] JOCHEN BRÜNING, VLADIMIR GEYLER AND KONSTANTIN PANKRASHKIN, *Spectra of self-adjoint extensions and applications to solvable Schrödinger operators*, Rev. Math. Phys. **20** (2008), no. 1, 1–70.
- [14] V. DERKACH, *On generalized resolvents of Hermitian relations in Krein spaces*, J. Math. Sci. **97** (1999), no. 5, 4420–4460.

- [15] V. DERKACH, S. HASSI AND H. DE SNOO, *Singular perturbations of self-adjoint operators*, Math. Phys. Anal. Geom. **6** (2003), no. 4, 349–384.
- [16] V. DERKACH, S. HASSI, M. MALAMUD AND H. DE SNOO, *Boundary relations and their Weyl families*, Trans. Amer. Math. Soc. **358** (2006), no. 12, 5351–5400.
- [17] V. DERKACH, S. HASSI, M. MALAMUD AND H. DE SNOO, *Boundary relations and generalized resolvents of symmetric operators*, Russ. J. Math. Phys. **16** (2009), no. 1, 17–60.
- [18] V. A. DERKACH AND M. M. MALAMUD, *Generalized Resolvents and the Boundary Value Problems for Hermitian Operators with Gaps*, J. Func. Anal. **95** (1991), no. 1, 1–95.
- [19] VLADIMIR DERKACH, SEPO HASSI, MARK MALAMUD AND HENK DE SNOO, *Boundary triplets and Weyl functions. Recent developments*, Operator Methods for Boundary Value Problems, London Math. Soc. Lecture Note Series, 2012, pp. 161–220.
- [20] VLADIMIR DERKACH, SEPO HASSI AND MARK M. MALAMUD, *Generalized boundary triples, I. Some classes of isometric and unitary boundary pairs and realization problems for subclasses of Nevanlinna functions*, Math. Nachr. **293** (2020), no. 7, 1278–1327.
- [21] A. DIJKSMA, P. KURASOV AND YU. SHONDIN, *High Order Singular Rank One Perturbations of a Positive Operator*, Integr. Equ. Oper. Theory **53** (2005), 209–245.
- [22] A. DIJKSMA AND H. LANGER, *Compressions of self-adjoint extensions of a symmetric operator and M.G. Krein's resolvent formula*, Integr. Equ. Oper. Theory **90** (2018), no. 41, 1–30.
- [23] A. DIJKSMA, H. LANGER, A. LUGER AND YU. SHONDIN, *Minimal realizations of scalar generalized Nevanlinna functions related to their basic factorization*, Spectral Methods for Operators of Mathematical Physics, Operator Theory: Advances and Applications, 2004.
- [24] M. GALLONE AND A. MICHELANGELI, *Self-adjoint extensions with Friedrichs lower bound*, Compl. Anal. Oper. Theory **14** (2020), no. 73, 23 p.
- [25] I. GOHBERG, P. LANCASTER AND L. RODMAN, *Indefinite Linear Algebra and Applications*, Birkhauser, 2005.
- [26] P. R. HALMOS, *Eigenvectors and adjoints*, Lin. Alg. Appl. **4** (1971), 11–15.
- [27] S. HASSI AND H. WIETSMA, *Minimal realizations of generalized Nevanlinna functions*, Opuscula Math. **36** (2016), no. 6, 749–768.
- [28] SEPO HASSI AND SERGH KUZHEL, *On symmetries in the theory of finite rank singular perturbations*, J. Func. Anal. **256** (2009), 777–809.
- [29] R. JURŠĖNAS, *Computation of the unitary group for the Rashba spin–orbit coupled operator, with application to point-interactions*, J. Phys. A: Math. Theor. **51** (2018), no. 1, 015203.
- [30] R. JURŠĖNAS, *The A-model with mutually equal model parameters can lead to a Hilbert space model*, Operators and Matrices **15** (2021), no. 4, 1319–1336.
- [31] RYTIS JURŠĖNAS, *On some extensions of the A-model*, Opuscula Mathematica **40** (2020), no. 5, 569–597.
- [32] P. KURASOV, \mathcal{H}_n -perturbations of self-adjoint operators and Krein's resolvent formula, Integr. Equ. Oper. Theory **45** (2003), no. 4, 437–460.
- [33] P. KURASOV AND YU. V. PAVLOV, *On field theory methods in singular perturbation theory*, Lett. Math. Phys. **64** (2003), no. 2, 171–184.
- [34] PAVEL KURASOV, *Triplet extensions I: Semibounded operators in the scale of Hilbert spaces*, Journal d'Analyse Mathématique **107** (2009), no. 1, 252–286.
- [35] H. LANGER AND B. TEXTORIUS, *On generalized resolvents and Q-functions of symmetric linear relations (subspaces) in Hilbert space*, Pacific. J. Math. **72** (1977), no. 1, 135–165.
- [36] B. SIMON, *Trace Ideals and Their Applications*, 2nd ed., Mathematical Surveys and Monographs, vol. 120, American Mathematical Society, 2005.