

ON THE RESOLUTION OF A PARABOLIC EQUATION IN A NONREGULAR DOMAIN OF \mathbb{R}^3

A. KHELOUFI, R. LABBAS AND B.-K. SADALLAH

(Communicated by M. Kirane)

Abstract. In this work we give new results of existence, uniqueness and maximal regularity of a solution to a parabolic equation set in a nonregular domain Q with Cauchy-Dirichlet boundary conditions, where $Q = \{(t, x_1) \in \mathbb{R}^2 : 0 < t < T; \varphi_1(t) < x_1 < \varphi_2(t)\} \times]0, b[\subseteq \mathbb{R}^3$ with some assumptions on the functions $(\varphi_i)_{i=1,2}$. The right-hand side term of the equation is taken in $L^2(Q)$. The method used is based on the approximation of the domain Q by a sequence of subdomains $(Q_n)_n$ which can be transformed into regular domains. This work is an extension of the one space variable case studied in [12].

1. Introduction

Let Ω be an open set of \mathbb{R}^2 defined by

$$\Omega = \{(t, x_1) \in \mathbb{R}^2 : 0 < t < T; \varphi_1(t) < x_1 < \varphi_2(t)\}$$

where T is a finite positive number, while φ_1 and φ_2 are continuous real-valued functions defined on $[0, T]$, Lipschitz continuous on $]0, T[$, and such that

$$\varphi_1(t) < \varphi_2(t)$$

for $t \in]0, T[$. φ_1 is allowed to coincide with φ_2 for $t = 0$ and for $t = T$. We also assume that $(\varphi_2 - \varphi_1)$ is a bounded function on $]0, T[$. For a fixed positive number b , let Q be the three-dimensional domain defined by

$$Q = \{(t, x_1) \in \mathbb{R}^2 : 0 < t < T; \varphi_1(t) < x_1 < \varphi_2(t)\} \times]0, b[,$$

with boundary $\partial Q = (\Gamma \times]0, b[) \cup (\Omega \times \{0\}) \cup (\Omega \times \{b\})$, Γ is the boundary of Ω (see Fig.1).

In this work, we study the existence and the regularity of the solution of the parabolic equation with Cauchy-Dirichlet boundary conditions

$$(P) \begin{cases} D_t u - D_{x_1}^2 u - D_{x_2}^2 u = f & \text{in } Q \\ u = 0 & \text{on } \partial Q - \Gamma_T, \end{cases}$$

Mathematics subject classification (2010): 35K05, 35K20.

Keywords and phrases: parabolic equation, nonregular domains, anisotropic Sobolev space.

This work has been supported in part by the EGIDE grant under the C.M.E.P Program. Project N $^{\circ}$. 08 MDU 735.

where Γ_T is the part of the boundary of Q where $t = T$. The right-hand side term f of the equation lies in $L^2(Q)$.

In Baderko [3] we can find domains of the same kind but which can not include our domain. In Sadallah [12] the same problem has been studied for a $2m$ -parabolic operator in the case of one space variable. Further references on the analysis of parabolic problems in non-cylindrical domains are: Savaré [13], Aref'ev and Bagirov [2], Hoffmann and Lewis [6], Labbas, Medeghri and Sadallah [7], [8], and Alkhutov [1]. There are many other works concerning boundary-value problems in nonsmooth domains (see, for example, Grisvard [5] and the references therein).

We are especially interested in question of what conditions the functions $(\varphi_i)_{i=1,2}$ must verify in order that Problem (P) has a solution with optimal regularity, that is a solution u belonging to the anisotropic Sobolev space

$$H_V^{1,2}(Q) = \{u \in H^{1,2}(Q) : u|_{\partial Q - \Gamma_T} = 0\}$$

with

$$H^{1,2}(Q) = \{u \in L^2(Q) : D_t u, D_{x_1}^j u, D_{x_2}^j u, D_{x_1} D_{x_2} u \in L^2(Q), j = 1, 2\}?$$

An idea to solve Problem (P) consists in transforming the parabolic equation in the nonregular domain Q into a variable-coefficient equation in a regular domain. However, in order to perform this, one must assume that $\varphi_1(0) < \varphi_2(0)$ and $\varphi_1(T) < \varphi_2(T)$.

So, in Section 2, we prove that Problem (P) admits a (unique) solution when Q could be transformed into a regular domain by means of a regular change of variable, i.e., we suppose that $\varphi_1(0) < \varphi_2(0)$ and $\varphi_1(T) < \varphi_2(T)$.

In Section 3 we approximate Q by a sequence (Q_{α_n}) of such domains and we establish an *a priori* estimate of the type

$$\|u_n\|_{H^{1,2}(Q_{\alpha_n})} \leq K \|f\|_{L^2(Q_{\alpha_n})},$$

where u_n is the solution of Problem (P) in Q_{α_n} and K is a constant independent of n . Finally, in Section 4 we take limits in (Q_{α_n}) in order to reach the domain Q .

The main assumptions on the functions $(\varphi_i)_{i=1,2}$ are

$$\varphi_i'(t)(\varphi_2(t) - \varphi_1(t)) \rightarrow 0 \text{ as } t \rightarrow 0, i = 1, 2 \tag{1.1}$$

and

$$\varphi_i'(t)(\varphi_2(t) - \varphi_1(t)) \rightarrow 0 \text{ as } t \rightarrow T, i = 1, 2 \tag{1.2}$$

2. Resolution of Problem (P) in a reference domain Q_α

In this section, we replace Q by

$$Q_\alpha = \{(t, x_1) \in \mathbb{R}^2 : \alpha < t < T - \alpha, \varphi_1(t) < x_1 < \varphi_2(t)\} \times]0, b[,$$

with $\alpha > 0$. Thus, we have

$$\begin{cases} \varphi_1(\alpha) < \varphi_2(\alpha) \\ \varphi_1(T - \alpha) < \varphi_2(T - \alpha), \end{cases}$$

(see Fig.1).




Figure 1.

We can find a change of variable ψ mapping Q_α into the parallelepiped

$$P_\alpha =]\alpha, T - \alpha[\times]0, 1[\times]0, b[,$$

which leaves the variable t unchanged. ψ is defined as $\psi : Q_\alpha \rightarrow P_\alpha$ such that:

$$(t, x_1, x_2) \mapsto \psi(t, x_1, x_2) = \left(\tau, y_1, y_2 \right) = \left(t, \frac{x_1 - \varphi_1(t)}{\varphi_2(t) - \varphi_1(t)}, x_2 \right).$$

The mapping ψ transforms the parabolic equation in the domain Q_α into a variable-coefficient parabolic equation in the parallelepiped P_α . Indeed, the equation

$$D_t u - D_{x_1}^2 u - D_{x_2}^2 u = f$$

in Q_α is equivalent to the following

$$D_\tau v + a(\tau, y_1) D_{y_1} v - c(\tau) D_{y_1}^2 v - D_{y_2}^2 v = g$$

in P_α , where a and c are defined by

$$a(\tau, y_1) = \frac{(\varphi_1'(\tau) - \varphi_2'(\tau)) y_1 - \varphi_1'(\tau)}{\varphi_2(\tau) - \varphi_1(\tau)} \quad \text{and} \quad c(\tau) = \frac{1}{(\varphi_2(\tau) - \varphi_1(\tau))^2},$$

and

$$\begin{aligned} f(t, x_1, x_2) &= g(\tau, y_1, y_2), \\ u(t, x_1, x_2) &= v(\tau, y_1, y_2). \end{aligned}$$

Since the functions a, c and $\varphi_2 - \varphi_1$ are bounded, it is easy to check the following.

LEMMA 2.1. $u \in H^{1,2}(Q_\alpha)$ if and only if $v \in H^{1,2}(P_\alpha)$.

In other words, ψ preserves the Sobolev space $H^{1,2}$.

The boundary conditions on v which correspond to the boundary conditions on u are the following

$$v|_{\partial P_\alpha - \Gamma_{T-\alpha}} = 0,$$

where $\Gamma_{T-\alpha}$ is the part of the boundary of P_α where $t = T - \alpha$.

In the sequel, the variables (τ, v_1, v_2) will be denoted again by (t, x_1, x_2) .

THEOREM 2.1. *The operator*

$$L' = D_t + aD_{x_1} - cD_{x_1}^2 - D_{x_2}^2$$

is an isomorphism from $H_Y^{1,2}(P_\alpha)$ into $L^2(P_\alpha)$, with

$$H_Y^{1,2}(P_\alpha) = \{u \in H^{1,2}(P_\alpha) : u|_{\partial P_\alpha - \Gamma_{T-\alpha}} = 0\}.$$

Consider the simplified problem

$$(P') \begin{cases} D_t v - c(t)D_{x_1}^2 v - D_{x_2}^2 v = g & \text{in } P_\alpha, \\ v|_{\partial P_\alpha - \Gamma_{T-\alpha}} = 0. \end{cases}$$

Note that $g \in L^2(P_\alpha)$ if and only if $f \in L^2(Q_\alpha)$.

LEMMA 2.2. *For every $g \in L^2(P_\alpha)$, there exists a unique $v \in H_Y^{1,2}(P_\alpha)$ solution of (P') .*

Proof. Since the coefficient $c(t)$ is continuous in $\overline{P_\alpha}$, the optimal regularity is given by Ladyzhenskaya-Solonnikov-Ural'tseva [9].

LEMMA 2.3. *Let B be an operator, $B : H_Y^{1,2}(P_\alpha) \rightarrow L^2(P_\alpha)$, defined by*

$$Bv = a(t, x_1)D_{x_1} v.$$

The operator B is compact.

Proof. P_α has the "horn property" of Besov [4], so

$$\begin{aligned} D_{x_1} : H_Y^{1,2}(P_\alpha) &\rightarrow H^{\frac{1}{2},1}(P_\alpha) \\ v &\mapsto D_{x_1} v \end{aligned}$$

is continuous. Since P_α is bounded, the canonical injection is compact from $H^{\frac{1}{2},1}(P_\alpha)$ into $L^2(P_\alpha)$, where (see [10]):

$$H^{\frac{1}{2},1}(P_\alpha) = L^2(\alpha, T - \alpha; H^1([0, 1[\times]0, b[)]) \cap H^{\frac{1}{2}}(\alpha, T - \alpha; L^2([0, 1[\times]0, b[))).$$

For the complete definitions of the $H^{r,s}$ Hilbertian Sobolev spaces see for instance [4]. Consider the composition

$$D_{x_1} : H_Y^{1,2}(P_\alpha) \rightarrow H^{\frac{1}{2},1}(P_\alpha) \rightarrow L^2(P_\alpha)$$

$$v \mapsto D_{x_1} v \mapsto D_{x_1} v,$$

then D_{x_1} is a compact operator from $H_Y^{1,2}(P_\alpha)$ into $L^2(P_\alpha)$. Since $a(\cdot, \cdot)$ is a bounded function, the operator aD_{x_1} is also compact from $H_Y^{1,2}(P_\alpha)$ into $L^2(P_\alpha)$.

Lemma 2.2 shows that the operator $L_1 = D_t - cD_{x_1}^2 - D_{x_2}^2$ is an isomorphism from $H_Y^{1,2}(P_\alpha)$ into $L^2(P_\alpha)$; on the other hand the operator aD_{x_1} is compact, consequently, $L_1 + aD_{x_1}$ is a Fredholm operator from $H_Y^{1,2}(P_\alpha)$ into $L^2(P_\alpha)$. Thus the invertibility of $L_1 + aD_{x_1}$ follows from its injectivity.

LEMMA 2.4. *The space*

$$D([\alpha, T - \alpha]; H^4([0, 1[\times]0, b[) \cap H_0^1([0, 1[\times]0, b[)), \text{ (see [10, p.13]),}$$

is dense in the subspace of $H^{1,2}([\alpha, T - \alpha[\times]0, 1[\times]0, b[)$ defined by

$$u = 0 \text{ on }]\alpha, T - \alpha[\times \{0\} \times]0, b[\text{ and }]\alpha, T - \alpha[\times \{1\} \times]0, b[.$$

It is a particular case of Theorem 2.1 [10].

We shall need the following result in order to justify the calculus of Section 3.

LEMMA 2.5. *The space*

$$\{u \in H^4(P_\alpha); u|_{\partial P_\alpha - \Gamma_{T-\alpha}} = 0\}$$

is dense in the space

$$\{u \in H^{1,2}(P_\alpha); u|_{\partial P_\alpha - \Gamma_{T-\alpha}} = 0\}.$$

Proof. Let Γ_α be the part of the boundary of P_α where $t = \alpha$. The previous lemma shows that the space

$$\{u \in H^4(P_\alpha); u|_{\partial P_\alpha - \Gamma_{T-\alpha} - \Gamma_\alpha} = 0\}$$

is dense in the space

$$\{u \in H^{1,2}(P_\alpha); u|_{\partial P_\alpha - \Gamma_{T-\alpha} - \Gamma_\alpha} = 0\}.$$

So, if

$$u \in \{u \in H^{1,2}(P_\alpha); u|_{\partial P_\alpha - \Gamma_{T-\alpha} - \Gamma_\alpha} = 0\},$$

then there exists a sequence

$$(u_n) \in \{u \in H^4(P_\alpha); u|_{\partial P_\alpha - \Gamma_{T-\alpha} - \Gamma_\alpha} = 0\}$$

such that

$$u_n \rightharpoonup u \text{ weakly in } H^{1,2}(P_\alpha), \text{ as } n \rightarrow \infty.$$

Let (e_n) be a sequence of $C^\infty([\alpha, T - \alpha])$ such that

$$e_n(t) = \begin{cases} 1, & \text{if } t \geq \alpha + \frac{1}{n}, \\ 0, & \text{if } t \leq \alpha + \frac{1}{2n}. \end{cases}$$

The sequence $(e_n u_n)$ belongs to

$$\{u \in H^4(P_\alpha); u|_{\partial P_\alpha - \Gamma_{T-\alpha}} = 0\}.$$

In addition

$$e_n u_n \rightharpoonup u \text{ weakly in } H^{1,2}(P_\alpha), \text{ as } n \rightarrow \infty.$$

REMARK 2.1. In Lemma 2.5, we can replace P_α by Q_α with the help of the change of variable ψ defined above.

3. A priori estimate

Now, we shall prove an *a priori* estimate which will allow us to take limits in α_n . We denote by $u_n \in H^{1,2}(Q_{\alpha_n})$ the solution of Problem (P) corresponding to the right-hand side $f_n = f|_{Q_{\alpha_n}} \in L^2(Q_{\alpha_n})$ in

$$Q_{\alpha_n} = \Omega_{\alpha_n} \times]0, b[,$$

where

$$\Omega_{\alpha_n} = \{(t, x_1) \in \mathbb{R}^2 : \alpha_n < t < T - \alpha_n, \varphi_1(t) < x_1 < \varphi_2(t)\},$$

with $(\alpha_n)_n$ a sequence decreasing to zero.

PROPOSITION 3.1. *There exists a constant K_1 independent of n such that*

$$\|u_n\|_{H^{1,2}(Q_{\alpha_n})} \leq K_1 \|f_n\|_{L^2(Q_{\alpha_n})} \leq K_1 \|f\|_{L^2(Q)}.$$

In order to prove Proposition 3.1, we need some preliminary results.

LEMMA 3.1. *Let $] \alpha, \beta[\subset \mathbb{R}$. There exists a constant K_2 (independent of α and β) such that*

$$\|u^{(j)}\|_{L^2(] \alpha, \beta])}^2 \leq (\beta - \alpha)^{2(2-j)} K_2 \|u^{(2)}\|_{L^2(] \alpha, \beta])}^2, \quad j = 0, 1,$$

for every $u \in H^2(\alpha, \beta) \cap H_0^1(\alpha, \beta)$, where $u^{(1)}$ (respectively $u^{(2)}$) is the first (respectively the second) derivative of u on $] \alpha, \beta[$ and $u^{(0)} = u$.

The proof of this inequality may be found, for instance, in Nečas [11].

LEMMA 3.2. For every $\varepsilon > 0$, chosen such that $(\varphi_2(t) - \varphi_1(t)) \leq \varepsilon$, there exists a constant C_1 independent of n such that

$$\|D_{x_1}^j u_n\|_{L^2(Q_{\alpha_n})}^2 \leq C_1 \varepsilon^{2(2-j)} \|D_{x_1}^2 u_n\|_{L^2(Q_{\alpha_n})}^2, j = 0, 1.$$

Proof. Replacing in Lemma 3.1 u by u_n and (α, β) by $(\varphi_1(t), \varphi_2(t))$, for a fixed t , we obtain

$$\begin{aligned} \int_{\varphi_1(t)}^{\varphi_2(t)} (D_{x_1}^j u_n)^2 dx_1 &\leq K_2 (\varphi_2(t) - \varphi_1(t))^{2(2-j)} \int_{\varphi_1(t)}^{\varphi_2(t)} (D_{x_1}^2 u_n)^2 dx_1 \\ &\leq K_2 \varepsilon^{2(2-j)} \int_{\varphi_1(t)}^{\varphi_2(t)} (D_{x_1}^2 u_n)^2 dx_1. \end{aligned}$$

Integrating in the previous inequality with respect to t , then with respect to x_2 , we get the desired result with $C_1 = K_2$.

Proof of Proposition 3.1. Let us denote the inner product in $L^2(Q_{\alpha_n})$ by $\langle \cdot, \cdot \rangle$, then we have

$$\begin{aligned} \|f_n\|_{L^2(Q_{\alpha_n})}^2 &= \langle D_t u_n - D_{x_1}^2 u_n - D_{x_2}^2 u_n, D_t u_n - D_{x_1}^2 u_n - D_{x_2}^2 u_n \rangle \\ &= \|D_t u_n\|_{L^2(Q_{\alpha_n})}^2 + \|D_{x_1}^2 u_n\|_{L^2(Q_{\alpha_n})}^2 + \|D_{x_2}^2 u_n\|_{L^2(Q_{\alpha_n})}^2 \\ &\quad - 2\langle D_t u_n, D_{x_1}^2 u_n \rangle - 2\langle D_t u_n, D_{x_2}^2 u_n \rangle + 2\langle D_{x_1}^2 u_n, D_{x_2}^2 u_n \rangle. \end{aligned}$$

1) *Estimation of $-2\langle D_t u_n, D_{x_1}^2 u_n \rangle$.* We have

$$D_t u_n D_{x_1}^2 u_n = D_{x_1} (D_t u_n D_{x_1} u_n) - \frac{1}{2} D_t (D_{x_1} u_n)^2.$$

Then

$$\begin{aligned} -2\langle D_t u_n, D_{x_1}^2 u_n \rangle &= -2 \int_{Q_{\alpha_n}} D_t u_n D_{x_1}^2 u_n dt dx_1 dx_2 \\ &= -2 \int_{Q_{\alpha_n}} D_{x_1} (D_t u_n D_{x_1} u_n) dt dx_1 dx_2 \\ &\quad + \int_{Q_{\alpha_n}} D_t (D_{x_1} u_n)^2 dt dx_1 dx_2 \\ &= \int_{\partial Q_{\alpha_n}} \left[(D_{x_1} u_n)^2 v_t - 2D_t u_n D_{x_1} u_n v_{x_1} \right] d\sigma, \end{aligned}$$

where v_t, v_{x_1}, v_{x_2} are the components of the unit outward normal vector at ∂Q_{α_n} . We shall rewrite the boundary integral making use of the boundary conditions. On the parts of the boundary of Q_{α_n} where $t = \alpha_n, x_2 = 0$ and $x_2 = b$, we have $u_n = 0$ and consequently $D_{x_1} u_n = 0$. The corresponding boundary integral vanishes. On the part

of the boundary where $t = T - \alpha_n$, we have $v_{x_1} = 0$ and $v_t = 1$. Accordingly the corresponding boundary integral

$$A = \int_0^b \int_{\varphi_1(T-\alpha_n)}^{\varphi_2(T-\alpha_n)} (D_{x_1} u_n)^2 dx_1 dx_2$$

is nonnegative. On the part of the boundary where $x_1 = \varphi_i(t)$, $i = 1, 2$, we have $u_n = 0$. By differentiating with respect to t we obtain

$$D_t u_n = -\varphi'_i(t) D_{x_1} u_n.$$

Consequently, the corresponding boundary integrals I_1 and I_2 are the following:

$$I_1 = - \int_0^b \int_{\alpha_n}^{T-\alpha_n} \varphi'_1(t) [D_{x_1} u_n(t, \varphi_1(t), x_2)]^2 dt dx_2$$

$$I_2 = \int_0^b \int_{\alpha_n}^{T-\alpha_n} \varphi'_2(t) [D_{x_1} u_n(t, \varphi_2(t), x_2)]^2 dt dx_2.$$

We have

$$-2\langle D_t u_n, D_{x_1}^2 u_n \rangle \geq -|I_1| - |I_2|. \tag{3.1}$$

LEMMA 3.3. *There exists a constant K_4 independent of n such that*

$$|I_i| \leq K_4 \varepsilon \|D_{x_1}^2 u_n\|_{L^2(Q_{\alpha_n})}^2, \quad i = 1, 2.$$

Proof. We convert the boundary integral I_1 into a surface integral by setting

$$\begin{aligned} [D_{x_1} u_n(t, \varphi_1(t), x_2)]^2 &= - \frac{\varphi_2(t) - x_1}{\varphi_2(t) - \varphi_1(t)} [D_{x_1} u_n(t, x_1, x_2)]^2 \Big|_{x_1=\varphi_1(t)}^{x_1=\varphi_2(t)} \\ &= - \int_{\varphi_1(t)}^{\varphi_2(t)} D_{x_1} \left\{ \frac{\varphi_2(t) - x_1}{\varphi_2(t) - \varphi_1(t)} [D_{x_1} u_n]^2 \right\} dx_1 \\ &= -2 \int_{\varphi_1(t)}^{\varphi_2(t)} \frac{\varphi_2(t) - x_1}{\varphi_2(t) - \varphi_1(t)} D_{x_1} u_n \cdot D_{x_1}^2 u_n dx_1 \\ &\quad + \int_{\varphi_1(t)}^{\varphi_2(t)} \frac{1}{\varphi_2(t) - \varphi_1(t)} [D_{x_1} u_n]^2 dx_1. \end{aligned}$$

Then, we have

$$I_1 = - \int_0^b \int_{\alpha_n}^{T-\alpha_n} \varphi'_1(t) [D_{x_1} u_n(t, \varphi_1(t), x_2)]^2 dt dx_2$$

$$= - \int_{Q_{\alpha_n}} \frac{\varphi'_1(t)}{\varphi_2(t) - \varphi_1(t)} [D_{x_1} u_n(t, x_1, x_2)]^2 dt dx_1 dx_2$$

$$+ 2 \int_{Q_{\alpha_n}} \frac{\varphi_2(t) - x_1}{\varphi_2(t) - \varphi_1(t)} \varphi'_1(t) (D_{x_1} u_n) (D_{x_1}^2 u_n) dt dx_1 dx_2.$$

Thanks to Lemma 3.2, we can write

$$\int_{\varphi_1(t)}^{\varphi_2(t)} [D_{x_1} u_n]^2 dx_1 \leq K_2 [\varphi_2(t) - \varphi_1(t)]^2 \int_{\varphi_1(t)}^{\varphi_2(t)} [D_{x_1}^2 u_n]^2 dx_1.$$

Therefore

$$\int_{\varphi_1(t)}^{\varphi_2(t)} [D_{x_1} u_n]^2 \frac{|\varphi_1'|}{\varphi_2 - \varphi_1} dx_1 \leq K_2 |\varphi_1'| [\varphi_2 - \varphi_1] \int_{\varphi_1(t)}^{\varphi_2(t)} [D_{x_1}^2 u_n]^2 dx_1,$$

consequently

$$\begin{aligned} |I_1| &\leq K_2 \int_{Q_{\varphi_n}} |\varphi_1'| [\varphi_2 - \varphi_1] (D_{x_1}^2 u_n)^2 dt dx_1 dx_2 \\ &\quad + 2 \int_{Q_{\varphi_n}} |\varphi_1'| |D_{x_1} u_n| |D_{x_1}^2 u_n| dt dx_1 dx_2, \end{aligned}$$

since $\left| \frac{\varphi_2(t) - x_1}{\varphi_2(t) - \varphi_1(t)} \right| \leq 1$. Using the inequality

$$2|\varphi_1' D_{x_1} u_n| |D_{x_1}^2 u_n| \leq \varepsilon (D_{x_1}^2 u_n)^2 + \frac{1}{\varepsilon} (\varphi_1')^2 (D_{x_1} u_n)^2$$

for all $\varepsilon > 0$, we obtain

$$\begin{aligned} |I_1| &\leq K_2 \int_{Q_{\varphi_n}} |\varphi_1'| [\varphi_2 - \varphi_1] (D_{x_1}^2 u_n)^2 dt dx_1 dx_2 \\ &\quad + \int_{Q_{\varphi_n}} \varepsilon (D_{x_1}^2 u_n)^2 dt dx_1 dx_2 + \frac{1}{\varepsilon} \int_{Q_{\varphi_n}} (\varphi_1')^2 (D_{x_1} u_n)^2 dt dx_1 dx_2. \end{aligned}$$

Lemma 3.2 yields

$$\frac{1}{\varepsilon} \int_{Q_{\varphi_n}} (\varphi_1')^2 (D_{x_1} u_n)^2 dt dx_1 dx_2 \leq K_2 \frac{1}{\varepsilon} \int_{Q_{\varphi_n}} (\varphi_1')^2 [\varphi_2 - \varphi_1]^2 (D_{x_1}^2 u_n)^2 dt dx_1 dx_2.$$

Thus,

$$\begin{aligned} |I_1| &\leq K_2 \int_{Q_{\varphi_n}} \left[|\varphi_1'| |\varphi_2 - \varphi_1| + \frac{1}{\varepsilon} (\varphi_1')^2 |\varphi_2 - \varphi_1|^2 \right] (D_{x_1}^2 u_n)^2 dt dx_1 dx_2 \\ &\quad + \int_{Q_{\varphi_n}} \varepsilon (D_{x_1}^2 u_n)^2 dt dx_1 dx_2 \\ &\leq (2K_2 + 1) \varepsilon \int_{Q_{\varphi_n}} (D_{x_1}^2 u_n)^2 dt dx_1 dx_2, \end{aligned}$$

since $|\varphi_1' (\varphi_2 - \varphi_1)| \leq \varepsilon$. Finally, taking $K_4 = (2K_2 + 1)$, we obtain

$$|I_1| \leq K_4 \varepsilon \|D_{x_1}^2 u_n\|_{L^2(Q_{\varphi_n})}.$$

The inequality

$$|I_2| \leq K_4 \varepsilon \|D_{x_1}^2 u_n\|_{L^2(Q_{\alpha_n})},$$

can be proved by a similar argument.

2) *Estimation of* $-2\langle D_t u_n, D_{x_2}^2 u_n \rangle$. We have

$$D_t u_n D_{x_2}^2 u_n = D_{x_2} (D_t u_n D_{x_2} u_n) - \frac{1}{2} D_t (D_{x_2} u_n)^2.$$

Then

$$\begin{aligned} -2\langle D_t u_n, D_{x_2}^2 u_n \rangle &= -2 \int_{Q_{\alpha_n}} D_t u_n D_{x_2}^2 u_n dt dx_1 dx_2 \\ &= -2 \int_{Q_{\alpha_n}} D_{x_2} (D_t u_n D_{x_2} u_n) dt dx_1 dx_2 \\ &\quad + \int_{Q_{\alpha_n}} D_t (D_{x_2} u_n)^2 dt dx_1 dx_2 \\ &= \int_{\partial Q_{\alpha_n}} \left[(D_{x_2} u_n)^2 v_t - 2D_t u_n D_{x_2} u_n v_{x_2} \right] d\sigma. \end{aligned}$$

Using the Cauchy-Dirichlet boundary conditions, we see that the above boundary integral is nonnegative. Consequently

$$-2\langle D_t u_n, D_{x_2}^2 u_n \rangle \geq 0. \quad (3.2)$$

3) *Estimation of* $2\langle D_{x_1}^2 u_n, D_{x_2}^2 u_n \rangle$. We have

$$D_{x_1}^2 u_n \cdot D_{x_2}^2 u_n = D_{x_1} (D_{x_1} u_n \cdot D_{x_2}^2 u_n) - D_{x_2} (D_{x_1} u_n \cdot D_{x_1} D_{x_2} u_n) + (D_{x_1} D_{x_2} u_n)^2.$$

Then

$$\begin{aligned} 2\langle D_{x_1}^2 u_n, D_{x_2}^2 u_n \rangle &= 2 \int_{Q_{\alpha_n}} D_{x_1}^2 u_n \cdot D_{x_2}^2 u_n dt dx_1 dx_2 \\ &= 2 \int_{Q_{\alpha_n}} D_{x_1} (D_{x_1} u_n \cdot D_{x_2}^2 u_n) dt dx_1 dx_2 \\ &\quad - 2 \int_{Q_{\alpha_n}} D_{x_2} (D_{x_1} u_n \cdot D_{x_1} D_{x_2} u_n) dt dx_1 dx_2 \\ &\quad + 2 \int_{Q_{\alpha_n}} (D_{x_1} D_{x_2} u_n)^2 dt dx_1 dx_2 \\ &= 2 \int_{Q_{\alpha_n}} (D_{x_1} D_{x_2} u_n)^2 dt dx_1 dx_2 \\ &\quad + 2 \int_{\partial Q_{\alpha_n}} [D_{x_1} u_n D_{x_2}^2 u_n v_{x_1} - D_{x_1} u_n \cdot D_{x_1} D_{x_2} u_n v_{x_2}] d\sigma. \end{aligned}$$

Thanks to the boundary conditions, the above boundary integral vanishes. Consequently

$$2\langle D_{x_1}^2 u_n, D_{x_2}^2 u_n \rangle = 2 \|D_{x_1} D_{x_2} u_n\|_{L^2(Q_{\alpha_n})}^2. \quad (3.3)$$

Summing up the estimates (3.1), (3.2) and (3.3) of the inner products and making use of Lemma 3.3, we then obtain

$$\begin{aligned} \|f_n\|_{L^2(Q_{\alpha_n})}^2 &\geq \|D_t u_n\|_{L^2(Q_{\alpha_n})}^2 + \|D_{x_1}^2 u_n\|_{L^2(Q_{\alpha_n})}^2 + \|D_{x_2}^2 u_n\|_{L^2(Q_{\alpha_n})}^2 \\ &\quad - |I_1| - |I_2| + 2 \|D_{x_1} D_{x_2} u_n\|_{L^2(Q_{\alpha_n})}^2 \\ &\geq \|D_t u_n\|_{L^2(Q_{\alpha_n})}^2 + (1 - 2K_4\varepsilon) \|D_{x_1}^2 u_n\|_{L^2(Q_{\alpha_n})}^2 \\ &\quad + \|D_{x_2}^2 u_n\|_{L^2(Q_{\alpha_n})}^2 + 2 \|D_{x_1} D_{x_2} u_n\|_{L^2(Q_{\alpha_n})}^2. \end{aligned}$$

Then, it is sufficient to choose ε such that $(1 - 2K_4\varepsilon) > 0$ to get a constant $K_0 > 0$ independent of n such that

$$\|f_n\|_{L^2(Q_{\alpha_n})} \geq K_0 \|u_n\|_{H^{1,2}(Q_{\alpha_n})},$$

and since

$$\|f_n\|_{L^2(Q_{\alpha_n})} \leq \|f\|_{L^2(Q)},$$

there exists a constant $K_1 > 0$, independent of n satisfying

$$\|u_n\|_{H^{1,2}(Q_{\alpha_n})} \leq K_1 \|f_n\|_{L^2(Q_{\alpha_n})} \leq K_1 \|f\|_{L^2(Q)}.$$

This completes the proof of Proposition 3.1.

4. Passage to the limit

We are now able to prove the main result of this work

THEOREM 4.1. *Assume that φ_1 and φ_2 fulfil conditions (1.1) and (1.2). Then the heat operator*

$$L = D_t - D_{x_1}^2 - D_{x_2}^2$$

is an isomorphism from $H_V^{1,2}(Q)$ into $L^2(Q)$.

Proof. Choose a sequence Q_{α_n} $n = 1, 2, \dots$ of reference domains (see Section 2) such that $Q_{\alpha_n} \subseteq Q$ with (α_n) a sequence decreasing to 0, as $n \rightarrow \infty$. Then we have $Q_{\alpha_n} \rightarrow Q$, as $n \rightarrow \infty$.

Consider the solution $u_{\alpha_n} \in H^{1,2}(Q_{\alpha_n})$ of the Cauchy-Dirichlet problem

$$f(x) = \begin{cases} D_t u_{\alpha_n} - D_{x_1}^2 u_{\alpha_n} - D_{x_2}^2 u_{\alpha_n} = f & \text{in } Q_{\alpha_n}, \\ u_{\alpha_n} / \partial Q - \Gamma_{T-\alpha_n} = 0, \end{cases}$$

with $\Gamma_{T-\alpha_n}$ is the part of the boundary of Q_{α_n} where $t = T - \alpha_n$. Such a solution u_{α_n} exists by Theorem 2.1. Let $\widetilde{u_{\alpha_n}}$ the 0-extension of u_{α_n} to Q . In virtue of Proposition 3.1, we know that there exists a constant C such that

$$\|\widetilde{u_{\alpha_n}}\|_{L^2(Q)} + \|\widetilde{D_t u_{\alpha_n}}\|_{L^2(Q)} + \sum_{\substack{i,j=0 \\ 1 \leq i+j \leq 2}}^2 \left\| \widetilde{D_{x_1}^i D_{x_2}^j u_{\alpha_n}} \right\|_{L^2(Q)} \leq C \|f\|_{L^2(Q)}.$$

This means that $\widetilde{u_{\alpha_n}}, \widetilde{D_t u_{\alpha_n}}, \widetilde{D_{x_1}^i D_{x_2}^j u_{\alpha_n}}$ for $1 \leq i + j \leq 2$ are bounded functions in $L^2(Q)$. So for a suitable increasing sequence of integers $n_k, k = 1, 2, \dots$, there exist functions

$$u, v \text{ and } v_{i,j}, \quad 1 \leq i + j \leq 2$$

in $L^2(Q)$ such that

$$\begin{aligned} \widetilde{u_{\alpha_{n_k}}} &\rightarrow u \text{ weakly in } L^2(Q), \quad k \rightarrow \infty, \\ \widetilde{D_t u_{\alpha_{n_k}}} &\rightarrow v \text{ weakly in } L^2(Q), \quad k \rightarrow \infty, \\ \widetilde{D_{x_1}^i D_{x_2}^j u_{\alpha_{n_k}}} &\rightarrow v_{i,j} \text{ weakly in } L^2(Q), \quad 1 \leq i + j \leq 2, k \rightarrow \infty. \end{aligned}$$

Clearly,

$$v = D_t u, \quad v_{i,j} = D_{x_1}^i D_{x_2}^j u, \quad 1 \leq i + j \leq 2$$

in the sense of distributions in Q . So, $u \in H^{1,2}(Q)$ and

$$D_t u - D_{x_1}^2 u - D_{x_2}^2 u = f \text{ in } Q.$$

On the other hand, the solution u satisfies the boundary conditions $u|_{\partial Q - \Gamma_T} = 0$ since

$$\forall n \in \mathbb{N}, u|_{Q_{\alpha_n}} = u_{\alpha_n}.$$

This proves the existence of a solution to Problem (P).

Notice that we have the estimate

$$\|u\|_{H^{1,2}(Q)}^2 \leq K \|f\|_{L^2(Q)}^2,$$

which implies the uniqueness of the solution.

REMARK 4.1. The result given in Theorem 4.1 holds true only under the assumption (1.1) (respectively, (1.2)), if $\varphi_1(0) = \varphi_2(0)$ and $\varphi_1(T) < \varphi_2(T)$ (respectively, if $\varphi_1(0) < \varphi_2(0)$ and $\varphi_1(T) = \varphi_2(T)$).

REMARK 4.2. Note that this work may be extended at least in the following directions:

1. The nonregular domain Q may be replaced by a non-cylindrical domain (conical domain, for example).
2. The function f on the right-hand side of the equation of Problem (P), may be taken in $L^p(\Omega)$, where $p \in]1, \infty[$. The method used here does not seem to be appropriate for the space $L^p(\Omega)$ when $p \neq 2$.
3. The operator L may be replaced by a high order operator.

REFERENCES

- [1] YU. A. ALKHUTOV, *L_p - estimates of solutions of the Dirichlet problem for the heat equation in a ball*, Journal of Mathematical Sciences, **142**, 3 (2007), 2021–2032.
- [2] V. N. AREF'EV, L. A. BAGIROV, *Solutions of the heat equation in domains with singularities*, Mathematical Notes, **64**, 2 (1998), 139–153.
- [3] E. A. BADERKO, *On the solution of the boundary value problems for parabolic equations of high order in domains with curvilinear lateral boundaries*, Diff. Urav., **12**, 10 (1976), 1781–1792.
- [4] V. BESOV, *The continuation of function in L_p^1 and W_p^1* , Proc. Steklov Inst. Math., **89** (1967), 5–17.
- [5] P. GRISVARD, *Elliptic Problems in Nonsmooth Domains*, Monographs and Studies in Mathematics 24, Pitman, Boston, 1985.
- [6] S. HOFMANN, J. L. LEWIS, *The L^p regularity problems for the heat equation in non-cylindrical domains*, Journal of Functional Analysis, **220** (2005), 1–54.
- [7] R. LABBAS, A. MEDEGHRI, B.-K. SADALLAH, *On a parabolic equation in a triangular domain*, Applied Mathematics and Computation, **130** (2002), 511–523.
- [8] R. LABBAS, A. MEDEGHRI, B.-K. SADALLAH, *An L^p approach for the study of degenerate parabolic equation*, EJDE, **2005**, 36 (2005), 1–20.
- [9] LADYZHENSKAYA-SOLONNIKOV-URAL'TSEVA, *Linear and Quasi-Linear Equations of Parabolic Type*. A.M.S., providence, Rhode Island, 1968.
- [10] J. L. LIONS, E. MAGENES, *Problèmes aux Limites Non Homogènes et Applications*, 1, 2, Dunod, Paris, 1968.
- [11] J. NEČAS, *Les Méthodes Directes en Théorie des Equations Elliptiques*, Masson, Paris, 1967.
- [12] B.-K. SADALLAH, *Etude d'un Problème $2m$ -parabolique dans des domaines plan non rectangulaires*, Boll. Un. Mat. Ital., **5**, 2-B (1983), 51–112.
- [13] G. SAVARÉ, *Parabolic problems with mixed variable lateral conditions: an abstract approach*, J. Math. Pures Appl., **76**, (1997), 321–351.

(Received May 7, 2009)

Arezki Kheloufi
 Department of Sciences and Techniques, Faculty of Technology
 Béjaia University
 6000. Béjaia
 Algeria
 e-mail: arezkinet2000@yahoo.fr

Rabah Labbas
 Laboratoire de Mathématiques
 Université du Havre, U.F.R.S.T.
 BP 540, 76058 Le Havre Cedex
 France
 e-mail: Rabah.Labbas@univ-lehavre.fr

Boubaker-Khaled Sadallah
 Department of Mathematics
 E.N.S.
 16050 Kouba, Algiers
 Algeria
 e-mail: sadallah@ens-kouba.dz