

## ON WEIGHTED FRACTIONAL INEQUALITIES USING GENERALIZED KATUGAMPOLA FRACTIONAL INTEGRAL OPERATOR

SATISH K. PANCHAL, VAIJANATH L. CHINCHANE AND ASHA B. NALE

(Communicated by S. K. Ntouyas)

*Abstract.* In this paper, we obtain some new weighted fractional inequalities which are presented by M. Houas in the paper (Certain weighted integral inequalities involving fractional hypergeometric operator, *Scientia, series A: Mathematical Science* 27(2016),87-97), using generalized Katugampola fractional integral operator.

### 1. Introduction

Fractional Calculus is traditionally associated to non-integers where the order of derivative or integral is considered to be non-integer. Fractional calculus have found significant and important applications in science and technology. The fractional integral inequalities gained more attention due to it's application in continuation solution, uniqueness of solution of fractional differential equations. During the past few years, many researchers have investigated well known fractional integral inequalities and its applications using Riemann-Liouville, Hadamard, Saigo, Erdélyi-Kober, Katugampola, k-fractional integral, k-Hadamard integral and generalized k-fractional integral, see [2, 3, 4, 5, 6, 8, 9, 12, 13, 14, 15, 17, 18, 21]. In [7], Curiel and Galue introduced the Gauss hypergeometric function operator. Recently, V. L. Chinchane, et al. [4], Baleanu et al. [3] proposed fractional integral inequalities using the generalized k-fractional integral operator in terms of the Gauss hypergeometric functions. In [10], M. Houas obtained certain weighted integral inequalities involving the fractional hypergeometric operators.

Recently, A. B. Nale, et al. [16] investigated new fractional integral inequalities for convex functions using generalized Katugampola fractional integral. T. A Aljaaidi, et al. [1] and J. V. Sousa et al. [22] have established Grüss-type inequalities using generalized Katugampola fractional integral. E. Set, et al. [19] established several Chebyshev type inequalities using generalized Katugampola fractional integral operator. Motivated from above work, the aim of this paper is to obtain some new weighted fractional integral inequalities using generalized Katugampola fractional integral operators. The paper has been organized as follows. In Section 2, we define basic definitions and proposition related to generalized Katugampola fractional derivatives and integrals. In Section 3, we give weighted fractional integral inequalities by employing generalized Katugampola fractional integral operator. In section 4, we give concluding remarks.

*Mathematics subject classification* (2010): 26D10, 26A33, 05A30.

*Keywords and phrases:* Inequalities, generalized Katugampola fractional integral operator.

## 2. Preliminaries

In this section, we give some basic definition and mathematical preliminaries of Generalized Katugampola fractional integral, see [1, 13, 19, 22].

DEFINITION 2.1. Consider the space  $X_c^p(a, b)$  ( $c \in \mathbb{R}, 1 \leq p < \infty$ ), of those complex valued Lebesgue measurable functions  $f$  on  $(a, b)$  for which the norm  $\|f\|_{X_c^p} < \infty$ , such that

$$\|f\|_{X_c^p} = \left( \int_a^b |x^c f|^p \frac{dx}{x} \right)^{\frac{1}{p}}, \quad (1 \leq p < \infty),$$

and

$$\|f\|_{X_c^p} = \sup_{x \in (a, b)} [x^c |f|].$$

In particular, when  $c = \frac{1}{p}$ , the space  $X_c^p(a, b)$  coincides with the space  $L^p(a, b)$ .

DEFINITION 2.2. The left and right sided fractional integrals of a function  $f$  where  $f \in X_c^p(a, b)$ ,  $\alpha > 0$  and  $\beta, \rho, \eta, k \in \mathbb{R}$ , are defined respectively by

$${}^\rho \mathcal{J}_{a+; \eta, k}^{\alpha, \beta} f(x) = \frac{\rho^{1-\beta} x^k}{\Gamma(\alpha)} \int_a^x \frac{\tau^{\rho(\eta+1)-1}}{(x^\rho - \tau^\rho)^{1-\alpha}} f(\tau) d\tau, \quad 0 \leq a < x < b \leq \infty, \quad (2.1)$$

and

$${}^\rho \mathcal{J}_{b-; \eta, k}^{\alpha, \beta} f(x) = \frac{\rho^{1-\beta} x^{k\rho}}{\Gamma(\alpha)} \int_x^b \frac{\tau^{k+\rho-1}}{(\tau^\rho - x^\rho)^{1-\alpha}} f(\tau) d\tau, \quad 0 \leq a < x < b \leq \infty, \quad (2.2)$$

if the integral exist.

To represent and discuss our new results in this paper we use the left sided fractional integrals, the right sided fractional can be proved similarly, also we consider  $a = 0$ , in (2.1), to obtain

$${}^\rho \mathcal{J}_{\eta, k}^{\alpha, \beta} f(x) = \frac{\rho^{1-\beta} x^k}{\Gamma(\alpha)} \int_0^x \frac{\tau^{\rho(\eta+1)-1}}{(x^\rho - \tau^\rho)^{1-\alpha}} f(\tau) d\tau. \quad (2.3)$$

The above fractional integrals has the following composition (index) formulae

$$\begin{aligned} {}^\rho \mathcal{J}_{a+; \eta_1, k_1}^{\alpha_1, \beta_1} {}^\rho \mathcal{J}_{a+; \eta_2, -\rho\eta_1}^{\alpha_2, \beta_2} f(x) &= {}^\rho \mathcal{J}_{a+; \eta_2, k_1}^{\alpha_1+\alpha_2, \beta_1+\beta_2} f(x), \\ {}^\rho \mathcal{J}_{b-; \eta_1, -\rho\eta_2}^{\alpha_1, \beta_1} {}^\rho \mathcal{J}_{b-; \eta_2, k_2}^{\alpha_2, \beta_2} f(x) &= {}^\rho \mathcal{J}_{a+; \eta_1, k_2}^{\alpha_1+\alpha_2, \beta_1+\beta_2} f(x). \end{aligned} \quad (2.4)$$

Here, we recall definition of beta function as:

DEFINITION 2.3. The beta function  $B(\alpha, \beta)$  is defined ([20], section 1.1)

$$B(\alpha, \beta) = \begin{cases} \int_0^1 t^{\alpha-1} (1-t)^{\beta-1} dt & (\Re(\alpha) > 0; \Re(\beta) > 0) \\ \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)} & (\alpha, \beta \in \mathbb{C}/\mathbb{Z}_0^-) \end{cases} \quad (2.5)$$

For the convenience of establishing our results we define the following function as in [19, 22], using (2.5) and let  $x > 0, \alpha > 0, \rho, k, \beta, \eta \in \mathbb{R}$ , then

$$\Lambda_{x,k}^{\rho,\beta}(\alpha, \eta) = \frac{\rho^{1-\beta} x^k}{\Gamma(\alpha)} \int_0^x \frac{\tau^{\rho(\eta+1)-1}}{(x^\rho - \tau^\rho)^{1-\alpha}} d\tau = \frac{\Gamma(\eta+1)}{\Gamma(\eta+\alpha+1)} \rho^{-\beta} x^{k+\rho(\eta+\alpha)}. \quad (2.6)$$

REMARK 2.1. The fractional integral (2.1) contain five well-known fractional integral as its particular cases, see [1, 12, 13, 16, 19, 22].

1. Setting  $k = 0, \eta = 0, a = 0$  and taking the limit  $\rho \rightarrow 1$  in (2.1), the integral operator (2.1) reduces to the Riemann-Liouville fractional integral.
2. Setting  $k = 0, \eta = 0$  and taking the limit  $\rho \rightarrow 1$  in (2.1), the integral operator (2.1) reduces to the Liouville fractional integral.
3. Setting  $\beta = \alpha, k = 0, \eta = 0$  and taking the limit  $\rho \rightarrow 0^+$  with L' Hospital rule in (2.1), the integral operator (2.1) reduces to the Hadamard fractional integral see [11].
4. Setting  $\beta = 0, k = -\rho(\alpha + \eta)$  in (2.1), the integral operator (2.1) reduces to the Erdelyi-Kober fractional integral.
5. Setting  $\beta = \alpha, k = 0$  and  $\eta = 0$  in (2.1), the integral operator (2.1) reduces to the Katugampola fractional integral.

### 3. Main results

Here, we obtain new fractional integral inequalities using generalized Katugampola fractional integral operators.

THEOREM 3.1. *Let  $f$  be positive and continuous function on  $[0, \infty)$ , such that*

$$(\sigma^\xi f^\xi(\tau) - \tau^\xi f^\xi(\sigma))(f^{\varpi-\lambda}(\tau) - f^{\varpi-\lambda}(\sigma)) \geq 0, \quad (3.1)$$

and  $w : [0, \infty) \rightarrow \mathbb{R}^+$  be positive continuous function. Then for all  $\alpha \geq 0, x > 0, \beta, \rho, \eta, k \in \mathbb{R}$ , we have

$$\begin{aligned} & \rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) f^{\xi+\lambda}(x)]^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) x^\xi f^\varpi(x)] \\ & \leq \rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) f^{\xi+\varpi}(x)]^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) x^\xi f^\lambda(x)]. \end{aligned} \quad (3.2)$$

*Proof.* Since  $f$  be positive and continuous function on  $[0, \infty)$ , then for all  $\xi > 0, \varpi \geq 0, \lambda > 0, \tau, \sigma \in (0, x), x > 0$ , from (3.1)

$$\begin{aligned} & \sigma^\xi f^{\varpi-\lambda}(\sigma) f^\xi(\tau) + \tau^\xi f^{\varpi-\lambda}(\tau) f^\xi(\sigma) \\ & \leq \sigma^\xi f^{\varpi+\xi-\lambda}(\tau) + \tau^\xi f^{\varpi+\xi-\lambda}(\sigma). \end{aligned} \quad (3.3)$$

Multiplying both sides of equation (3.3) by  $\frac{\rho^{1-\beta}x^k}{\Gamma(\alpha)} \frac{\tau^{\rho(\eta+1)-1}}{(x^\rho - \tau^\rho)^{1-\alpha}} w(\tau) f^\lambda(\tau)$ ,  $\tau \in (0, x)$ ,  $x > 0$  which is positive, and integrating the obtained result with respect to  $\tau$  from 0 to  $x$ , we get

$$\begin{aligned} & \frac{\rho^{1-\beta}x^k}{\Gamma(\alpha)} \int_0^x \frac{\tau^{\rho(\eta+1)-1}}{(x^\rho - \tau^\rho)^{1-\alpha}} \sigma^\xi f^{\varpi-\lambda}(\sigma) w(\tau) f^{\lambda+\xi}(\tau) d\tau \\ & + \frac{\rho^{1-\beta}x^k}{\Gamma(\alpha)} \int_0^x \frac{\tau^{\rho(\eta+1)-1}}{(x^\rho - \tau^\rho)^{1-\alpha}} \tau^\xi f^{\varpi}(\tau) f^\xi(\sigma) w(\tau) d\tau \\ & \leq \frac{\rho^{1-\beta}x^k}{\Gamma(\alpha)} \int_0^x \frac{\tau^{\rho(\eta+1)-1}}{(x^\rho - \tau^\rho)^{1-\alpha}} \sigma^\xi f^{\varpi+\xi-\lambda}(\tau) w(\tau) f^\lambda(\tau) d\tau \\ & + \frac{\rho^{1-\beta}x^k}{\Gamma(\alpha)} \int_0^x \frac{\tau^{\rho(\eta+1)-1}}{(x^\rho - \tau^\rho)^{1-\alpha}} \tau^\xi f^{\varpi+\xi-\lambda}(\sigma) w(\tau) f^\lambda(\tau) d\tau, \end{aligned} \quad (3.4)$$

consequently,

$$\begin{aligned} & \sigma^\xi f^{\varpi-\lambda}(\sigma) {}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) f^{\lambda+\xi}(x)] + f^\xi(\sigma) {}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) \tau^\xi f^{\varpi}(x)] \\ & \geq \sigma^\xi {}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) f^{\xi+\varpi}(x)] + f^{\xi+\varpi-\lambda}(\sigma) {}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) x^\xi f^\lambda(x)]. \end{aligned} \quad (3.5)$$

Multiplying both sides of equation (3.5) by  $\frac{\rho^{1-\beta}x^k}{\Gamma(\alpha)} \frac{\sigma^{\rho(\eta+1)-1}}{(x^\rho - \sigma^\rho)^{1-\alpha}} w(\sigma) f^\lambda(\sigma)$ ,  $\sigma \in (0, x)$ ,  $x > 0$  which is positive and integrating the obtained result with respect to  $\sigma$  from 0 to  $x$ , we get

$$\begin{aligned} & {}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) f^{\xi+\lambda}(x)] {}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) x^\xi f^{\varpi}(x)] \\ & + {}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) x^\xi f^{\varpi}(x)] {}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) f^{\xi+\lambda}(x)] \\ & \geq {}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) f^{\xi+\varpi}(x)] {}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) x^\xi f^\lambda(x)] \\ & + {}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) x^\xi f^\lambda(x)] {}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) f^{\xi+\varpi}(x)]. \end{aligned} \quad (3.6)$$

This completes the proof of inequality (3.2).  $\square$

Now, we give our main result.

**THEOREM 3.2.** *Let  $f$  be positive and continuous function on  $[0, \infty)$  and satisfies (3.1). Let  $w : [0, \infty) \rightarrow \mathbb{R}^+$  be positive continuous function. Then for all  $\alpha, \theta \geq 0$ ,  $x > 0$ ,  $\beta, \pi, \rho, \eta, k \in \mathbb{R}$ , we have*

$$\begin{aligned} & {}^\rho \mathcal{J}_{\eta,k}^{\theta,\pi} [w(x) x^\xi f^{\varpi}(x)] {}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) f^{\xi+\lambda}(x)] \\ & + {}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) x^\xi f^{\varpi}(x)] {}^\rho \mathcal{J}_{\eta,k}^{\theta,\pi} [w(x) f^{\xi+\lambda}(x)] \\ & \leq {}^\rho \mathcal{J}_{\eta,k}^{\theta,\pi} [w(x) x^\xi f^\lambda(x)] {}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) f^{\xi+\varpi}(x)] \\ & + {}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) x^\xi f^\lambda(x)] {}^\rho \mathcal{J}_{\eta,k}^{\theta,\pi} [w(x) f^{\xi+\varpi}(x)]. \end{aligned} \quad (3.7)$$

*Proof.* Now multiplying both sides of (3.3) by  $\frac{\rho^{1-\pi}x^k}{\Gamma(\theta)} \frac{\sigma^{\rho(\eta+1)-1}}{(x^\rho - \sigma^\rho)^{1-\theta}} w(\sigma) f^\lambda(\sigma)$ ,  $\sigma \in (0, x)$ ,  $x > 0$ , this function remains positive under the conditions stated with the theorem. Integrating the obtained result with respect to  $\sigma$  from 0 to  $x$ , we get

$$\begin{aligned} & \frac{f^{\overline{\omega}}(\tau) \rho^{1-\pi} x^k}{\Gamma(\theta)} \int_0^x \frac{\sigma^{\rho(\eta+1)-1}}{(x^\rho - \sigma^\rho)^{1-\pi}} w(\sigma) \sigma^\xi f^{\overline{\omega}}(\sigma) d\sigma \\ & + \frac{\tau^\xi f^{\overline{\omega}-\lambda}(\tau) \rho^{1-\pi} x^k}{\Gamma(\theta)} \int_0^x \frac{\sigma^{\rho(\eta+1)-1}}{(x^\rho - \sigma^\rho)^{1-\pi}} w(\sigma) f^{\xi+\lambda}(\sigma) d\sigma \\ & \leq \frac{f^{\overline{\omega}+\xi-\lambda}(\tau) \rho^{1-\pi} x^k}{\Gamma(\theta)} \int_0^x \frac{\sigma^{\rho(\eta+1)-1}}{(x^\rho - \sigma^\rho)^{1-\pi}} w(\sigma) \sigma^\xi f^\lambda d\sigma \\ & + \frac{\tau^\xi \rho^{1-\pi} x^k}{\Gamma(\theta)} \int_0^x \frac{\sigma^{\rho(\eta+1)-1}}{(x^\rho - \sigma^\rho)^{1-\pi}} w(\sigma) f^{\overline{\omega}+\xi}(\sigma) d\sigma, \end{aligned} \tag{3.8}$$

consequently

$$\begin{aligned} & f^\xi(\tau)^\rho \mathcal{J}_{\eta,k}^{\theta,\pi} [w(x) x^\xi f^{\overline{\omega}}(x)] + \sigma^\xi f^{\overline{\omega}-\lambda}(x)^\rho \mathcal{J}_{\eta,k}^{\theta,\pi} [w(x) f^{\xi+\lambda}(x)] \\ & \geq f^{\xi+\overline{\omega}-\lambda}(\tau)^\rho \mathcal{J}_{\eta,k}^{\theta,\pi} [w(x) x^\xi f^\lambda(x)] + \tau^\xi \rho \mathcal{J}_{\eta,k}^{\theta,\pi} [w(x) f^{\xi+\overline{\omega}}(x)]. \end{aligned} \tag{3.9}$$

Multiplying both sides of equation (3.9) by  $\frac{\rho^{1-\beta}x^k}{\Gamma(\alpha)} \frac{\tau^{\rho(\eta+1)-1}}{(x^\rho - \tau^\rho)^{1-\alpha}} w(\tau) f^\lambda(\tau)$ ,  $\tau \in (0, x)$ ,  $x > 0$  which is positive, and integrating the obtained result with respect to  $\sigma$  from 0 to  $x$ , we get

$$\begin{aligned} & \rho \mathcal{J}_{\eta,k}^{\theta,\pi} [w(x) x^\xi f^{\overline{\omega}}(x)] \frac{\rho^{1-\beta} x^k}{\Gamma(\alpha)} \int_0^x \frac{\tau^{\rho(\eta+1)-1}}{(x^\rho - \tau^\rho)^{1-\alpha}} \tau^\xi w(\tau) f^{\xi+\lambda}(\tau) d\tau \\ & + \rho \mathcal{J}_{\eta,k}^{\theta,\pi} [w(x) f^{\xi+\lambda}(x)] \frac{\rho^{1-\beta} x^k}{\Gamma(\alpha)} \int_0^x \frac{\tau^{\rho(\eta+1)-1}}{(x^\rho - \tau^\rho)^{1-\alpha}} w(\tau) \tau^\xi f^{\overline{\omega}}(\tau) d\tau \\ & \leq \rho \mathcal{J}_{\eta,k}^{\theta,\pi} [w(x) x^\xi f^\lambda(x)] \frac{\rho^{1-\beta} x^k}{\Gamma(\alpha)} \int_0^x \frac{\tau^{\rho(\eta+1)-1}}{(x^\rho - \tau^\rho)^{1-\alpha}} w(\tau) f^{\overline{\omega}+\xi}(\tau) d\tau \\ & + \rho \mathcal{J}_{\eta,k}^{\theta,\pi} [w(x) f^{\xi+\overline{\omega}}(x)] \frac{\rho^{1-\beta} x^k}{\Gamma(\alpha)} \int_0^x \frac{\tau^{\rho(\eta+1)-1}}{(x^\rho - \tau^\rho)^{1-\alpha}} w(\tau) \tau^\xi f^\lambda(\tau) d\tau, \end{aligned} \tag{3.10}$$

This completes the proof of Theorem 3.2.  $\square$

**THEOREM 3.3.** *Let  $f, h$  be positive and continuous functions on  $[0, \infty)$ , such that*

$$(h^\xi(\sigma) f^\xi(\tau) - h^\xi(\tau) f^\xi(\sigma))(f^{\overline{\omega}-\lambda}(\tau) - f^{\overline{\omega}-\lambda}(\sigma)) \geq 0, \tag{3.11}$$

*and let  $w : [0, \infty) \rightarrow \mathbb{R}^+$  be positive continuous function. Then for all  $\alpha \geq 0$ ,  $x > 0$ ,  $\beta, \rho, \eta, k \in \mathbb{R}$ , we have*

$$\begin{aligned} & \rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) f^\xi(x)]^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) h^\xi(x) f^{\overline{\omega}}(x)] \\ & \leq \rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) f^{\overline{\omega}+\xi}(x)]^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) h^\xi(x) f^\lambda(x)]. \end{aligned} \tag{3.12}$$

*Proof.* Let  $(\tau, \sigma) \in (0, \sigma)$ ,  $x > 0$ , for any  $\varpi > \lambda > 0$ ,  $\xi > 0$ . Then from (3.11)

$$h^\xi(\sigma)f^{\varpi-\lambda}(\sigma)f^\xi(\tau)+h^\xi(\tau)f^\xi(\sigma)f^{\varpi-\lambda}(\tau) \leq h^\xi(\sigma)f^{\varpi+\xi-\lambda}(\tau)+h^\xi(\tau)f^{\varpi+\xi-\lambda}(\sigma). \tag{3.13}$$

Multiplying of both sides of (3.13) by  $\frac{\rho^{1-\beta}x^k}{\Gamma(\alpha)} \frac{\tau^{\rho(\eta+1)-1}}{(x^\rho-\tau^\rho)^{1-\alpha}} w(\tau)f^\lambda(\tau)$ , then integrating the resulting inequality with respect to  $\tau$  over  $(0, x)$ , we obtain

$$\begin{aligned} & h^\xi(\sigma)f^{\varpi-\lambda}(\sigma)^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta}[w(x)f^{\xi+\lambda}(x)] + f^\xi(\sigma)^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta}[w(x)h^\xi(x)f^\varpi(x)] \\ & \leq h^\xi(\sigma)^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta}[w(x)f^{\varpi+\xi}(x)] + f^{\xi+\varpi-\lambda}(\sigma)^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta}[w(x)h^\xi(x)f^\lambda(x)]. \end{aligned} \tag{3.14}$$

Multiplying of both sides of (3.14) by  $\frac{\rho^{1-\beta}x^k}{\Gamma(\alpha)} \frac{\sigma^{\rho(\eta+1)-1}}{(x^\rho-\sigma^\rho)^{1-\alpha}} w(\sigma)f^\lambda(\sigma)$ , then integrating the resulting inequality with respect to  $\sigma$  over  $(0, x)$ , we obtain

$$\begin{aligned} & \rho \mathcal{J}_{\eta,k}^{\alpha,\beta}[w(x)f^{\xi+\lambda}(x)] \frac{\rho^{1-\beta}x^k}{\Gamma(\alpha)} \int_0^x \frac{\sigma^{\rho(\eta+1)-1}}{(x^\rho-\sigma^\rho)^{1-\alpha}} w(\sigma)h^\xi(\sigma)f^\varpi(\sigma)d\sigma \\ & + \rho \mathcal{J}_{\eta,k}^{\alpha,\beta}[w(x)h^\xi(x)f^\varpi(x)] \frac{\rho^{1-\beta}x^k}{\Gamma(\alpha)} \int_0^x \frac{\sigma^{\rho(\eta+1)-1}}{(x^\rho-\sigma^\rho)^{1-\alpha}} f^{\xi+\lambda}(\sigma)w(\sigma)d\sigma \\ & \leq \rho \mathcal{J}_{\eta,k}^{\alpha,\beta}[w(x)f^{\varpi+\xi}(x)] \frac{\rho^{1-\beta}x^k}{\Gamma(\alpha)} \int_0^x \frac{\sigma^{\rho(\eta+1)-1}}{(x^\rho-\sigma^\rho)^{1-\alpha}} f^\lambda(\sigma)w(\sigma)h^\xi(\sigma)d\sigma \\ & + \rho \mathcal{J}_{\eta,k}^{\alpha,\beta}[w(x)h^\xi(x)f^\lambda(x)] \frac{\rho^{1-\beta}x^k}{\Gamma(\alpha)} \int_0^x \frac{\sigma^{\rho(\eta+1)-1}}{(x^\rho-\sigma^\rho)^{1-\alpha}} w(\sigma)f^{\varpi+\xi}(\sigma)d\sigma, \end{aligned} \tag{3.15}$$

which implies that

$$\begin{aligned} & \rho \mathcal{J}_{\eta,k}^{\alpha,\beta}[w(x)f^{\xi+\lambda}(x)]^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta}[w(x)h^\xi(x)f^\varpi(x)] \\ & + \rho \mathcal{J}_{\eta,k}^{\alpha,\beta}[w(x)h^\xi(x)f^\varpi(x)]^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta}[f^{\xi+\lambda}(x)w(x)] \\ & \leq \rho \mathcal{J}_{\eta,k}^{\alpha,\beta}[w(x)f^{\varpi+\xi}(x)]^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta}[f^\lambda(x)w(x)h^\xi(x)] \\ & + \rho \mathcal{J}_{\eta,k}^{\alpha,\beta}[w(x)h^\xi(x)f^\lambda(x)]^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta}[w(x)f^{\varpi+\xi}(x)]. \end{aligned} \tag{3.16}$$

This completes proof of inequality (3.12).  $\square$

**THEOREM 3.4.** Let  $f, h$  be two positive and continuous functions on  $[0, \infty)$  and satisfying (3.11). Let  $w : [0, \infty) \rightarrow \mathbb{R}^+$  be positive continuous function. Then for all  $\alpha, \theta \geq 0$ ,  $x > 0$ ,  $\beta, \pi, \rho, \eta, k \in \mathbb{R}$ , we have

$$\begin{aligned} & \rho \mathcal{J}_{\eta,k}^{\theta,\pi}[w(x)h^\xi(x)f^\varpi(x)]^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta}[w(x)f^{\xi+\lambda}(x)] \\ & + \rho \mathcal{J}_{\eta,k}^{\theta,\pi}[w(x)f^{\xi+\lambda}(x)]^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta}[w(x)h^\xi(x)f^\varpi(x)] \\ & \leq \rho \mathcal{J}_{\eta,k}^{\theta,\pi}[w(x)h^\xi(x)f^\lambda(x)]^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta}[w(x)f^{\varpi+\xi}(x)] \\ & + \rho \mathcal{J}_{\eta,k}^{\theta,\pi}[w(x)f^{\xi+\varpi-\lambda}(x)]^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta}[w(x)h^\xi(x)f^\lambda(x)]. \end{aligned} \tag{3.17}$$

*Proof.* Multiplying the inequality (3.14) by  $\frac{\rho^{1-\pi}x^k}{\Gamma(\theta)} \frac{\sigma^{\rho(\eta+1)-1}}{(x^\rho-\sigma^\rho)^{1-\theta}} w(\sigma) f^\lambda(\sigma)$ ,  $\sigma \in (0, x)$ ,  $x > 0$ , this function remains positive under the conditions stated with the theorem. Integrating the obtained result with respect to  $\sigma$  from 0 to  $x$ , we get

$$\begin{aligned} & \rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) f^{\xi+\lambda}(x)] \frac{\rho^{1-\pi}x^k}{\Gamma(\theta)} \int_0^x \frac{\sigma^{\rho(\eta+1)-1}}{(x^\rho-\sigma^\rho)^{1-\theta}} w(\sigma) h^\xi(\sigma) f^\varpi(\sigma) d\sigma \\ & + \rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) h^\xi(x) f^\varpi(x)] \frac{\rho^{1-\pi}x^k}{\Gamma(\theta)} \int_0^x \frac{\sigma^{\rho(\eta+1)-1}}{(x^\rho-\sigma^\rho)^{1-\theta}} f^{\xi+\lambda}(\sigma) w(\sigma) d\sigma \\ & \leq \rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) f^{\varpi+\xi}(x)] \frac{\rho^{1-\pi}x^k}{\Gamma(\theta)} \int_0^x \frac{\sigma^{\rho(\eta+1)-1}}{(x^\rho-\sigma^\rho)^{1-\theta}} f^\lambda(\sigma) w(\sigma) h^\xi(\sigma) d\sigma \\ & + \rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) h^\xi(x) f^\lambda(x)] \frac{\rho^{1-\pi}x^k}{\Gamma(\theta)} \int_0^x \frac{\sigma^{\rho(\eta+1)-1}}{(x^\rho-\sigma^\rho)^{1-\theta}} w(\sigma) f^{\varpi+\xi}(\sigma) d\sigma, \end{aligned} \tag{3.18}$$

which implies that

$$\begin{aligned} & \rho \mathcal{J}_{\eta,k}^{\theta,\pi} [w(x) h^\xi(x) f^\varpi(x)]^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) f^{\xi+\lambda}(x)] \\ & + \rho \mathcal{J}_{\eta,k}^{\theta,\pi} [w(x) f^{\xi+\lambda}(x)]^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) h^\xi(x) f^\varpi(x)] \\ & \leq \rho \mathcal{J}_{\eta,k}^{\theta,\pi} [w(x) h^\xi(x) f^\lambda(x)]^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) f^{\varpi+\xi}(x)] \\ & + \rho \mathcal{J}_{\eta,k}^{\theta,\pi} [w(x) f^{\xi+\varpi}(x)]^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) h^\xi(x) f^\lambda(x)]. \end{aligned} \tag{3.19}$$

Hence result is proved.  $\square$

Next, we shall propose a new generalization of weighted fractional integral inequalities using a family of  $n$  positive functions defined on  $[0, \infty)$ .

**THEOREM 3.5.** *Let  $f_r$ ,  $i = 1, \dots, n$  be  $n$  positive and continuous functions on  $[0, \infty)$  such that*

$$(\sigma^\xi f_r^\xi(\tau) - \tau^\xi f_r^\xi(\sigma))(f_r^{\varpi-\lambda_r}(\tau) - f_r^{\varpi-\lambda_r}(\sigma)) \geq 0. \tag{3.20}$$

Let  $w : [0, \infty) \rightarrow \mathbb{R}^+$ . Then for all  $\alpha \geq 0$ ,  $x > 0$ ,  $\beta, \rho, \eta, k \in \mathbb{R}$ ,  $\xi > 0$ ,  $\varpi \geq \lambda_r > 0$ ,  $r \in \{1, \dots, n\}$ , the following fractional inequality

$$\begin{aligned} & \rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) f_r^\xi(x) \prod_{i=1}^n f_i^{\lambda_i}(x)]^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) x^\xi f_r^\varpi(x) \prod_{i \neq r}^n f_i^{\lambda_i}(x)] \\ & \leq \rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) x^\xi \prod_{i=1}^n f_i^{\lambda_i}(x)]^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) f_r^{\varpi+\xi}(x) \prod_{i \neq r}^n f_i^{\lambda_i}(x)], \end{aligned} \tag{3.21}$$

is valid.

*Proof.* Suppose  $f_i$ ,  $i = 1, \dots, n$  be  $n$  positive and continuous functions on  $[0, \infty)$ , then for any fixed  $r \in \{1, \dots, n\}$  and for any  $\xi > 0$ ,  $\varpi \geq \lambda_r > 0$ ,  $\tau, \sigma \in (0, x)$ ,  $x > 0$ , from (3.20), we obtain

$$\begin{aligned} & \sigma^\xi f_r^{\varpi-\lambda_r}(\sigma) f_r^\xi(\tau) + \tau^\xi f_r^\xi(\sigma) f_r^{\varpi-\lambda_r}(\tau) \\ & \leq \sigma^\xi f_r^{\varpi+\xi-\lambda_r}(\tau) + \tau^\xi f_r^{\varpi+\xi-\lambda_r}(\sigma). \end{aligned} \tag{3.22}$$

Now multiplying both sides of (3.22) by  $\frac{\rho^{1-\beta}x^k}{\Gamma(\alpha)} \frac{\tau^{\rho(\eta+1)-1}}{(x^\rho - \tau^\rho)^{1-\alpha}} w(\tau) \prod_{i=1}^n f_i^{\lambda_i}(\tau)$ , then integrating the resulting inequality with respect to  $\tau$  over  $(0, x)$ , we obtain

$$\begin{aligned} & \sigma^\xi f_r^{\overline{\omega}-\lambda_r}(\sigma) \frac{\rho^{1-\beta}x^k}{\Gamma(\alpha)} \int_0^x \frac{\tau^{\rho(\eta+1)-1}}{(x^\rho - \tau^\rho)^{1-\alpha}} [w(\tau) f_r^\xi(\tau) \prod_{i=1}^n f_i^{\lambda_i}(\tau)] d\tau \\ & + f_r^\xi(\sigma) \frac{\rho^{1-\beta}x^k}{\Gamma(\alpha)} \int_0^x \frac{\tau^{\rho(\eta+1)-1}}{(x^\rho - \tau^\rho)^{1-\alpha}} [w(\tau) \tau^\xi \prod_{i \neq r}^n f_i^{\lambda_i}(\tau)] d\tau \\ & \leq \sigma^\xi \frac{\rho^{1-\beta}x^k}{\Gamma(\alpha)} \int_0^x \frac{\tau^{\rho(\eta+1)-1}}{(x^\rho - \tau^\rho)^{1-\alpha}} [w(\tau) f_r^{\overline{\omega}+\xi}(\tau) \prod_{i \neq r}^n f_i^{\lambda_i}(\tau)] d\tau \\ & + f_r^{\overline{\omega}+\xi-\lambda_r}(\sigma) \frac{\rho^{1-\beta}x^k}{\Gamma(\alpha)} \int_0^x \frac{\tau^{\rho(\eta+1)-1}}{(x^\rho - \tau^\rho)^{1-\alpha}} [w(\tau) \tau^\xi \prod_{i=1}^n f_i^{\lambda_i}(\tau)] d\tau, \end{aligned} \tag{3.23}$$

consequently

$$\begin{aligned} & \sigma^\xi f_r^{\overline{\omega}-\lambda_r}(\sigma)^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) f_r^\xi(x) \prod_{i=1}^n f_i^{\lambda_i}(x)] \\ & + f_r^\xi(\sigma)^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) x^\xi \prod_{i \neq r}^n f_i^{\lambda_i}(x)] \\ & \leq \sigma^\xi \rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) f_r^{\overline{\omega}+\xi}(x) \prod_{i \neq r}^n f_i^{\lambda_i}(x)] \\ & + f_r^{\overline{\omega}+\xi-\lambda_r}(\sigma)^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) x^\xi \prod_{i=1}^n f_i^{\lambda_i}(x)]. \end{aligned} \tag{3.24}$$

Again, multiplying the inequality (3.24) by  $\frac{\rho^{1-\beta}x^k}{\Gamma(\alpha)} \frac{\sigma^{\rho(\eta+1)-1}}{(x^\rho - \sigma^\rho)^{1-\alpha}} w(\sigma) \prod_{i=1}^n f_i^{\lambda_i}(\sigma)$ ,  $\sigma \in (0, x)$ ,  $x > 0$ , this function remains positive under the conditions stated with the theorem. Integrating the obtained result with respect to  $\sigma$  from 0 to  $x$ , we get

$$\begin{aligned} & \rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) f_r^\xi(x) \prod_{i=1}^n f_i^{\lambda_i}(x)] \frac{\rho^{1-\beta}x^k}{\Gamma(\alpha)} \int_0^x \frac{\sigma^{\rho(\eta+1)-1}}{(x^\rho - \sigma^\rho)^{1-\alpha}} w(\sigma) \sigma^\xi f_r^{\overline{\omega}}(\sigma) \prod_{i \neq r}^n f_i^{\lambda_i}(\sigma) d\sigma \\ & + \rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) x^\xi \prod_{i \neq r}^n f_i^{\lambda_i}(x)] \frac{\rho^{1-\beta}x^k}{\Gamma(\alpha)} \int_0^x \frac{\sigma^{\rho(\eta+1)-1}}{(x^\rho - \sigma^\rho)^{1-\alpha}} w(\sigma) f_r^\xi(\sigma) \prod_{i=1}^n f_i^{\lambda_i}(\sigma) d\sigma \\ & \leq \rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) f_r^{\overline{\omega}+\xi}(x) \prod_{i \neq r}^n f_i^{\lambda_i}(x)] \frac{\rho^{1-\beta}x^k}{\Gamma(\alpha)} \int_0^x \frac{\sigma^{\rho(\eta+1)-1}}{(x^\rho - \sigma^\rho)^{1-\alpha}} w(\sigma) \sigma^\xi \prod_{i=1}^n f_i^{\lambda_i}(\sigma) d\sigma \\ & + \rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) x^\xi \prod_{i=1}^n f_i^{\lambda_i}(x)] \frac{\rho^{1-\beta}x^k}{\Gamma(\alpha)} \int_0^x \frac{\sigma^{\rho(\eta+1)-1}}{(x^\rho - \sigma^\rho)^{1-\alpha}} w(\sigma) f_r^{\overline{\omega}+\xi}(\sigma) \prod_{i \neq r}^n f_i^{\lambda_i}(\sigma) d\sigma, \end{aligned} \tag{3.25}$$

therefore,

$$\begin{aligned} & \rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) f_r^\xi(x) \prod_{i=1}^n f_i^{\lambda_i}(x)]^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [x^\xi f_r^{\overline{\omega}}(x) \prod_{i \neq r}^n f_i^{\lambda_i}(x)] \\ & + \rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) x^\xi \prod_{i \neq r}^n f_i^{\lambda_i}(x)]^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) f_r^\xi(x) \prod_{i=1}^n f_i^{\lambda_i}(x)] \\ & \leq \rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) f_r^{\overline{\omega}+\xi}(x) \prod_{i \neq r}^n f_i^{\lambda_i}(x)]^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) x^\xi \prod_{i=1}^n f_i^{\lambda_i}(x)] \\ & + \rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) x^\xi \prod_{i=1}^n f_i^{\lambda_i}(x)]^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) f_r^{\overline{\omega}+\xi}(x) \prod_{i \neq r}^n f_i^{\lambda_i}(x)]. \end{aligned} \tag{3.26}$$



This ends the inequality (3.21).  $\square$

**THEOREM 3.6.** *Let  $f_i, i = 1, \dots, n$  be  $n$  positive and continuous functions on  $[0, \infty)$  and satisfying (3.20). Let  $w : [0, \infty) \rightarrow \mathbb{R}^+$  be positive continuous function. Then for all  $\alpha, \theta \geq 0, x > 0, \beta, \pi, \rho, \eta, k \in \mathbb{R}, \xi > 0, \varpi \geq \lambda_r > 0, r \in \{1, \dots, n\}$ , we have that the inequality*

$$\begin{aligned} & \rho \mathcal{J}_{\eta,k}^{\theta,\pi} [w(x)x^\xi f_r^\varpi(x) \prod_{i \neq r} f_i^{\lambda_i}(x)]^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x)f_r^\xi(x) \prod_{i=1}^n f_i^{\lambda_i}(x)] \\ & \quad + \rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x)x^\xi f_r^\varpi(x) \prod_{i \neq r} f_i^{\lambda_i}(x)]^\rho \mathcal{J}_{\eta,k}^{\theta,\pi} [w(x)f_r^\xi(x) \prod_{i=1}^n f_i^{\lambda_i}(x)] \\ & \leq \rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x)f_r^{\varpi+\xi}(x) \prod_{i \neq r} f_i^{\lambda_i}(x)]^\rho \mathcal{J}_{\eta,k}^{\theta,\pi} [w(x)x^\xi \prod_{i=1}^n f_i^{\lambda_i}(x)] \\ & \quad + \rho \mathcal{J}_{\eta,k}^{\theta,\pi} [w(x)f_r^{\varpi+\xi}(x) \prod_{i \neq r} f_i^{\lambda_i}(x)]^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x)x^\xi \prod_{i=1}^n f_i^{\lambda_i}(x)], \end{aligned} \tag{3.27}$$

is valid.

*Proof.* We multiplying the inequality (3.24) by  $\frac{\rho^{1-\pi} x^k}{\Gamma(\theta)} \frac{\sigma^{\rho(\eta+1)-1}}{(x^\rho - \sigma^\rho)^{1-\theta}} w(\sigma) \prod_{i=1}^n f_i^{\lambda_i}(\sigma)$ ,  $\sigma \in (0, x), x > 0$ , this function remains positive under the conditions stated with the theorem. Integrating the obtained result with respect to  $\sigma$  from 0 to  $x$ , we get

$$\begin{aligned} & \rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x)f_r^\xi(x) \prod_{i=1}^n f_i^{\lambda_i}(x)]^\rho \frac{\rho^{1-\pi} x^k}{\Gamma(\theta)} \int_0^x \frac{\sigma^{\rho(\eta+1)-1}}{(x^\rho - \sigma^\rho)^{1-\theta}} w(\sigma) \sigma^\xi f_r^\varpi(\sigma) \prod_{i \neq r} f_i^{\lambda_i}(\sigma) d\sigma \\ & \quad + \rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x)x^\xi \prod_{i \neq r} f_i^{\lambda_i}(x)]^\rho \frac{\rho^{1-\pi} x^k}{\Gamma(\theta)} \int_0^x \frac{\sigma^{\rho(\eta+1)-1}}{(x^\rho - \sigma^\rho)^{1-\theta}} w(\sigma) f_r^\xi(\sigma) \prod_{i=1}^n f_i^{\lambda_i}(\sigma) d\sigma \\ & \leq \rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x)f_r^{\varpi+\xi}(x) \prod_{i \neq r} f_i^{\lambda_i}(x)]^\rho \frac{\rho^{1-\pi} x^k}{\Gamma(\theta)} \int_0^x \frac{\sigma^{\rho(\eta+1)-1}}{(x^\rho - \sigma^\rho)^{1-\theta}} w(\sigma) \sigma^\xi \prod_{i=1}^n f_i^{\lambda_i}(\sigma) d\sigma \\ & \quad + \rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x)x^\xi \prod_{i=1}^n f_i^{\lambda_i}(x)]^\rho \frac{\rho^{1-\pi} x^k}{\Gamma(\theta)} \int_0^x \frac{\sigma^{\rho(\eta+1)-1}}{(x^\rho - \sigma^\rho)^{1-\theta}} w(\sigma) f_r^{\varpi+\xi}(\sigma) \prod_{i \neq r} f_i^{\lambda_i}(\sigma) d\sigma, \end{aligned} \tag{3.28}$$

which gives the inequality (3.27).  $\square$

**THEOREM 3.7.** *Let  $h, f_i, i = 1, \dots, n$  be positive and continuous functions on  $[0, \infty)$ , such that*

$$(h^\xi(\sigma) f_r^\xi(\tau) - h^\xi(\tau) f_r^\xi(\sigma)) (f_r^{\varpi-\lambda_r}(\tau) - f_r^{\varpi-\lambda_r}(\sigma)) \geq 0. \tag{3.29}$$

*Let  $w : [0, \infty) \rightarrow \mathbb{R}^+$  be positive continuous function. Then for all  $\alpha \geq 0, x > 0, \beta, \rho, \eta, k \in \mathbb{R}, \xi > 0, \varpi \geq \lambda_r > 0, r \in \{1, \dots, n\}$ , the following fractional inequality*

$$\begin{aligned} & \rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x)f_r^\xi(x) \prod_{i=1}^n f_i^{\lambda_i}(x)]^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x)h^\xi(x) f_r^\varpi(x) \prod_{i \neq r} f_i^{\lambda_i}(x)] \\ & \leq \rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x)h^\xi(x) \prod_{i=1}^n f_i^{\lambda_i}(x)]^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x)f_r^{\varpi+\xi}(x) \prod_{i \neq r} f_i^{\lambda_i}(x)], \end{aligned} \tag{3.30}$$

is valid.

*Proof.* Let  $\tau, \sigma \in (0, x)$ ,  $x > 0$ , for any  $\xi > 0$ ,  $\varpi \geq \lambda_i > 0$ ,  $r \in \{1, 2, \dots, n\}$  then from (3.29), we have

$$\begin{aligned} & h^\xi(\sigma) f_r^{\varpi-\lambda_r}(\sigma) f_r^\xi(\tau) + f_r^\xi(\sigma) h_r^\xi(\tau) f_r^{\varpi-\lambda_r}(\tau) \\ & \leq h^\xi(\sigma) f_r^{\varpi+\xi-\lambda_r}(\tau) + f_r^{\varpi+\xi-\lambda_r}(\sigma) h^\xi(\tau). \end{aligned} \tag{3.31}$$

Multiplying both sides of (3.31) by  $\frac{\rho^{1-\beta} x^k}{\Gamma(\alpha)} \frac{\tau^{\rho(\eta+1)-1}}{(x^\rho-\tau^\rho)^{1-\alpha}} w(\tau) \prod_{i=1}^n f_i^{\lambda_i}(\tau)$ , then integrating the resulting inequality with respect to  $\tau$  over  $(0, x)$ , we obtain

$$\begin{aligned} & h^\xi(\sigma) f_r^{\varpi-\lambda_r}(\sigma)^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) f_r^\xi(x) \prod_{i=1}^n f_i^{\lambda_i}(x)] \\ & + f_r^\xi(\sigma)^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) h^\xi(x) f_r^\varpi(x) \prod_{i \neq r}^n f_i^{\lambda_i}(x)] \\ & \leq h^\xi(\sigma)^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) h^\xi(x) \prod_{i=1}^n f_i^{\lambda_i}(x)] \\ & + f_r^{\varpi+\xi-\lambda_r}(\sigma)^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) f_r^{\varpi+\xi}(x) \prod_{i \neq r}^n f_i^{\lambda_i}(x)]. \end{aligned} \tag{3.32}$$

Multiplying both sides of (3.32) by  $\frac{\rho^{1-\beta} x^k}{\Gamma(\alpha)} \frac{\sigma^{\rho(\eta+1)-1}}{(x^\rho-\sigma^\rho)^{1-\alpha}} w(\sigma) \prod_{i=1}^n f_i^{\lambda_i}(\sigma)$ , then integrating the resulting inequality with respect to  $\sigma$  over  $(0, x)$ , we have

$$\begin{aligned} & 2^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) f_r^\xi(x) \prod_{i=1}^n f_i^{\lambda_i}(x)]^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) h^\xi(x) f_r^\varpi(x) \prod_{i \neq r}^n f_i^{\lambda_i}(x)] \\ & \leq 2^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) f_r^{\varpi+\xi}(x) \prod_{i \neq r}^n f_i^{\lambda_i}(x)]^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) h^\xi(x) \prod_{i=1}^n f_i^{\lambda_i}(x)]. \end{aligned} \tag{3.33}$$

This completes the proof of Theorem 3.7.  $\square$

**THEOREM 3.8.** *Let  $h, f_i, i = 1, \dots, n$  be positive and continuous functions on  $[0, \infty)$ , such that*

$$(h^\xi(\sigma) f_r^\xi(\tau) - h^\xi(\tau) f_r^\xi(\sigma)) (f_r^{\varpi-\lambda_r}(\tau) - f_r^{\varpi-\lambda_r}(\sigma)) \geq 0. \tag{3.34}$$

*Let  $w : [0, \infty) \rightarrow \mathbb{R}^+$  be positive continuous function. Then for all  $\alpha, \theta \geq 0$ ,  $x > 0$ ,  $\beta, \pi, \rho, \eta, k \in \mathbb{R}$ ,  $\xi > 0$ ,  $\varpi \geq \lambda_r > 0$ ,  $r \in \{1, \dots, n\}$ , the following fractional inequality*

$$\begin{aligned} & {}^\rho \mathcal{J}_{\eta,k}^{\theta,\pi} [w(x) h^\xi(x) f_r^\varpi(x) \prod_{i \neq r}^n f_i^{\lambda_i}(x)] {}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) f_r^\xi(x) \prod_{i=1}^n f_i^{\lambda_i}(x)] \\ & + {}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) h^\xi(x) f_r^\varpi(x) \prod_{i \neq r}^n f_i^{\lambda_i}(x)] {}^\rho \mathcal{J}_{\eta,k}^{\theta,\pi} [w(x) f_r^\xi(x) \prod_{i=1}^n f_i^{\lambda_i}(x)] \\ & \leq {}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) f_r^{\varpi+\xi}(x) \prod_{i \neq r}^n f_i^{\lambda_i}(x)] {}^\rho \mathcal{J}_{\eta,k}^{\theta,\pi} [w(x) h^\xi(x) \prod_{i=1}^n f_i^{\lambda_i}(x)] \\ & {}^\rho \mathcal{J}_{\eta,k}^{\theta,\pi} [w(x) f_r^{\varpi+\xi}(x) \prod_{i \neq r}^n f_i^{\lambda_i}(x)] {}^\rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x) h^\xi(x) \prod_{i=1}^n f_i^{\lambda_i}(x)], \end{aligned} \tag{3.35}$$

*is valid.*

*Proof.* Multiplying both sides of (3.31) by  $\frac{\rho^{1-\pi} x^k}{\Gamma(\theta)} \frac{\sigma^{\rho(\eta+1)-1}}{(x^\rho-\sigma^\rho)^{1-\theta}} w(\sigma) \prod_{i=1}^n f_i^{\lambda_i}(\sigma)$ , then integrating the resulting inequality with respect to  $\sigma$  over  $(0, x)$ , we have

$$\begin{aligned} & f_r^\xi(\tau)^\rho \mathcal{J}_{\eta,k}^{\theta,\pi} [w(x) h^\xi(x) f_r^\varpi(x) \prod_{i \neq r}^n f_i^{\lambda_i}(x)] + h^\xi(\tau) f_r^{\varpi-\lambda_r}(\tau)^\rho \mathcal{J}_{\eta,k}^{\theta,\pi} [w(x) f_r^\xi(x) \prod_{i=1}^n f_i^{\lambda_i}(x)] \\ & \leq f_r^{\varpi+\xi-\lambda_r}(\tau)^\rho \mathcal{J}_{\eta,k}^{\theta,\pi} [w(x) h^\xi(x) \prod_{i=1}^n f_i^{\lambda_i}(x)] + h^\xi(\tau)^\rho \mathcal{J}_{\eta,k}^{\theta,\pi} [w(x) f_r^{\varpi+\xi}(x) \prod_{i \neq r}^n f_i^{\lambda_i}(x)]. \end{aligned} \tag{3.36}$$

Multiplying both sides of (3.36) by  $\frac{\rho^{1-\beta}x^k}{\Gamma(\alpha)} \frac{\tau^{\rho(\eta+1)-1}}{(x^\rho-\tau^\rho)^{1-\alpha}} w(\tau) \prod_{i=1}^n f_i^{\lambda_i}(\tau)$ , then integrating the resulting inequality with respect to  $\tau$  over  $(0, x)$ , we have

$$\begin{aligned} & \rho \mathcal{J}_{\eta,k}^{\theta,\pi} [w(x)h^\xi(x)f_r^\omega(x)\prod_{i\neq r}^n f_i^{\lambda_i}(x)] \rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x)f_r^\xi(x)\prod_{i=1}^n f_i^{\lambda_i}(x)] \\ & \quad + \rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x)h^\xi(x)f_r^\omega(x)\prod_{i\neq r}^n f_i^{\lambda_i}(x)] \rho \mathcal{J}_{\eta,k}^{\theta,\pi} [w(x)f_r^\xi(x)\prod_{i=1}^n f_i^{\lambda_i}(x)] \\ & \leq \rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x)f_r^{\omega+\xi}(x)\prod_{i\neq r}^n f_i^{\lambda_i}(x)] \rho \mathcal{J}_{\eta,k}^{\theta,\pi} [w(x)h^\xi(x)\prod_{i=1}^n f_i^{\lambda_i}(x)] \\ & \quad \rho \mathcal{J}_{\eta,k}^{\theta,\pi} [w(x)f_r^{\omega+\xi}(x)\prod_{i\neq r}^n f_i^{\lambda_i}(x)] \rho \mathcal{J}_{\eta,k}^{\alpha,\beta} [w(x)h^\xi(x)\prod_{i=1}^n f_i^{\lambda_i}(x)]. \end{aligned} \quad (3.37)$$

This completes the proof.  $\square$

#### 4. Concluding remarks

Several fractional integral inequalities have been investigated by employing the different fractional integral operators. In this paper, we established weighted fractional integral inequalities using generalized Katugampola fractional integral operators. This works give some contribution to the theory of fractional integral inequalities and fractional calculus.

#### REFERENCES

- [1] T. A. ALJAAIDI AND D. B. PACHPATTE, *Osone Grüss-type inequalities using generalized Katugampola fractional integral*, AIMS MATH., 5 (2), (2020), 1011–1024. doi: 10.3934/math.2020070.
- [2] G. A. ANASTASSIOU, *Fractional Differentiation Inequalities*, Springer Publishing Company, New York, NY, 2009.
- [3] D. BALEANU, S. D. PUROHIT AND P. AGARWAL, *On fractional integral inequalities involving hypergeometric operators*, Chin. J. Math., Article ID609476, 2014,(2014), 1–5. <https://doi.org/10.1155/2014/609476>.
- [4] V. L. CHINCHANE AND D. B. PACHPATTE, *On some integral inequalities using Hadamard fractional integral*, Malaya J. Mat., 1 (1), (2012), 62–66.
- [5] V. L. CHINCHANE, *On Chebyshev type inequalities using generalized k-fractional integral operator*, Progr. Fract. Differ. Appl., 3 (3), (2017), 219–226.
- [6] V. L. CHINCHANE AND D. B. PACHPATTE, *On some Grüss-type fractional inequalities using Saigo fractional integral operator*, Journal of Mathematics, Article ID 527910, 2014 (2014), 1–9. <https://doi.org/10.1155/2014/527910>.
- [7] L. CURIEL AND L. GALUE, *A generalization of the integral operators involving the Gruss hypergeometric function*, Rev. Tech. Ingr. Unlv. Zulla, 1 (19), (1996), 17–22.
- [8] Z. DAHMANI, *The Riemann-Liouville operator to generate some new inequalities*, Int. J. Nonlinear Sci., 12 (4) (2011), 452–455.
- [9] Z. DAHMANI, *Some results associate with fractional integrals involving the extended Chebyshev*, Acta Univ. Apulensis Math. Inform., 27 (2011), 217–224.
- [10] M. HOUAS, *Certain weighted integral inequalities involving the fractional hypergeometric operators*, Scientia Series A: Mathematical Science, 27(2016), 87–97.
- [11] A. A. KILBAS, H. M. SRIVASATAVA AND J. J. TRUJILLO, *Theory and Application of Fractional Differential Equation*, North-Holland Mathematiccal Studies Vol. Elsevier Science Publishers, Amsterdam, London and New York, 2006.
- [12] U. N. KATUGAMPOLA, *A new approach to generalized fractional derivatives*, Bull. Math. Anal. Appl., 6 (4), (2014), 1–15.

- [13] U. N. KATUGAMPOLA, *new fractional integral unifying six existing fractional integral*, arXiv:1612.08596 [math.CA] (2016), 1–6.
- [14] V. KIRYAKOVA, *On two Saigo's fractional integral operator in the class of univalent functions*, Fract. Calc. Appl. Anal., 9 (2), (2006).
- [15] V. S. KIRYAKOVA, *Generalized Fractional Calculus and Applications*, Pitman Res. Notes in Math. Ser., 301, Longman, Harlow, 1994. <https://doi.org/10.1017/S0013091500006325>.
- [16] A. B. NALE, S. K. PANCHAL AND V. L. CHINCHANE, *Certain fractional integral inequalities using generalized Katugampola fractional integral operator*, Malaya J. Mat., 3 (8), (2020), 809–814.
- [17] A. R. PRABHAKARAN AND K. SRINIVASA RAO, *Saigo operator of fractional integration of Hypergeometric functions*, Int. J. Pure Appl. Math., 81 (5), (2012), 755–763.
- [18] M. SAIGO, *A remark on integral operators involving the Gauss hypergeometric functions*, Math. Rep. Kyushu Univ, 11, (1978), 135.
- [19] E. SET, J. CHOI AND L. MUMCU, *Chebyshev type inequalities involving generalized Katugampola fractional integral operators*, Tamkang J. Mathe., 4 (50), (2019), 381–390.
- [20] H. M. SRIVASTAVA AND J. CHOI, *Zeta and q-Zeta function and associated series and integral*, Elsevier Science Publishers, Amsterdam, London, New York, 2012.
- [21] J. TARIBOON, S. K. NTOUYAS AND W. SUDSUTAD, *Some New Riemann-Liouville Fractional Integral Inequalities*, Int. J. Math. Math. Sci., Article ID 869434, 2014 (2014), 1–6.
- [22] J. VANTERLER DA C. SOUSA D. S. OLIVEIRA E. CAPELAS DE OLIVEIRA, *Grüss- type inequalities by means of generalized fractional integrals*, Bull. Braz. Math. Soc., New 50, (2019), 1029–1047.

(Received July 15, 2020)

Satish K. Panchal  
Department of Mathematics  
Dr. Babasaheb Ambedkar Marathwada University  
Aurangabad-431 004, India  
e-mail: drskpanchal@gmail.com

Vaijanath L. Chinchane  
Department of Mathematics  
Deogiri Institute of Engineering and Management Studies  
Aurangabad-431005, India  
e-mail: chinchane85@gmail.com

Asha B. Nale  
Department of Mathematics  
Dr. Babasaheb Ambedkar Marathwada University  
Aurangabad-431 004, India  
e-mail: ashabnale@gmail.com