

OSTROWSKI AND TRAPEZOID TYPE INEQUALITIES FOR RIEMANN-LIOUVILLE FRACTIONAL INTEGRALS OF ABSOLUTELY CONTINUOUS FUNCTIONS WITH BOUNDED DERIVATIVES

SILVESTRU SEVER DRAGOMIR

(Communicated by B. Ahmad)

Abstract. In this paper we establish some Ostrowski and trapezoid type inequalities for the Riemann-Liouville fractional integrals of absolutely continuous functions with bounded derivatives. Applications for mid-point and trapezoid inequalities are provided as well. They generalize the known results holding for the classical Riemann integral. Some examples for convex functions are also given.

1. Introduction

In 2002 [12], we proved the following *Ostrowski type inequality* for convex functions $f : [a, b] \rightarrow \mathbb{R}$ and $x \in (a, b)$,

$$\begin{aligned} & \frac{1}{2} \left[(b-x)^2 f'_+(x) - (x-a)^2 f'_-(x) \right] \\ & \leq \int_a^b f(t) dt - (b-a)f(x) \\ & \leq \frac{1}{2} \left[(b-x)^2 f'_-(b) - (x-a)^2 f'_+(a) \right]. \end{aligned} \tag{1.1}$$

In particular, we have the *mid-point inequalities*

$$\begin{aligned} & \frac{1}{8} \left[f'_+ \left(\frac{a+b}{2} \right) - f'_- \left(\frac{a+b}{2} \right) \right] (b-a)^2 \\ & \leq \int_a^b f(t) dt - (b-a)f \left(\frac{a+b}{2} \right) \\ & \leq \frac{1}{8} [f'_-(b) - f'_+(a)] (b-a)^2, \end{aligned} \tag{1.2}$$

with the constant $\frac{1}{8}$ as best possible in both inequalities.

Mathematics subject classification (2010): 26D15, 26D10, 26D07, 26A33.

Keywords and phrases: Riemann-Liouville fractional integrals, absolutely continuous functions, Ostrowski type inequalities, trapezoid inequalities, bounded derivatives.

In the same year [13], we also obtained the following *generalized trapezoid type inequality* for convex functions $f : [a, b] \rightarrow \mathbb{R}$ and $x \in (a, b)$,

$$\begin{aligned} & \frac{1}{2} \left[(b-x)^2 f'_+(x) - (x-a)^2 f'_-(x) \right] \\ & \leq (x-a)f(a) + (b-x)f(b) - \int_a^b f(t) dt \\ & \leq \frac{1}{2} \left[(b-x)^2 f'_-(b) - (x-a)^2 f'_+(a) \right]. \end{aligned} \quad (1.3)$$

In particular, we have the *trapezoid inequality*

$$\begin{aligned} & \frac{1}{8} \left[f'_+ \left(\frac{a+b}{2} \right) - f'_- \left(\frac{a+b}{2} \right) \right] (b-a)^2 \\ & \leq \frac{f(a) + f(b)}{2} (b-a) - \int_a^b f(t) dt \\ & \leq \frac{1}{8} [f'_-(b) - f'_+(a)] (b-a)^2, \end{aligned} \quad (1.4)$$

with the constant $\frac{1}{8}$ as best possible in both inequalities.

These results were generalized in the following manner:

THEOREM 1. (Dragomir, 2003 [14]) *Let $f : [a, b] \rightarrow \mathbb{R}$ be an absolutely continuous function on $[a, b]$ and $x \in [a, b]$. Suppose that there exist the functions $m_i, M_i : [a, b] \rightarrow \mathbb{R}$ ($i = 1, 2$) with the properties:*

$$m_1(x) \leq f'(t) \leq M_1(x) \text{ for a.e. } t \in [a, x] \quad (1.5)$$

and

$$m_2(x) \leq f'(t) \leq M_2(x) \text{ for a.e. } t \in (x, b]. \quad (1.6)$$

Then we have the inequalities:

$$\begin{aligned} & \frac{1}{2(b-a)} \left[m_1(x)(x-a)^2 - M_2(x)(b-x)^2 \right] \\ & \leq f(x) - \frac{1}{b-a} \int_a^b f(t) dt \\ & \leq \frac{1}{2(b-a)} \left[M_1(x)(x-a)^2 - m_2(x)(b-x)^2 \right]. \end{aligned} \quad (1.7)$$

The constant $\frac{1}{2}$ is sharp on both sides.

If we assume global bounds for the derivative, then we have:

COROLLARY 1. (Dragomir, 2003 [14]) *If $f : [a, b] \rightarrow \mathbb{R}$ is absolutely continuous on $[a, b]$ and the derivative $f' : [a, b] \rightarrow \mathbb{R}$ is bounded above and below, that is,*

$$-\infty < m \leq f'(t) \leq M < \infty \text{ for a.e. } t \in [a, b], \quad (1.8)$$

then we have the inequality

$$\begin{aligned} \frac{1}{2(b-a)} \left[m(x-a)^2 - M(b-x)^2 \right] &\leq f(x) - \frac{1}{b-a} \int_a^b f(t) dt \\ &\leq \frac{1}{2(b-a)} \left[M(x-a)^2 - m(b-x)^2 \right] \end{aligned} \quad (1.9)$$

for all $x \in [a, b]$. The constant $\frac{1}{2}$ is the best in both inequalities.

In particular, we have

$$\left| \frac{1}{b-a} \int_a^b f(t) dt - f\left(\frac{a+b}{2}\right) \right| \leq \frac{1}{8} (M-m)(b-a), \quad (1.10)$$

with $\frac{1}{8}$ as the best possible constant.

In order to extend these results for fractional integrals we need the following definitions.

Let $f : [a, b] \rightarrow \mathbb{C}$ be a complex valued Lebesgue integrable function on the real interval $[a, b]$. The Riemann-Liouville fractional integrals are defined for $\alpha > 0$ by

$$J_{a+}^{\alpha} f(x) = \frac{1}{\Gamma(\alpha)} \int_a^x (x-t)^{\alpha-1} f(t) dt$$

for $a < x \leq b$ and

$$J_{b-}^{\alpha} f(x) = \frac{1}{\Gamma(\alpha)} \int_x^b (t-x)^{\alpha-1} f(t) dt$$

for $a \leq x < b$, where Γ is the Gamma function. For $\alpha = 0$, they are defined as

$$J_{a+}^0 f(x) = J_{b-}^0 f(x) = f(x) \text{ for } x \in (a, b).$$

For several Ostrowski type inequalities for Riemann-Liouville fractional integrals see [1]–[6], [19]–[29] and the references therein.

Motivated by the above results, we obtain in this paper some inequalities for the Riemann-Liouville fractional integrals of absolutely continuous functions with bounded derivatives and of convex functions. Applications for mid-point and trapezoid inequalities are provided as well. Some examples for convex functions are also given.

2. Some identities

We have the following representation:

LEMMA 1. Let $f : [a, b] \rightarrow \mathbb{C}$ be an absolutely continuous function on $[a, b]$.

(i) For any $x \in (a, b)$ we have

$$\begin{aligned} &J_{a+}^{\alpha} f(x) + J_{b-}^{\alpha} f(x) \\ &= \frac{1}{\Gamma(\alpha+1)} \left[(x-a)^{\alpha} f(a) + (b-x)^{\alpha} f(b) \right] \\ &\quad + \frac{1}{\Gamma(\alpha+1)} \left[\int_a^x (x-t)^{\alpha} f'(t) dt - \int_x^b (t-x)^{\alpha} f'(t) dt \right]. \end{aligned} \quad (2.1)$$

(ii) For any $x \in (a, b)$ we have

$$\begin{aligned} J_{x-}^{\alpha} f(a) + J_{x+}^{\alpha} f(b) & \\ = \frac{1}{\Gamma(\alpha+1)} [(x-a)^{\alpha} + (b-x)^{\alpha}] f(x) & \\ + \frac{1}{\Gamma(\alpha+1)} \left[\int_x^b (b-t)^{\alpha} f'(t) dt - \int_a^x (t-a)^{\alpha} f'(t) dt \right]. & \end{aligned} \quad (2.2)$$

Proof. (i) Since $f : [a, b] \rightarrow \mathbb{C}$ is an absolutely continuous function on $[a, b]$, then the Lebesgue integrals

$$\int_a^x (x-t)^{\alpha} f'(t) dt \text{ and } \int_x^b (t-x)^{\alpha} f'(t) dt$$

exist and integrating by parts, we have

$$\begin{aligned} \frac{1}{\Gamma(\alpha+1)} \int_a^x (x-t)^{\alpha} f'(t) dt & \\ = \frac{1}{\Gamma(\alpha)} \int_a^x (x-t)^{\alpha-1} f(t) dt - \frac{1}{\Gamma(\alpha+1)} (x-a)^{\alpha} f(a) & \\ = J_{a+}^{\alpha} f(x) - \frac{1}{\Gamma(\alpha+1)} (x-a)^{\alpha} f(a) & \end{aligned} \quad (2.3)$$

for $a < x \leq b$ and

$$\begin{aligned} \frac{1}{\Gamma(\alpha+1)} \int_x^b (t-x)^{\alpha} f'(t) dt & \\ = \frac{1}{\Gamma(\alpha+1)} (b-x)^{\alpha} f(b) - \frac{1}{\Gamma(\alpha)} \int_x^b (t-x)^{\alpha-1} f(t) dt & \\ = \frac{1}{\Gamma(\alpha+1)} (b-x)^{\alpha} f(b) - J_{b-}^{\alpha} f(x) & \end{aligned} \quad (2.4)$$

for $a \leq x < b$.

From (2.3) we have

$$J_{a+}^{\alpha} f(x) = \frac{1}{\Gamma(\alpha+1)} (x-a)^{\alpha} f(a) + \frac{1}{\Gamma(\alpha+1)} \int_a^x (x-t)^{\alpha} f'(t) dt$$

for $a < x \leq b$ and from (2.4) we have

$$J_{b-}^{\alpha} f(x) = \frac{1}{\Gamma(\alpha+1)} (b-x)^{\alpha} f(b) - \frac{1}{\Gamma(\alpha+1)} \int_x^b (t-x)^{\alpha} f'(t) dt,$$

for $a \leq x < b$, which by addition give (2.1).

(ii) We have

$$J_{x+}^{\alpha} f(b) = \frac{1}{\Gamma(\alpha)} \int_x^b (b-t)^{\alpha-1} f(t) dt$$

for $a \leq x < b$ and

$$J_{x-}^{\alpha} f(a) = \frac{1}{\Gamma(\alpha)} \int_a^x (t-a)^{\alpha-1} f(t) dt$$

for $a < x \leq b$.

Since $f : [a, b] \rightarrow \mathbb{C}$ is an absolutely continuous function $[a, b]$, then the Lebesgue integrals

$$\int_a^x (t-a)^{\alpha} f'(t) dt \text{ and } \int_x^b (b-t)^{\alpha} f'(t) dt$$

exist and integrating by parts, we have

$$\begin{aligned} & \frac{1}{\Gamma(\alpha+1)} \int_a^x (t-a)^{\alpha} f'(t) dt & (2.5) \\ &= \frac{1}{\Gamma(\alpha+1)} (x-a)^{\alpha} f(x) - \frac{1}{\Gamma(\alpha)} \int_a^x (t-a)^{\alpha-1} f(t) dt \\ &= \frac{1}{\Gamma(\alpha+1)} (x-a)^{\alpha} f(x) - J_{x-}^{\alpha} f(a) \end{aligned}$$

for $a < x \leq b$ and

$$\begin{aligned} & \frac{1}{\Gamma(\alpha+1)} \int_x^b (b-t)^{\alpha} f'(t) dt & (2.6) \\ &= \frac{1}{\Gamma(\alpha)} \int_x^b (b-t)^{\alpha-1} f(t) dt - \frac{1}{\Gamma(\alpha+1)} (b-x)^{\alpha} f(x) \\ &= J_{x+}^{\alpha} f(b) - \frac{1}{\Gamma(\alpha+1)} (b-x)^{\alpha} f(x) \end{aligned}$$

for $a \leq x < b$.

From (2.5) we have

$$J_{x-}^{\alpha} f(a) = \frac{1}{\Gamma(\alpha+1)} (x-a)^{\alpha} f(x) - \frac{1}{\Gamma(\alpha+1)} \int_a^x (t-a)^{\alpha} f'(t) dt \tag{2.7}$$

for $a < x \leq b$ and from (2.6)

$$J_{x+}^{\alpha} f(b) = \frac{1}{\Gamma(\alpha+1)} (b-x)^{\alpha} f(x) + \frac{1}{\Gamma(\alpha+1)} \int_x^b (b-t)^{\alpha} f'(t) dt, \tag{2.8}$$

for $a \leq x < b$, which by addition produce (2.2). \square

COROLLARY 2. *Let $f : [a, b] \rightarrow \mathbb{C}$ be an absolutely continuous function on $[a, b]$. We have the midpoint equalities*

$$\begin{aligned} & J_{a+}^{\alpha} f\left(\frac{a+b}{2}\right) + J_{b-}^{\alpha} f\left(\frac{a+b}{2}\right) & (2.9) \\ &= \frac{1}{2^{\alpha-1} \Gamma(\alpha+1)} \frac{f(a) + f(b)}{2} \\ &+ \frac{1}{\Gamma(\alpha+1)} \left[\int_a^{\frac{a+b}{2}} \left(\frac{a+b}{2} - t\right)^{\alpha} f'(t) dt - \int_{\frac{a+b}{2}}^b \left(t - \frac{a+b}{2}\right)^{\alpha} f'(t) dt \right] \end{aligned}$$

and

$$\begin{aligned} & J_{\frac{a+b}{2}-}^{\alpha} f(a) + J_{\frac{a+b}{2}+}^{\alpha} f(b) \\ &= \frac{1}{2^{\alpha-1} \Gamma(\alpha+1)} f\left(\frac{a+b}{2}\right) (b-a)^{\alpha} \\ & \quad + \frac{1}{\Gamma(\alpha+1)} \left[\int_{\frac{a+b}{2}}^b (b-t)^{\alpha} f'(t) dt - \int_a^{\frac{a+b}{2}} (t-a)^{\alpha} f'(t) dt \right] \end{aligned} \quad (2.10)$$

and the trapezoid equality

$$\begin{aligned} \frac{J_{b-}^{\alpha} f(a) + J_{a+}^{\alpha} f(b)}{2} &= \frac{1}{\Gamma(\alpha+1)} \frac{f(b) + f(a)}{2} (b-a)^{\alpha} \\ & \quad + \frac{1}{\Gamma(\alpha+1)} \int_a^b \frac{(b-t)^{\alpha} - (t-a)^{\alpha}}{2} f'(t) dt. \end{aligned} \quad (2.11)$$

Proof. Equality (2.9) follows by (2.1) for $x = \frac{a+b}{2}$ while the equality (2.10) follows by (2.2).

For $x = b$ in (2.7) we have

$$J_{b-}^{\alpha} f(a) = \frac{1}{\Gamma(\alpha+1)} (b-a)^{\alpha} f(b) - \frac{1}{\Gamma(\alpha+1)} \int_a^b (t-a)^{\alpha} f'(t) dt$$

while from (2.8) we have for $x = a$ that

$$J_{a+}^{\alpha} f(b) = \frac{1}{\Gamma(\alpha+1)} (b-a)^{\alpha} f(a) + \frac{1}{\Gamma(\alpha+1)} \int_a^b (b-t)^{\alpha} f'(t) dt.$$

If we add these two equalities and divide by 2, we get (2.11). \square

3. Inequalities for functions with bounded derivatives

We have the following result that provides upper and lower bounds for the Ostrowski and trapezoid differences:

THEOREM 2. *Let $f : [a, b] \rightarrow \mathbb{R}$ be an absolutely continuous function on $[a, b]$. If $x \in (a, b)$ and there exists the real numbers $m_1(x)$, $M_1(x)$, $m_2(x)$, $M_2(x)$ such that*

$$m_1(x) \leq f'(t) \leq M_1(x) \text{ for a.e. } t \in (a, x) \quad (3.1)$$

and

$$m_2(x) \leq f'(t) \leq M_2(x) \text{ for a.e. } t \in (x, b) \quad (3.2)$$

then

$$\begin{aligned} & \frac{1}{\Gamma(\alpha+2)} \left[m_2(x)(b-x)^{\alpha+1} - M_1(x)(x-a)^{\alpha+1} \right] \\ & \leq \frac{1}{\Gamma(\alpha+1)} \left[(x-a)^\alpha f(a) + (b-x)^\alpha f(b) \right] - J_{a+}^\alpha f(x) - J_{b-}^\alpha f(x) \\ & \leq \frac{1}{\Gamma(\alpha+2)} \left[M_2(x)(b-x)^{\alpha+1} - m_1(x)(x-a)^{\alpha+1} \right] \end{aligned} \quad (3.3)$$

and

$$\begin{aligned} & \frac{1}{\Gamma(\alpha+2)} \left[m_2(x)(b-x)^{\alpha+1} - M_1(x)(x-a)^{\alpha+1} \right] \\ & \leq J_{x-}^\alpha f(a) + J_{x+}^\alpha f(b) - \frac{1}{\Gamma(\alpha+1)} \left[(x-a)^\alpha + (b-x)^\alpha \right] f(x) \\ & \leq \frac{1}{\Gamma(\alpha+2)} \left[M_2(x)(b-x)^{\alpha+1} - m_1(x)(x-a)^{\alpha+1} \right]. \end{aligned} \quad (3.4)$$

Proof. We have

$$\begin{aligned} & \frac{1}{\Gamma(\alpha+1)} \left[(x-a)^\alpha f(a) + (b-x)^\alpha f(b) \right] - J_{a+}^\alpha f(x) - J_{b-}^\alpha f(x) \\ & = \frac{1}{\Gamma(\alpha+1)} \left[\int_x^b (t-x)^\alpha f'(t) dt - \int_a^x (x-t)^\alpha f'(t) dt \right] \end{aligned} \quad (3.5)$$

for any $x \in (a, b)$.

Using the conditions (3.1) and (3.2) we have

$$m_2(x) \int_x^b (t-x)^\alpha dt \leq \int_x^b (t-x)^\alpha f'(t) dt \leq M_2(x) \int_x^b (t-x)^\alpha dt$$

and

$$m_1(x) \int_a^x (x-t)^\alpha dt \leq \int_a^x (x-t)^\alpha f'(t) dt \leq M_1(x) \int_a^x (x-t)^\alpha dt$$

namely

$$\frac{1}{\alpha+1} m_2(x)(b-x)^{\alpha+1} \leq \int_x^b (t-x)^\alpha f'(t) dt \leq \frac{1}{\alpha+1} M_2(x)(b-x)^{\alpha+1}$$

and

$$\frac{1}{\alpha+1} m_1(x)(x-a)^{\alpha+1} \leq \int_a^x (x-t)^\alpha f'(t) dt \leq \frac{1}{\alpha+1} M_1(x)(x-a)^{\alpha+1}.$$

These imply that

$$\begin{aligned} & \frac{1}{\alpha+1} \left[m_2(x)(b-x)^{\alpha+1} - M_1(x)(x-a)^{\alpha+1} \right] \\ & \leq \int_x^b (t-x)^\alpha f'(t) dt - \int_a^x (x-t)^\alpha f'(t) dt \\ & \leq \frac{1}{\alpha+1} \left[M_2(x)(b-x)^{\alpha+1} - m_1(x)(x-a)^{\alpha+1} \right] \end{aligned}$$

that is equivalent to

$$\begin{aligned} & \frac{1}{\Gamma(\alpha+2)} \left[m_2(x) (b-x)^{\alpha+1} - M_1(x) (x-a)^{\alpha+1} \right] \\ & \leq \frac{1}{\Gamma(\alpha+1)} \left[\int_x^b (t-x)^\alpha f'(t) dt - \int_a^x (x-t)^\alpha f'(t) dt \right] \\ & \leq \frac{1}{\Gamma(\alpha+2)} \left[M_2(x) (b-x)^{\alpha+1} - m_1(x) (x-a)^{\alpha+1} \right]. \end{aligned}$$

By using the equality (3.5) we get (3.3).

From (2.2) we have

$$\begin{aligned} & J_{x-}^\alpha f(a) + J_{x+}^\alpha f(b) - \frac{1}{\Gamma(\alpha+1)} [(x-a)^\alpha + (b-x)^\alpha] f(x) \\ & = \frac{1}{\Gamma(\alpha+1)} \left[\int_x^b (b-t)^\alpha f'(t) dt - \int_a^x (t-a)^\alpha f'(t) dt \right]. \end{aligned} \quad (3.6)$$

In a similar way, we have

$$\frac{1}{\alpha+1} m_2(x) (b-x)^{\alpha+1} \leq \int_x^b (b-t)^\alpha f'(t) dt \leq \frac{1}{\alpha+1} M_2(x) (b-x)^{\alpha+1}$$

and

$$\frac{1}{\alpha+1} m_1(x) (x-a)^{\alpha+1} \leq \int_a^x (t-a)^\alpha f'(t) dt \leq \frac{1}{\alpha+1} M_1(x) (x-a)^{\alpha+1},$$

which implies that

$$\begin{aligned} & \frac{1}{\Gamma(\alpha+2)} \left[m_2(x) (b-x)^{\alpha+1} - M_1(x) (x-a)^{\alpha+1} \right] \\ & \leq \frac{1}{\Gamma(\alpha+1)} \left[\int_x^b (b-t)^\alpha f'(t) dt - \int_a^x (t-a)^\alpha f'(t) dt \right] \\ & \leq \frac{1}{\Gamma(\alpha+2)} \left[M_2(x) (b-x)^{\alpha+1} - m_1(x) (x-a)^{\alpha+1} \right] \end{aligned}$$

and by (3.6) we get (3.4). \square

REMARK 1. If we take $\alpha = 1$ in (3.3), then we get

$$\begin{aligned} & \frac{1}{2} \left[m_2(x) (b-x)^2 - M_1(x) (x-a)^2 \right] \\ & \leq (x-a)f(a) + (b-x)f(b) - \int_a^b f(t) dt \\ & \leq \frac{1}{2} \left[M_2(x) (b-x)^2 - m_1(x) (x-a)^2 \right] \end{aligned} \quad (3.7)$$

for any $x \in (a, b)$. If we take $\alpha = 1$ in (3.4), then we get (1.7).

COROLLARY 3. Let $f : [a, b] \rightarrow \mathbb{R}$ be an absolutely continuous function on $[a, b]$. If there exists the real numbers m_1, M_1, m_2, M_2 such that

$$m_1 \leq f'(t) \leq M_1 \text{ for a.e. } t \in \left(a, \frac{a+b}{2}\right) \tag{3.8}$$

and

$$m_2 \leq f'(t) \leq M_2 \text{ for a.e. } t \in \left(\frac{a+b}{2}, b\right) \tag{3.9}$$

then

$$\begin{aligned} & \frac{1}{2^{\alpha+1}\Gamma(\alpha+2)}(b-a)^{\alpha+1}(m_2 - M_1) \tag{3.10} \\ & \leq \frac{1}{2^{\alpha-1}\Gamma(\alpha+1)} \frac{f(a)+f(b)}{2} (b-a)^\alpha - J_{a^+}^\alpha f\left(\frac{a+b}{2}\right) - J_{b^-}^\alpha f\left(\frac{a+b}{2}\right) \\ & \leq \frac{1}{2^{\alpha+1}\Gamma(\alpha+2)}(b-a)^{\alpha+1}(M_2 - m_1) \end{aligned}$$

and

$$\begin{aligned} & \frac{1}{2^{\alpha+1}\Gamma(\alpha+2)}(b-a)^{\alpha+1}(m_2 - M_1) \tag{3.11} \\ & \leq J_{\frac{a+b}{2}^-}^\alpha f(a) + J_{\frac{a+b}{2}^+}^\alpha f(b) - \frac{1}{2^{\alpha-1}\Gamma(\alpha+1)} f\left(\frac{a+b}{2}\right) (b-a)^\alpha \\ & \leq \frac{1}{2^{\alpha+1}\Gamma(\alpha+2)}(b-a)^{\alpha+1}(M_2 - m_1). \end{aligned}$$

In particular, we have the simpler inequalities:

COROLLARY 4. Let $f : [a, b] \rightarrow \mathbb{R}$ be an absolutely continuous function on $[a, b]$. If there exists the real numbers m, M , such that $m \leq f'(t) \leq M$ for a.e. $t \in (a, b)$, then

$$\begin{aligned} & \left| \frac{1}{2^{\alpha-1}\Gamma(\alpha+1)} \frac{f(a)+f(b)}{2} (b-a)^\alpha - J_{a^+}^\alpha f\left(\frac{a+b}{2}\right) - J_{b^-}^\alpha f\left(\frac{a+b}{2}\right) \right| \tag{3.12} \\ & \leq \frac{1}{2^{\alpha+1}\Gamma(\alpha+2)}(b-a)^{\alpha+1}(M - m) \end{aligned}$$

and

$$\begin{aligned} & \left| J_{\frac{a+b}{2}^-}^\alpha f(a) + J_{\frac{a+b}{2}^+}^\alpha f(b) - \frac{1}{2^{\alpha-1}\Gamma(\alpha+1)} f\left(\frac{a+b}{2}\right) (b-a)^\alpha \right| \tag{3.13} \\ & \leq \frac{1}{2^{\alpha+1}\Gamma(\alpha+2)}(b-a)^{\alpha+1}(M - m). \end{aligned}$$

REMARK 2. If we take $\alpha = 1$ in (3.12), then we get

$$\left| \frac{f(a)+f(b)}{2} (b-a) - \int_a^b f(t) dt \right| \leq \frac{1}{8} (b-a)^2 (M - m) \tag{3.14}$$

while from (3.13) we get (1.10).

We also have the following trapezoid type result:

THEOREM 3. *Let $f : [a, b] \rightarrow \mathbb{R}$ be an absolutely continuous function on $[a, b]$. If there exists the real numbers m, M , such that $m \leq f'(t) \leq M$ for a.e. $t \in (a, b)$, then*

$$\begin{aligned} & \left| \frac{1}{\Gamma(\alpha+1)} \frac{f(b)+f(a)}{2} (b-a)^\alpha - \frac{J_{b-}^\alpha f(a) + J_{a+}^\alpha f(b)}{2} \right| \\ & \leq \frac{2^\alpha - 1}{2^{\alpha+1} \Gamma(\alpha+2)} (M-m) (b-a)^{\alpha+1}. \end{aligned} \quad (3.15)$$

Proof. We have by (2.11) that

$$\begin{aligned} & \frac{1}{\Gamma(\alpha+1)} \frac{f(b)+f(a)}{2} (b-a)^\alpha - \frac{J_{b-}^\alpha f(a) + J_{a+}^\alpha f(b)}{2} \\ & = \frac{1}{\Gamma(\alpha+1)} \int_a^b \frac{(t-a)^\alpha - (b-t)^\alpha}{2} f'(t) dt. \end{aligned}$$

Observe also that

$$\begin{aligned} & \int_a^b \frac{(t-a)^\alpha - (b-t)^\alpha}{2} \left(f'(t) - \frac{m+M}{2} \right) dt \\ & = \int_a^b \frac{(t-a)^\alpha - (b-t)^\alpha}{2} f'(t) dt - \frac{m+M}{2} \int_a^b \frac{(t-a)^\alpha - (b-t)^\alpha}{2} dt \end{aligned}$$

and since

$$\int_a^b [(t-a)^\alpha - (b-t)^\alpha] dt = \frac{(b-a)^{\alpha+1}}{\alpha+1} - \frac{(b-a)^{\alpha+1}}{\alpha+1} = 0,$$

then we have the following identity of interest

$$\begin{aligned} & \frac{1}{\Gamma(\alpha+1)} \frac{f(b)+f(a)}{2} (b-a)^\alpha - \frac{J_{b-}^\alpha f(a) + J_{a+}^\alpha f(b)}{2} \\ & = \frac{1}{\Gamma(\alpha+1)} \int_a^b \frac{(t-a)^\alpha - (b-t)^\alpha}{2} \left(f'(t) - \frac{m+M}{2} \right) dt. \end{aligned} \quad (3.16)$$

By taking the modulus in (3.16), we get

$$\begin{aligned} & \left| \frac{1}{\Gamma(\alpha+1)} \frac{f(b)+f(a)}{2} (b-a)^\alpha - \frac{J_{b-}^\alpha f(a) + J_{a+}^\alpha f(b)}{2} \right| \\ & \leq \frac{1}{\Gamma(\alpha+1)} \int_a^b \left| \frac{(t-a)^\alpha - (b-t)^\alpha}{2} \right| \left| f'(t) - \frac{m+M}{2} \right| dt \\ & \leq \frac{1}{4} (M-m) \frac{1}{\Gamma(\alpha+1)} \int_a^b |(b-t)^\alpha - (t-a)^\alpha| dt. \end{aligned} \quad (3.17)$$

The function $h : [a, b] \rightarrow [0, \infty)$, $h(t) = |(b-t)^\alpha - (t-a)^\alpha|$ is symmetric on $[a, b]$, then

$$\begin{aligned} & \int_a^b |(b-t)^\alpha - (t-a)^\alpha| dt \\ &= 2 \int_a^{\frac{a+b}{2}} [(b-t)^\alpha - (t-a)^\alpha] dt \\ &= 2 \left[-\frac{(b-t)^{\alpha+1}}{\alpha+1} \Big|_a^{\frac{a+b}{2}} - \frac{(t-a)^{\alpha+1}}{\alpha+1} \Big|_a^{\frac{a+b}{2}} \right] \\ &= 2 \left[-\frac{(b-\frac{a+b}{2})^{\alpha+1}}{\alpha+1} + \frac{(b-a)^{\alpha+1}}{\alpha+1} - \frac{(\frac{a+b}{2}-a)^{\alpha+1}}{\alpha+1} \right] \\ &= 2 \left[\frac{(b-a)^{\alpha+1}}{\alpha+1} - \frac{(b-a)^{\alpha+1}}{2^\alpha(\alpha+1)} \right] = \frac{2^\alpha - 1}{2^{\alpha-1}(\alpha+1)} (b-a)^{\alpha+1}. \end{aligned}$$

By making use of (3.17) we then obtain (3.15). \square

4. Inequalities for convex functions

We have the following result for convex functions:

THEOREM 4. *Let $f : [a, b] \rightarrow \mathbb{R}$ be a convex function and $x \in (a, b)$, then we have the inequalities*

$$\begin{aligned} & \frac{1}{\Gamma(\alpha+2)} \left[f'_+(x)(b-x)^{\alpha+1} - f'_-(x)(x-a)^{\alpha+1} \right] \\ & \leq \frac{1}{\Gamma(\alpha+1)} \left[(x-a)^\alpha f(a) + (b-x)^\alpha f(b) \right] - J_{a+}^\alpha f(x) - J_{b-}^\alpha f(x) \\ & \leq \frac{1}{\Gamma(\alpha+2)} \left[f'_-(b)(b-x)^{\alpha+1} - f'_+(a)(x-a)^{\alpha+1} \right] \end{aligned} \quad (4.1)$$

and

$$\begin{aligned} & \frac{1}{\Gamma(\alpha+2)} \left[f'_+(x)(b-x)^{\alpha+1} - f'_-(x)(x-a)^{\alpha+1} \right] \\ & \leq J_{x-}^\alpha f(a) + J_{x+}^\alpha f(b) - \frac{1}{\Gamma(\alpha+1)} \left[(x-a)^\alpha + (b-x)^\alpha \right] f(x) \\ & \leq \frac{1}{\Gamma(\alpha+2)} \left[f'_-(b)(b-x)^{\alpha+1} - f'_+(a)(x-a)^{\alpha+1} \right], \end{aligned} \quad (4.2)$$

where $f'_\pm(\cdot)$ are the lateral derivatives of f .

Proof. Since f is convex, then the derivative f' exists almost everywhere on $[a, b]$ and

$$f'_+(a) \leq f'(t) \leq f'_-(x) \text{ for a.e. } t \in (a, x)$$

and

$$f'_+(x) \leq f'(t) \leq f'_-(b) \text{ for a.e. } t \in (x, b).$$

Now, writing the inequalities (3.3) and (3.4) for $m_1(x) = f'_+(a)$, $M_1(x) = f'_-(x)$, $m_2(x) = f'_+(x)$ and $M_2(x) = f'_-(b)$ we get the desired results (4.1) and (4.2). \square

REMARK 3. If we take $\alpha = 1$ in (4.1) and (4.2), then we recapture (1.3) and (1.1) that hold for convex functions.

COROLLARY 5. Let $f : [a, b] \rightarrow \mathbb{R}$ be a convex function, then we have the inequalities

$$\begin{aligned} 0 &\leq \frac{1}{2^{\alpha+1}\Gamma(\alpha+2)} \left[f'_+ \left(\frac{a+b}{2} \right) - f'_- \left(\frac{a+b}{2} \right) \right] (b-a)^{\alpha+1} & (4.3) \\ &\leq \frac{1}{2^{\alpha-1}\Gamma(\alpha+1)} \frac{f(a)+f(b)}{2} (b-a)^\alpha - J_{a^+}^\alpha f \left(\frac{a+b}{2} \right) - J_{b^-}^\alpha f \left(\frac{a+b}{2} \right) \\ &\leq \frac{1}{2^{\alpha+1}\Gamma(\alpha+2)} [f'_-(b) - f'_+(a)] (b-a)^{\alpha+1}, \end{aligned}$$

$$\begin{aligned} 0 &\leq \frac{1}{2^{\alpha+1}\Gamma(\alpha+2)} \left[f'_+ \left(\frac{a+b}{2} \right) - f'_- \left(\frac{a+b}{2} \right) \right] (b-a)^{\alpha+1} & (4.4) \\ &\leq J_{\frac{a+b}{2}^-}^\alpha f(a) + J_{\frac{a+b}{2}^+}^\alpha f(b) - \frac{1}{2^{\alpha-1}\Gamma(\alpha+1)} f \left(\frac{a+b}{2} \right) (b-a)^\alpha \\ &\leq \frac{1}{2^{\alpha+1}\Gamma(\alpha+2)} [f'_-(b) - f'_+(a)] (b-a)^{\alpha+1}, \end{aligned}$$

and

$$\begin{aligned} 0 &\leq \frac{1}{\Gamma(\alpha+1)} \frac{f(b)+f(a)}{2} (b-a)^\alpha - \frac{J_{b^-}^\alpha f(a) + J_{a^+}^\alpha f(b)}{2} & (4.5) \\ &\leq \frac{2^\alpha - 1}{2^{\alpha+1}\Gamma(\alpha+2)} (f'_-(b) - f'_+(a)) (b-a)^{\alpha+1}. \end{aligned}$$

If we take $\alpha = 1$ in (4.3) and (4.4), then we recapture the midpoint and trapezoid inequalities for convex functions mentioned in the introduction.

5. Conclusion

In this paper, by making use of some fundamental identities for the Riemann-Liouville fractional integrals, we established some Ostrowski and trapezoid type inequalities for these integrals of absolutely continuous functions with bounded derivatives. Applications for mid-point and trapezoid inequalities were provided as well. They generalize the known results holding for the classical Riemann integral. Some natural applications for the important case of convex functions were also given.

REFERENCES

- [1] A. AGLIĆ ALJINOVIĆ, *Montgomery identity and Ostrowski type inequalities for Riemann-Liouville fractional integral*, J. Math. **2014**, Art. ID 503195, 6 pp.
- [2] T. M. APOSTOL, *Mathematical Analysis*, Second Edition, Addison-Wesley Publishing Company, 1975.
- [3] A. O. AKDEMIR, *Inequalities of Ostrowski's type for m - and (α, m) -logarithmically convex functions via Riemann-Liouville fractional integrals*, J. Comput. Anal. Appl. **16** (2014), no. 2, 375–383.
- [4] G. A. ANASTASSIOU, *Fractional representation formulae under initial conditions and fractional Ostrowski type inequalities*, Demonstr. Math. **48** (2015), no. 3, 357–378.
- [5] G. A. ANASTASSIOU, *The reduction method in fractional calculus and fractional Ostrowski type inequalities*, Indian J. Math. **56** (2014), no. 3, 333–357.
- [6] H. BUDAK, M. Z. SARIKAYA, E. SET, *Generalized Ostrowski type inequalities for functions whose local fractional derivatives are generalized s -convex in the second sense*, J. Appl. Math. Comput. Mech. **15** (2016), no. 4, 11–21.
- [7] P. CERONE AND S. S. DRAGOMIR, *Midpoint-type rules from an inequalities point of view. Handbook of analytic-computational methods in applied mathematics*, 135–200, Chapman & Hall/CRC, Boca Raton, FL, 2000.
- [8] S. S. DRAGOMIR, *The Ostrowski's integral inequality for Lipschitzian mappings and applications*, Comput. Math. Appl. **38** (1999), no. 11–12, 33–37.
- [9] S. S. DRAGOMIR, *The Ostrowski integral inequality for mappings of bounded variation*, Bull. Austral. Math. Soc. **60** (1999), no. 3, 495–508.
- [10] S. S. DRAGOMIR, *On the midpoint quadrature formula for mappings with bounded variation and applications*, Kragujevac J. Math. **22** (2000), 13–19.
- [11] S. S. DRAGOMIR, *On the Ostrowski's integral inequality for mappings with bounded variation and applications*, Math. Ineq. Appl. **4** (2001), No. 1, 59–66. Preprint: RGMIA Res. Rep. Coll. **2** (1999), Art. 7, online <http://rgmia.org/papers/v2n1/v2n1-7.pdf>.
- [12] S. S. DRAGOMIR, *An inequality improving the first Hermite-Hadamard inequality for convex functions defined on linear spaces and applications for semi-inner products*, J. Inequal. Pure Appl. Math. **3** (2002), no. 2, Article 31, 8 pp., online <http://www.emis.de/journals/JIPAM/article183.html?sid=183>.
- [13] S. S. DRAGOMIR, *An inequality improving the second Hermite-Hadamard inequality for convex functions defined on linear spaces and applications for semi-inner products*, J. Inequal. Pure Appl. Math. **3** (2002), no. 3, Art. 35, 8 pp., online https://www.emis.de/journals/JIPAM/images/080_01_JIPAM/080_01.pdf.
- [14] S. S. DRAGOMIR, *Improvements of Ostrowski and generalised trapezoid inequality in terms of the upper and lower bounds of the first derivative*, Tamkang J. Math. **34** (2003), no. 3, 213–222.
- [15] S. S. DRAGOMIR, *Refinements of the generalised trapezoid and Ostrowski inequalities for functions of bounded variation*, Arch. Math. (Basel) **91** (2008), no. 5, 450–460.
- [16] S. S. DRAGOMIR, *Refinements of the Ostrowski inequality in terms of the cumulative variation and applications*, Analysis (Berlin) **34** (2014), no. 2, 223–240. Preprint: RGMIA Res. Rep. Coll. **16** (2013), Art. 29, online <http://rgmia.org/papers/v16/v16a29.pdf>.
- [17] S. S. DRAGOMIR, *Ostrowski type inequalities for Lebesgue integral: a survey of recent results*, Australian J. Math. Anal. Appl. Volume **14**, Issue 1, Article 1, pp. 1–287, 2017., online <http://ajmaa.org/cgi-bin/paper.pl?string=v14n1/V14I1P1.tex>.
- [18] S. S. DRAGOMIR, *Ostrowski type inequalities for Riemann-Liouville fractional integrals of bounded variation, Hölder and Lipschitzian functions*, Preprint RGMIA Res. Rep. Coll. **20** (2017), Art. 48., online <http://rgmia.org/papers/v20/v20a48.pdf>.
- [19] A. GUEZANE-LAKOUD AND F. AISSAOUI, *New fractional inequalities of Ostrowski type*, Transylv. J. Math. Mech. **5** (2013), no. 2, 103–106.
- [20] A. KASHURI AND R. LIKO, *Ostrowski type fractional integral inequalities for generalized (s, m, φ) -preinvex functions*, Aust. J. Math. Anal. Appl. **13** (2016), no. 1, Art. 16, 11 pp.
- [21] M. A. NOOR, K. I. NOOR AND S. IFTIKHAR, *Fractional Ostrowski inequalities for harmonic h -preinvex functions*, Facta Univ. Ser. Math. Inform. **31** (2016), no. 2, 417–445.
- [22] M. Z. SARIKAYA AND H. FILIZ, *Note on the Ostrowski type inequalities for fractional integrals*, Vietnam J. Math. **42** (2014), no. 2, 187–190.

- [23] M. Z. SARIKAYA AND H. BUDAK, *Generalized Ostrowski type inequalities for local fractional integrals*, Proc. Amer. Math. Soc. **145** (2017), no. 4, 1527–1538.
- [24] E. SET, *New inequalities of Ostrowski type for mappings whose derivatives are s -convex in the second sense via fractional integrals*, Comput. Math. Appl. **63** (2012), no. 7, 1147–1154.
- [25] M. TUNÇ, *On new inequalities for h -convex functions via Riemann-Liouville fractional integration*, Filomat **27**:4 (2013), 559–565.
- [26] M. TUNÇ, *Ostrowski type inequalities for m - and (α, m) -geometrically convex functions via Riemann-Liouville fractional integrals*, Afr. Mat. **27** (2016), no. 5–6, 841–850.
- [27] H. YILDIRIM AND Z. KIRTAY, *Ostrowski inequality for generalized fractional integral and related inequalities*, Malaya J. Mat., **2** (3) (2014), 322–329.
- [28] C. YILDIZ, E. ÖZDEMİR AND Z. S. MUHAMET, *New generalizations of Ostrowski-like type inequalities for fractional integrals*, Kyungpook Math. J. **56** (2016), no. 1, 161–172.
- [29] H. YUE, *Ostrowski inequality for fractional integrals and related fractional inequalities*, Transylv. J. Math. Mech. **5** (2013), no. 1, 85–89.

(Received October 27, 2020)

Silvestru Sever Dragomir
Mathematics, College of Engineering & Science
Victoria University, PO Box 14428
Melbourne City, MC 8001, Australia
and
DST-NRF Centre of Excellence
in the Mathematical and Statistical Sciences
School of Computer Science & Applied Mathematics
University of the Witwatersrand, Private Bag 3
Johannesburg 2050, South Africa
e-mail: sever.dragomir@vu.edu.au