

ON STATISTICAL CONVERGENCE OF SEQUENCES OF FUNCTIONS IN 2-NORMED SPACES

SEVİM YEGÜL AND ERDİNÇ DÜNDAR

Abstract. Statistical convergence and statistical Cauchy sequence in 2-normed space were studied by Gürdal and Pehlivan [M. Gürdal, S. Pehlivan, *Statistical convergence in 2-normed spaces*, Southeast Asian Bulletin of Mathematics, (33) (2009), 257–264]. In this paper, we get analogous results of statistical convergence and statistical Cauchy sequence of functions and investigate some properties and relationships between them in 2-normed spaces.

1. Introduction

Throughout the paper, \mathbb{N} denotes the set of all positive integers, \mathbb{R} the set of all real numbers. The concept of convergence of a sequence of real numbers has been extended to statistical convergence independently by Fast [7] and Schoenberg [23]. Gökhan et al. [12] introduced the concepts of pointwise statistical convergence and statistical Cauchy sequence of real-valued functions. Balcerzak et al. [2] studied statistical convergence and ideal convergence for sequence of functions. Baláz et al. [1] investigated \mathcal{I} -convergence and \mathcal{I} -continuity of real functions. Gezer and Karakuş [11] investigated \mathcal{I} -pointwise and uniform convergence and \mathcal{I}^* -pointwise and uniform convergence of function sequences and then they examined the relation between them. Gökhan et al. [13] introduced the notion of pointwise and uniform statistical convergence of double sequences of real-valued functions. Dündar and Altay [4, 5] studied the concepts of pointwise and uniformly \mathcal{I} -convergence and \mathcal{I}^* -convergence of double sequences of functions and investigated some properties about them. Furthermore, Dündar [6] investigated some results of \mathcal{I}_2 -convergence of double sequences of functions.

The concept of 2-normed spaces was initially introduced by Gähler [9, 10] in the 1960's. Gürdal and Pehlivan [16] studied statistical convergence, statistical Cauchy sequence and investigated some properties of statistical convergence in 2-normed spaces. Sharma and Kumar [24] introduced statistical convergence, statistical Cauchy sequence, statistical limit points and statistical cluster points in probabilistic 2-normed space. Savaş and Gürdal [22] concerned with \mathcal{I} -convergence of sequences of functions in random 2-normed spaces and introduce the concepts of ideal uniform convergence and ideal pointwise convergence in the topology induced by random 2-normed spaces.

Mathematics subject classification (2010): 40A30, 40A35, 46A70.

Keywords and phrases: Statistical convergence, sequence of functions, statistical Cauchy sequence, 2-normed spaces.

Sarabadan and Talebi [21] presented various kinds of statistical convergence and \mathcal{I} -convergence for sequences of functions with values in 2-normed spaces and also defined the notion of \mathcal{I} -equistatistically convergence and study \mathcal{I} -equistatistically convergence of sequences of functions. Şahiner et al. [25] and Gürdal [18] studied \mathcal{I} -convergence in 2-normed spaces. Gürdal and Açıık [17] investigated \mathcal{I} -Cauchy and \mathcal{I}^* -Cauchy sequences in 2-normed spaces. Furthermore, a lot of development have been made in this area (see [3, 14, 15, 19, 20]).

2. Definitions and notations

Now, we recall the concept of density, statistical convergence, 2-normed space and some fundamental definitions and notations (See [2, 8, 10, 11, 12, 14, 15, 16, 21, 24]).

If $K \subseteq \mathbb{N}$, then K_n denotes the set $\{k \in K : k \leq n\}$ and $|K_n|$ denotes the cardinality of K_n . The natural density of K is given by $\delta(K) = \lim_{n \rightarrow \infty} \frac{1}{n} |K_n|$, if it exists.

Clearly, finite subsets have natural density zero and $\delta(K^c) = 1 - \delta(K)$ where $K^c = \mathbb{N} \setminus K$, i.e., the complement of K . If $K_1 \subseteq K_2$ and K_1 and K_2 have natural densities then $\delta(K_1) \leq \delta(K_2)$. Moreover, if $\delta(K_1) = \delta(K_2) = 1$, then $\delta(K_1 \cap K_2) = 1$.

The number sequence $x = (x_k)$ is statistically convergent to L provided that for every $\varepsilon > 0$ the set

$$K = K(\varepsilon) := \{k \in \mathbb{N} : |x_k - L| \geq \varepsilon\}$$

has natural density zero; in this case, we write $st - \lim x = L$.

We note following theorem which is useful in establishing our results.

THEOREM 1. [8] *The following statements are equivalent:*

(i) *x is statistically convergent sequence;*

(ii) *x is statistically Cauchy sequence;*

(iii) *x is sequence for which there is a convergent sequence y such that $x_n = y_n$,*

for a.a. n .

Let X be a real vector space of dimension d , where $2 \leq d < \infty$. A 2-norm on X is a function $\|\cdot, \cdot\| : X \times X \rightarrow \mathbb{R}$ which satisfies the following statements:

(i) $\|x, y\| = 0$ if and only if x and y are linearly dependent.

(ii) $\|x, y\| = \|y, x\|$.

(iii) $\|\alpha x, y\| = |\alpha| \|x, y\|$, $\alpha \in \mathbb{R}$.

(iv) $\|x, y + z\| \leq \|x, y\| + \|x, z\|$.

The pair $(X, \|\cdot, \cdot\|)$ is then called a 2-normed space. As an example of a 2-normed space we may take $X = \mathbb{R}^2$ being equipped with the 2-norm $\|x, y\| :=$ the area of the parallelogram based on the vectors x and y which may be given explicitly by the formula

$$\|x, y\| = |x_1 y_2 - x_2 y_1|; \quad x = (x_1, x_2), y = (y_1, y_2) \in \mathbb{R}^2.$$

In this study, we suppose X to be a 2-normed space having dimension d ; where $2 \leq d < \infty$.

Let $(X, \|\cdot, \cdot\|)$ be a finite dimensional 2-normed space and $u = \{u_1, \dots, u_d\}$ be a basis of X . We can define the norm $\|\cdot\|_\infty$ on X by

$$\|x\|_\infty = \max\{\|x, u_i\| : i = 1, \dots, d\}.$$

Associated to the derived norm $\|\cdot\|_\infty$, we can define the (closed) balls $B_u(x, \varepsilon)$ centered at x having radius ε by

$$B_u(x, \varepsilon) = \{y : \|x - y\|_\infty \leq \varepsilon\},$$

where $\|x - y\|_\infty = \max\{\|x - y, u_j\|, j = 1, \dots, d\}$.

Let X be a 2-normed space. A sequence (x_n) in X is said to be convergent to $L \in X$, if for every nonzero $z \in X$,

$$\lim_{n \rightarrow \infty} \|x_n - L, z\| = 0.$$

In this case, we write $\lim_{n \rightarrow \infty} x_n = L$ and call L the limit of (x_n) .

Let $\{x_n\}$ be a sequence in 2-normed space $(X, \|\cdot, \cdot\|)$. The sequence (x_n) is said to be statistically convergent to L , if for every $\varepsilon > 0$, the set

$$\{n \in \mathbb{N} : \|x_n - L, z\| \geq \varepsilon\}$$

has natural density zero for each nonzero z in X , in other words (x_n) statistically converges to L in 2-normed space $(X, \|\cdot, \cdot\|)$ if

$$\lim_{n \rightarrow \infty} \frac{1}{n} |\{n : \|x_n - L, z\| \geq \varepsilon\}| = 0,$$

for each nonzero z in X . It means that for each $z \in X$,

$$\|x_n - L, z\| < \varepsilon, \text{ a.a. } n.$$

In this case we write $st - \lim_{n \rightarrow \infty} \|x_n, z\| = \|L, z\|$.

A sequence (x_n) in 2-normed space $(X, \|\cdot, \cdot\|)$ is said to be statistically Cauchy sequence in X , if for every $\varepsilon > 0$ and every nonzero $z \in X$ there exists a number $N = N(\varepsilon, z)$ such that

$$\delta(\{n \in \mathbb{N} : \|x_n - x_N, z\| \geq \varepsilon\}) = 0,$$

i.e., for each nonzero $z \in X$,

$$\|x_n - x_N, z\| < \varepsilon, \text{ a.a. } n.$$

Let X and Y be two 2-normed spaces and assume that functions $f_n : X \rightarrow Y$ and $f : X \rightarrow Y$ are given. The sequence of functions $\{f_n\}_{n \in \mathbb{N}}$ is said to be convergent to f if $f_n(x) \xrightarrow{\|\cdot, \cdot\|_Y} f(x)$ for each $x \in X$. We write $f_n \xrightarrow{\|\cdot, \cdot\|_Y} f$. This can be expressed by the formula

$$(\forall y \in Y)(\forall x \in X)(\forall \varepsilon > 0)(\exists n_0 \in \mathbb{N})(\forall n \geq n_0)\|f_n(x) - f(x), y\| < \varepsilon.$$

3. Main results

In this paper, we study concepts of convergence, statistical convergence and statistical Cauchy sequence of functions and investigate some properties and relationships between them in 2-normed spaces.

Throughout the paper, we let X and Y be two 2-normed spaces, $\{f_n\}_{n \in \mathbb{N}}$ and $\{g_n\}_{n \in \mathbb{N}}$ be two sequences of functions and f, g be two functions from X to Y .

DEFINITION 1. The sequence $\{f_n\}_{n \in \mathbb{N}}$ is said to be (pointwise) statistical convergent to f , if for every $\varepsilon > 0$,

$$\lim_{n \rightarrow \infty} \frac{1}{n} |\{n \in \mathbb{N} : \|f_n(x) - f(x), z\| \geq \varepsilon\}| = 0,$$

for each $x \in X$ and each nonzero $z \in Y$. It means that for each $x \in X$ and each nonzero $z \in Y$,

$$\|f_n(x) - f(x), z\| < \varepsilon, \quad \text{a.a. } n.$$

In this case, we write

$$st - \lim_{n \rightarrow \infty} \|f_n(x), z\| = \|f(x), z\| \quad \text{or} \quad f_n \xrightarrow{\|\cdot\|_Y} st f.$$

REMARK 1. $\{f_n\}_{n \in \mathbb{N}}$ is any sequence of functions and f is any function from X to Y , then set

$$\{n \in \mathbb{N} : \|f_n(x) - f(x), z\| \geq \varepsilon, \text{ for each } x \in X \text{ and each } z \in Y\} = \emptyset,$$

since if $z = \vec{0}$ (0 vektor), $\|f_n(x) - f(x), z\| = 0 \not\geq \varepsilon$ so the above set is empty.

THEOREM 1. *If for each $x \in X$ and each nonzero $z \in Y$,*

$$st - \lim_{n \rightarrow \infty} \|f_n(x), z\| = \|f(x), z\| \quad \text{and} \quad st - \lim_{n \rightarrow \infty} \|f_n(x), z\| = \|g(x), z\|,$$

then $\|f_n(x), z\| = \|g_n(x), z\|$ (i.e., $f = g$), for each $x \in X$ and each nonzero $z \in Y$.

Proof. Assume $f \neq g$. Then $f - g \neq \vec{0}$, so there exists a $z \in Y$ such that f, g and z are linearly independent (such a z exists since $d \geq 2$). Therefore, for each $x \in X$ and each nonzero $z \in Y$,

$$\|f(x) - g(x), z\| = 2\varepsilon, \quad \text{with } \varepsilon > 0.$$

Now, for each $x \in X$ and each nonzero $z \in Y$, we get

$$\begin{aligned} 2\varepsilon &= \|f(x) - g(x), z\| = \|(f(x) - f_n(x)) + (f_n(x) - g(x)), z\| \\ &\leq \|f_n(x) - g(x), z\| + \|f_n(x) - f(x), z\| \end{aligned}$$

and so

$$\{n : \|f_n(x) - g(x), z\| < \varepsilon\} \subseteq \{n : \|f_n(x) - f(x), z\| \geq \varepsilon\}.$$

But, for each $x \in X$ and each nonzero $z \in Y$, $\delta(\{n : \|f_n(x) - g(x), z\| < \varepsilon\}) = 0$, then contradicting the fact that $f_n \xrightarrow{\|\cdot\|_Y} st g$. \square

THEOREM 2. *If $\{g_n\}_{(n \in \mathbb{N})}$ is a convergent sequence of functions such that $f_n = g_n$ a.a. n , then $\{f_n\}_{(n \in \mathbb{N})}$ is statistically convergent.*

Proof. Suppose that for each $x \in X$ and each nonzero $z \in Y$,

$$\delta(\{n \in \mathbb{N} : f_n(x) \neq g_n(x)\}) = 0 \quad \text{and} \quad \lim_{n \rightarrow \infty} \|g_n(x), z\| = \|f(x), z\|,$$

then for every $\varepsilon > 0$,

$$\begin{aligned} \{n \in \mathbb{N} : \|f_n(x) - f(x), z\| \geq \varepsilon\} &\subseteq \{n \in \mathbb{N} : \|g_n(x) - f(x), z\| \geq \varepsilon\} \\ &\cup \{n \in \mathbb{N} : f_n(x) \neq g_n(x)\}. \end{aligned}$$

Therefore,

$$\begin{aligned} \delta(\{n \in \mathbb{N} : \|f_n(x) - f(x), z\| \geq \varepsilon\}) &\leq \delta(\{n \in \mathbb{N} : \|g_n(x) - f(x), z\| \geq \varepsilon\}) \quad (1) \\ &\quad + \delta(\{n \in \mathbb{N} : f_n(x) \neq g_n(x)\}) \end{aligned}$$

Since $\lim_{n \rightarrow \infty} \|g_n(x), z\| = \|f(x), z\|$, for each $x \in X$ and each nonzero $z \in Y$. The set $\{n \in \mathbb{N} : \|g_n(x) - f(x), z\| \geq \varepsilon\}$ contain finite number of integers and so

$$\delta(\{n \in \mathbb{N} : \|g_n(x) - f(x), z\| \geq \varepsilon\}) = 0.$$

Using inequality (1) we get for every $\varepsilon > 0$

$$\delta(\{n \in \mathbb{N} : \|f_n(x) - f(x), z\| \geq \varepsilon\}) = 0,$$

for each $x \in X$ and each nonzero $z \in Y$ and so consequently

$$st - \lim_{n \rightarrow \infty} \|f_n(x), z\| = \|f(x), z\|. \quad \square$$

THEOREM 3. *Let $\alpha \in \mathbb{R}$. If for each $x \in X$ and each nonzero $z \in Y$,*

$$st - \lim_{n \rightarrow \infty} \|f_n(x), z\| = \|f(x), z\| \quad \text{and} \quad st - \lim_{n \rightarrow \infty} \|g_n(x), z\| = \|g(x), z\|,$$

then

$$(i) \quad st - \lim_{n \rightarrow \infty} \|f_n(x) + g_n(x), z\| = \|f(x) + g(x), z\| \quad \text{and}$$

$$(ii) \quad st - \lim_{n \rightarrow \infty} \|\alpha f_n(x), z\| = \|\alpha f(x), z\|.$$

Proof. (i) Suppose that

$$st - \lim_{n \rightarrow \infty} \|f_n(x), z\| = \|f(x), z\| \quad \text{and} \quad st - \lim_{n \rightarrow \infty} \|g_n(x), z\| = \|g(x), z\|$$

for each $x \in X$ and each nonzero $z \in Y$. Then, $\delta(K_1) = 0$ and $\delta(K_2) = 0$ where

$$K_1 = K_1(\varepsilon, z) : \left\{ n \in \mathbb{N} : \|f_n(x) - f(x), z\| \geq \frac{\varepsilon}{2} \right\}$$

and

$$K_2 = K_2(\varepsilon, z) : \left\{ n \in \mathbb{N} : \|g_n(x) - g(x), z\| \geq \frac{\varepsilon}{2} \right\}$$

for every $\varepsilon > 0$, each $x \in X$ and each nonzero $z \in Y$. Let

$$K = K(\varepsilon, z) = \{n \in \mathbb{N} : \|(f_n(x) + g_n(x)) - (f(x) + g(x)), z\| \geq \varepsilon\}.$$

To prove that $\delta(K) = 0$, it suffices to show that $K \subset K_1 \cup K_2$. Let $n_0 \in K$ then, for each $x \in X$ and each nonzero $z \in Y$,

$$\|(f_{n_0}(x) + g_{n_0}(x)) - (f(x) + g(x)), z\| \geq \varepsilon. \quad (2)$$

Suppose to the contrary, that $n_0 \notin K_1 \cup K_2$. Then, $n_0 \notin K_1$ and $n_0 \notin K_2$. If $n_0 \notin K_1$ and $n_0 \notin K_2$ then, for each $x \in X$ and each nonzero $z \in Y$,

$$\|f_{n_0}(x) - f(x), z\| < \frac{\varepsilon}{2} \quad \text{and} \quad \|g_{n_0}(x) - g(x), z\| < \frac{\varepsilon}{2}.$$

Then, we get

$$\begin{aligned} \|(f_{n_0}(x) + g_{n_0}(x)) - (f(x) + g(x)), z\| &\leq \|f_{n_0}(x) - f(x), z\| + \|g_{n_0}(x) - g(x), z\| \\ &< \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon, \end{aligned}$$

for each $x \in X$ and each nonzero $z \in Y$, which contradicts (2). Hence $n_0 \in K_1 \cup K_2$ and so $K \subset K_1 \cup K_2$.

(ii) Let $\alpha \in \mathbb{R}$ ($\alpha \neq 0$) and for each $x \in X$ and each nonzero $z \in Y$,

$$st - \lim_{n \rightarrow \infty} \|f_n(x), z\| = \|f(x), z\|.$$

Then, we get

$$\delta \left(\left\{ n \in \mathbb{N} : \|f_n(x) - f(x), z\| \geq \frac{\varepsilon}{|\alpha|} \right\} \right) = 0.$$

Therefore, for each $x \in X$ and each nonzero $z \in Y$, we have

$$\begin{aligned} \{n \in \mathbb{N} : \|\alpha f_n(x) - \alpha f(x), z\| \geq \varepsilon\} &= \{n \in \mathbb{N} : |\alpha| \|f_n(x) - f(x), z\| \geq \varepsilon\} \\ &= \left\{ n \in \mathbb{N} : \|f_n(x) - f(x), z\| \geq \frac{\varepsilon}{|\alpha|} \right\}. \end{aligned}$$

Hence, the right hand side of above equality equals 0. Therefore, for each $x \in X$ and each nonzero $z \in Y$, we have

$$st - \lim_{n \rightarrow \infty} \|\alpha f_n(x), z\| = \|\alpha f(x), z\|. \quad \square$$

Now, we give the concept of statistical Cauchy sequence and investigate relationships between statistical Cauchy sequence and statistical convergence in 2-normed space.

DEFINITION 2. The sequences of functions $\{f_n\}$ is said to be statistically Cauchy sequence, if for every $\varepsilon > 0$ and each nonzero $z \in Y$, there exist a number $k = k(\varepsilon, z)$ such that

$$\delta(\{n \in \mathbb{N} : \|f_n(x) - f_k(x), z\| \geq \varepsilon\}) = 0$$

for each $x \in X$ i.e.,

$$\|f_n(x) - f_k(x), z\| < \varepsilon, \quad \text{a.a. } n.$$

THEOREM 4. Let $\{f_n\}_{n \geq 1}$ be a statistically Cauchy sequence of functions in a finite dimensional 2-normed space $(X, \|\cdot, \cdot\|)$. Then, there exists a convergent sequence of functions $\{g_n\}_{n \geq 1}$ in $(X, \|\cdot, \cdot\|)$ such that $f_n = g_n$, for a.a. n .

Proof. First note that $\{f_n\}_{n \geq 1}$ is a statistically Cauchy sequence of functions in $(X, \|\cdot\|_\infty)$. Choose a natural number $k(1)$ such that the closed ball $B_u^1 = B_u(f_{k(1)}(x), 1)$ contains $f_n(x)$ for a.a. n and for each $x \in X$. Then, choose a natural number $k(2)$ such that the closed ball $B_2 = B_u(f_{k(1)}(x), \frac{1}{2})$ contains $f_n(x)$ for a.a. n and for each $x \in X$. Note that $B_u^2 = B_u^1 \cap B_2$ also contains $f_n(x)$ for a.a. n and for each $x \in X$. Thus, by continuing of this process, we can obtain a sequence $\{B_u^m\}_{m \geq 1}$ of nested closed balls such that $\text{diam}(B_u^m) \leq \frac{1}{2^m}$. Therefore,

$$\bigcap_{m=1}^{\infty} B_u^m = \{h(x)\},$$

where h is a function from X to Y . Since each B_u^m contains $f_n(x)$ for a.a. n and for each $x \in X$, we can choose a sequence of strictly increasing natural numbers $\{S_m\}_{m \geq 1}$ such that for each $x \in X$,

$$\frac{1}{n} |\{n \in \mathbb{N} : f_n(x) \notin B_u^m\}| < \frac{1}{m}, \text{ if } n > S_m.$$

Put $R_m = \{n \in \mathbb{N} : n > S_m, f_n(x) \notin B_u^m\}$ for each $x \in X$, for all $m \geq 1$ and $R = \bigcup_{m=1}^{\infty} R_m$. Now, for each $x \in X$, define the sequence of functions $\{g_n\}_{n \geq 1}$ as following

$$g_n(x) = \begin{cases} h(x), & \text{if } n \in R \\ f_n(x), & \text{otherwise.} \end{cases}$$

Note that, $\lim_{n \rightarrow \infty} g_n(x) = h(x)$, for each $x \in X$. In fact, for each $\varepsilon > 0$ and for each $x \in X$, choose a natural number m such that $\varepsilon > \frac{1}{m} > 0$. Then, for each $n > S_m$ and for each $x \in X$, $g_n(x) = h(x)$ or $g_n(x) = f_n(x) \in B_u^m$ and so in each case

$$\|g_n(x) - h(x)\|_\infty \leq \text{diam}(B_u^m) \leq \frac{1}{2^{m-1}}.$$

Since, for each $x \in X$, $\{n \in \mathbb{N} : g_n(x) \neq f_n(x)\} \subseteq \{n \in \mathbb{N} : f_n(x) \notin B_u^m\}$, we have

$$\frac{1}{n} |\{n \in \mathbb{N} : g_n(x) \neq f_n(x)\}| \leq \frac{1}{n} |\{n \in \mathbb{N} : f_n(x) \notin B_u^m\}| < \frac{1}{m},$$

and so

$$\delta(\{n \in \mathbb{N} : g_n(x) \neq f_n(x)\}) = 0.$$

Thus, $g_n(x) = f_n(x)$ for a.a. n and for each $x \in X$ in $(X, \|\cdot\|_\infty)$. Suppose that $\{u_1, \dots, u_d\}$ is a basis for $(X, \|\cdot, \cdot\|)$. Since, for each $x \in X$,

$$\lim_{n \rightarrow \infty} \|g_n(x) - h(x)\|_\infty = 0 \text{ and } \|g_n(x) - h(x), u_i\| \leq \|g_n(x) - h(x)\|_\infty$$

for all $1 \leq i \leq d$, then we have

$$\lim_{n \rightarrow \infty} \|g_n(x) - h(x), z\|_\infty = 0,$$

for each $x \in X$ and each nonzero $z \in X$. This completes the proof. \square

THEOREM 5. *The sequence $\{f_n\}$ is statistically convergent if and only if $\{f_n\}$ is a statistically Cauchy sequence of functions.*

Proof. Assume that f be function from X to Y and

$$st - \lim_{n \rightarrow \infty} \|f_n(x), z\| = \|f(x), z\|,$$

for each $x \in X$ and each nonzero $z \in Y$ and $\varepsilon > 0$. Then, for each $x \in X$ and each nonzero $z \in Y$, we have

$$\|f_n(x) - f(x), z\| < \frac{\varepsilon}{2}, \text{ a.a. } n.$$

If $k = k(\varepsilon, z)$ is chosen so that for each $x \in X$ and each nonzero $z \in Y$,

$$\|f_k(x) - f(x), z\| < \frac{\varepsilon}{2},$$

and so we have

$$\|f_n(x) - f_k(x), z\| \leq \|f_n(x) - f(x), z\| + \|f(x) - f_k(x), z\| < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon, \text{ a.a. } n.$$

Hence, $\{f_n\}$ is statistically Cauchy sequence of functions.

Now, assume that $\{f_n\}$ is statistically Cauchy sequence of function. By Theorem 4, there exists a convergent sequence $\{g_n\}_{n \in \mathbb{N}}$ from X to Y such that $f_n = g_n$ for a.a. n . By Theorem 2, we have

$$st - \lim \|f_n(x), z\| = \|f(x), z\|$$

for each $x \in X$ and each nonzero $z \in Y$. \square

Acknowledgement. This study is supported by Afyon Kocatepe University Scientific Research Coordination Unit with the project number 16.FEN.EDB.04.

REFERENCES

- [1] V. BALÁZ, J. ČERVEŇANSKÝ, P. KOSTYRKO, T. ŠALÁT, *I-convergence and I-continuity of real functions*, Acta Mathematica, Faculty of Natural Sciences, Constantine the Philosopher University, Nitra, **5** (2004), 43–50.
- [2] M. BALCERZAK, K. DEMS, A. KOMISARSKI, *Statistical convergence and ideal convergence for sequences of functions*, J. Math. Anal. Appl. **328**(1) (2007), 715–729.
- [3] H. ÇAKALLI AND S. ERSAN, *New types of continuity in 2-normed spaces*, Filomat **30** (3) (2016), 525–532.
- [4] E. DÜNDAR, B. ALTAY, *\mathcal{I}_2 -convergence of double sequences of functions*, Electronic Journal of Mathematical Analysis and Applications **3** (1) (2015), 111–121.
- [5] E. DÜNDAR, B. ALTAY, *\mathcal{I}_2 -uniform convergence of double sequences of functions*, Filomat **30** (5) (2016), 1273–1281.
- [6] E. DÜNDAR, *On some results of \mathcal{I}_2 -convergence of double sequences of functions*, Mathematical Analysis Sciences and Applications E-notes **3** (1) (2015), 44–52.
- [7] H. FAST, *Sur la convergence statistique*, Colloq. Math. **2** (1951), 241–244.
- [8] J. A. FRIDY, *On statistical convergence*, Analysis **5** (1985), 301–313.
- [9] S. GÄHLER, *2-metrische Räume und ihre topologische struktur*, Math. Nachr. **26** (1963), 115–148.
- [10] S. GÄHLER, *2-normed spaces*, Math. Nachr. **28** (1964), 1–43.
- [11] F. GEZER, S. KARAKUŞ, *\mathcal{I} and \mathcal{I}^* convergent function sequences*, Math. Commun. **10** (2005), 71–80.
- [12] A. GÖKHAN AND M. GÜNGÖR, *On pointwise statistical convergence*, Indian J. Pure Appl. Math. **33** (9) (2002), 1379–1384.

- [13] A. GÖKHAN, M. GÜNGÖR AND M. ET, *Statistical convergence of double sequences of real-valued functions*, Int. Math. Forum, **2** (8) (2007), 365–374.
- [14] H. GUNAWAN, M. MASHADI, *On finite dimensional 2-normed spaces*, Soochow J. Math. **27** (3) (2001), 321–329.
- [15] M. GÜRDAL, S. PEHLIVAN, *The statistical convergence in 2-Banach spaces*, Thai J. Math. **2** (1) (2004), 107–113.
- [16] M. GÜRDAL, S. PEHLIVAN, *Statistical convergence in 2-normed spaces*, Southeast Asian Bulletin of Mathematics **33**(2) (2009), 257–264.
- [17] M. GÜRDAL, I. AÇIK, *On \mathcal{I} -Cauchy sequences in 2-normed spaces*, Math. Inequal. Appl. **11** (2) (2008), 349–354.
- [18] M. GÜRDAL, *On ideal convergent sequences in 2-normed spaces*, Thai J. Math. **4** (1) (2006), 85–91.
- [19] M. MURSALEEN, S. A. MOHIUDDINE, *On ideal convergence in probabilistic normed spaces*, Math. Slovaca **62** (1) (2012), 49–62.
- [20] M. MURSALEEN, A. ALOTAIBI, *On \mathcal{I} -convergence in random 2-normed spaces*, Math. Slovaca **61** (6) (2011), 933–940.
- [21] S. SARABADAN, S. TALEBI, *Statistical convergence and ideal convergence of sequences of functions in 2-normed spaces*, Internat. J. Math. Math. Sci. **2011** (2011), 10 pages, doi:10.1155/2011/517841.
- [22] E. SAVAŞ, M. GÜRDAL, *Ideal Convergent Function Sequences in Random 2-Normed Spaces*, Filomat, **30** (3) (2016), 557–567.
- [23] I. J. SCHOENBERG, *The integrability of certain functions and related summability methods*, Amer. Math. Monthly **66** (1959), 361–375.
- [24] A. SHARMA, K. KUMAR, *Statistical convergence in probabilistic 2-normed spaces*, Mathematical Sciences, **2** (4) (2008), 373–390.
- [25] A. ŞAHINER, M. GÜRDAL, S. SALTAN, H. GUNAWAN, *Ideal convergence in 2-normed spaces*, Taiwanese J. Math. **11**(5) (2007), 1477–1484.

(Received August 2, 2016)

Sevim Yegül
Afyonkarahisar, Turkey
Afyon Kocatepe University
Afyonkarahisar, Turkey
e-mail: sevimyegull@gmail.com

Erdinç Diındar
Department of Mathematics
Afyon Kocatepe University
Afyonkarahisar, Turkey
e-mail: erdincdundar79@gmail.com,
edundar@aku.edu.tr