

## FURTHER RESULTS ABOUT NORMAL CRITERIA AND SHARED VALUES FOR FAMILIES OF MEROMORPHIC FUNCTIONS

JIANMING QI

*Abstract.* Let  $k$  be a positive integer and let  $\mathcal{F}$  be a family of meromorphic functions in the domain  $D$  all of whose zeros with multiplicity at least  $k$ . Let  $P$  be a polynomial and  $P$  have at least one simple zero,  $p = \deg(P) \geq k + 2$ . If, for each pair  $f, g \in \mathcal{F}$ ,  $P(f)G^m(f)$  and  $P(g)G^m(g)$  share a nonzero constant  $b$  ignoring multiplicity in  $D$ , where  $G(f) = P(f^{(k)}) + H(f)$  is a differential polynomial of  $f$  satisfying  $\frac{w}{\deg}|_H \leq \frac{kmq}{l+mq} + 1$  or  $w(H) - \deg(H) < qk$ , and  $q > l \geq k + 1$  is a positive integer, then  $\mathcal{F}$  is normal in  $D$ .

### 1. Introduction

Let  $D$  be a domain in  $\mathbb{C}$  and  $\mathcal{F}$  is a family of meromorphic in  $D$ .  $\mathcal{F}$  is said to be normal in  $D$ , in the sense of Montel, if each sequence  $\{f_n\} \subset \mathcal{F}$  had a subsequence  $\{f_{n_j}\}$  which converges spherically locally uniformly in  $D$ , to a meromorphic function or  $\infty$  (see [9], [23], [29]). Let  $P(z)$  be a polynomial or a finite complex number.  $\deg P(z)$  denotes the degree of the polynomial  $P(z)$ .

Suppose that  $f(z)$ ,  $g(z)$  are meromorphic functions in  $D$  and  $a \in \mathbb{C} \cup \{\infty\}$ . If  $f(z) = a$  if and only if  $g(z) = a$ , we say that  $f$  and  $g$  share  $a$  ignoring multiplicity (IM) (see [27]).

**DEFINITION 1.** Let  $D \subseteq \mathbb{C}$  be an arbitrary domain,  $m, l_1, l_2, \dots, l_m$  be non-negative integers and  $(0 \leq l_i \leq k)$ , if

$$M(f, f', \dots, f^{(k)}) = a(z) \prod_{i=1}^m f^{(l_i)},$$

where  $f$  is meromorphic and  $a$  is a holomorphic function in  $D$  ( $a \neq 0$ ), then  $M(f, f', \dots, f^{(k)})$  is called a differential monomial of degree  $\deg(M) := m$  and weight  $w(M) := \sum_{i=1}^m (1 + l_i)$ .

The summation  $H := M_1 + \dots + M_n$  of differential monomials  $M_j$  is called the differential polynomial. For the convenience in this article, deviating from the common definition, we definite the degree of  $\deg(H) := \min\{\deg(M_1), \dots, \deg(M_n)\}$  and weight  $w(H) := \max\{w(M_1), \dots, w(M_n)\}$ ,  $\frac{w}{\deg}|_H = \max\left\{\frac{w(M_1)}{\deg(M_1)}, \frac{w(M_2)}{\deg(M_2)}, \dots, \frac{w(M_n)}{\deg(M_n)}\right\}$ .

*Mathematics subject classification* (2010): 30D35, 30D45.

*Keywords and phrases:* Normal family, Nevanlinna theory, sharing values, meromorphic function.

DEFINITION 2. Let  $q$  be a positive integer, and  $b_j(z)$  ( $j = 1, 2, \dots, q-1$ ) be analytic function on a domain  $D$ .

$$P(f^{(k)})(z) = (f^{(k)}(z))^q + b_{q-1}(z)(f^{(k)}(z))^{q-1} + \dots + b_1(z)f^{(k)}(z),$$

$$G(f) = P(f^{(k)}) + H(f, f', \dots, f^{(k)}).$$

In 1959, Hayman [10] proposed:

Conjecture A If  $f$  is a transcendental meromorphic function, then  $f^n f'$  assumes every finite non-zero complex number infinitely often for any positive integer  $n$ .

Hayman [10, 11] himself confirmed it for  $n \geq 3$  and for  $n \geq 2$  in the case of entire  $f$ . Further, it was proved by Mues [17] when  $n \geq 2$ ; Clunie [6] when  $n \geq 1$  and  $f$  is entire; Bergweiler and Eremenko [2] verified the case when  $n = 1$  and  $f$  is of finite order, and finally by Chen and Fang [5] for the case  $n = 1$ .

Correspondingly, there is a conjecture of Hayman [11] related to above problem concerning the normality of  $\mathcal{F}$  (see [1]).

Conjecture B If each  $f \in \mathcal{F}$  satisfies  $f^n f' \neq a$  for a positive integer  $n$  and a finite non-zero complex number  $a$ , then  $\mathcal{F}$  is normal.

Concerning this conjecture, there are many significant results have been obtained by Yang and Zhang [30]; Gu [8]; Oshkin [18]; Li and Xie [15]; Pang [19]; Zalcman [31]. Chen and Fang [5] verified the conjecture B completely. Schick([24]) was the first author to draw a connection between value shared by functions in  $\mathcal{F}$  and the normality of the family  $\mathcal{F}$ . Moreover, many scholars had studied normality criterions such as Meng [3]; Lei and Fang [14]; Li and Gu [16]; Pang and Zalcman [20]; Xia and Xu [25].

In 2004, Fang and Zalcman [7] proved:

Theorem A Let  $\mathcal{F}$  be a family of meromorphic functions in a domain  $D$ , let  $n \geq 1$  be a positive integer, and  $b$  be a finite non-zero complex number. If, for each  $f, g \in \mathcal{F}$ ,  $f$  and  $g$  share 0 IM;  $f^n f'$  and  $g^n g'$  share  $b$  IM in  $D$ , then  $\mathcal{F}$  is normal in  $D$ .

Lately, Zhang [32] obtained:

Theorem B Let  $\mathcal{F}$  be a family of holomorphic functions in a domain  $D$ , let  $n \geq 1$  be a positive integer, and  $b$  be a finite complex number. If, for each  $f, g \in \mathcal{F}$ ,  $f^n(f-1)f'$  and  $g^n(g-1)g'$  share  $b$  IM, then  $\mathcal{F}$  is normal in  $D$ .

In 2008, Zhang [33] improved the condition of Theorem A, and obtained:

Theorem C Let  $\mathcal{F}$  be a family of meromorphic functions in a domain  $D$ , let  $n \geq 2$  be a positive integer, and  $b$  be a finite non-zero complex number. If, for each  $f, g \in \mathcal{F}$ ,  $f^n f'$  and  $g^n g'$  share  $b$  IM, then  $\mathcal{F}$  is normal in  $D$ .

There are examples showing that this result is not true if  $n = 1$ . Recently, Lei and Fang [13] extended Theorems A,B,C and obtained as follows:

**Theorem D.** Let  $\mathcal{F}$  be a family of meromorphic functions in the plane domain  $D$ , let  $P$  be a polynomial with either  $\deg(P) \geq 3$  or  $\deg(P) = 2$  and  $P$  having only one distinct zero. If, for each  $f, g \in \mathcal{F}$ ,  $P(f)f'$  and  $P(g)g'$  share a nonzero constant  $b$  IM in  $D$ , then  $\mathcal{F}$  is normal in  $D$ .

**Theorem E.** Let  $\mathcal{F}$  be a family of meromorphic functions in the domain  $D$ , all of whose poles are multiple, and let  $P$  be a polynomial with two distinct zeros. If, for each  $f, g \in \mathcal{F}$ ,  $P(f)f'$  and  $P(g)g'$  share complex number  $b$  IM in  $D$ , then  $\mathcal{F}$  is normal in  $D$ .

In 2012, Qi, Ding and Yang[22] extended the Theorem D and Theorem E and obtained as follows:

**Theorem D1.** Let  $k$  be a positive integer and let  $\mathcal{F}$  be a family of meromorphic functions in the plane domain  $D$  all of whose zeros with multiplicity at least  $k$ . Let  $P = a_p z^p + \dots + a_2 z^2 + z$  be a polynomial,  $a_p, a_2 \neq 0$  and  $p = \deg(P) \geq k + 2$ . If, for each  $f, g \in \mathcal{F}$ ,  $P(f)G(f)$  and  $P(g)G(g)$  share a non-zero constant  $b$  IM in  $D$ , where  $G(f) = f^{(k)} + H(f)$  be a differential polynomial of  $f$  satisfying  $\frac{w}{\deg} |H| \leq \frac{k}{l+1} + 1$  or  $w(H) - \deg(H) < k$ , then  $\mathcal{F}$  is normal in  $D$ .

**Theorem E1.** Let  $k$  be a positive integer, suppose that  $\mathcal{F}$  be a family of meromorphic functions in the plane domain  $D$  all of whose zeros and poles with multiplicity at least  $k$  and 2 respectively. Let  $P$  be a polynomial with two distinct zeros at least. If, for each  $f, g \in \mathcal{F}$ ,  $P(f)G(f)$  and  $P(g)G(g)$  share a constant  $b$  IM in  $D$ , where  $G(f) = f^{(k)} + H(f)$  be a differential polynomial of  $f$  with  $w(H) - \deg(H) < k$ , then  $\mathcal{F}$  is normal in  $D$ .

In 2009, Hu and Meng[12] obtained the following results.

**Theorem F.** Take positive integers  $n$  and  $k$  with  $n, k \geq 2$  and take a non-zero complex number  $a$ . Let  $\mathcal{F}$  be a family of meromorphic functions in the plane domain  $D$  such that each  $f \in \mathcal{F}$  has only zeros of multiplicity at least  $k$ . For each pair  $(f, g) \in \mathcal{F}$ , if  $f(f^{(k)})^n$  and  $g(g^{(k)})^n$  share  $a$  IM, then  $\mathcal{F}$  is normal in  $D$ .

Also in the final paper of Hu and Meng[12], Hu and Meng pointed that if the form  $f(f^{(k)})^n$  (resp.  $g(g^{(k)})^n$ ) in Theorem 1.1 was replaced by the form  $f^l(f^{(k)})^n$  (resp.  $g^l(g^{(k)})^n$ ) for an integer  $l \geq 2$ , the conclusion also holds.

Based on the ideas of Theorem D and Theorem E, it is natural to ask whether  $f(f^{(k)})^n$ , in Theorem F can be replaced by  $P(f)G(f)^n$ , where  $P(f)$  is polynomial about  $f$ ,  $G(f)$  is a differential polynomial about  $f$ . The main purpose of this paper is to investigate this problem. We prove the following results.

**THEOREM 1. (Main)** *Let  $m, k$  be two positive integers. Let  $\mathcal{F}$  be a family of meromorphic functions in the domain  $D$  all of whose zeros with multiplicity at least  $k$ . Let  $P$  be a polynomial and have at least one simple zero, and  $p = \deg(P) \geq k + 2$ . If, for each pair  $f, g \in \mathcal{F}$ ,  $P(f)G^m(f)$  and  $P(g)G^m(g)$  share a non-zero constant  $b$  IM in  $D$ , where  $G(f) = P(f^{(k)}) + H(f)$  be a differential polynomial of  $f$  satisfying*

$\frac{w}{\deg} |H \leq \frac{kmq}{l+mq} + 1$  or  $w(H) - \deg(H) < qk$ , and  $q > l \geq k+1$  is a positive integer, then  $\mathcal{F}$  is normal in  $D$ .

REMARK 1. If the polynomial  $P(z)$  has only one zero, the Theorem 1 is established for  $\deg(P) \geq k+1$ .

COROLLARY 1. Let  $k$  be two positive integers and  $\mathcal{F}$  be a family of meromorphic functions in the plane domain  $D$  all of whose zeros with multiplicity at least  $k$ . Let  $P$  be a polynomial as theorem 1. If, for each  $f, g \in \mathcal{F}$ ,  $P(f)(f^{(k)})^m$  and  $P(g)(g^{(k)})^m$  share a non-zero constant  $b$  IM in  $D$ , then  $\mathcal{F}$  is normal in  $D$ .

THEOREM 2. (Main) Let  $k$  be a positive integer, suppose that  $\mathcal{F}$  be a family of meromorphic functions in the domain  $D$  all of whose zeros and poles with multiplicity at least  $k$  and 2 respectively. Let  $P$  be a polynomial with two distinct zeros at least. If, for each  $f, g \in \mathcal{F}$ ,  $P(f)G^m(f)$  and  $P(g)G^m(g)$  share a constant  $b$  IM in  $D$ , where  $G(f) = P(f^{(k)}) + H(f)$  be a differential polynomial of  $f$  with  $w(H) - \deg(H) < qk$ , then  $\mathcal{F}$  is normal in  $D$ .

COROLLARY 2. Let  $k$  be a positive integer, suppose that  $\mathcal{F}$  be a family of meromorphic functions in the plane domain  $D$  all of whose zeros and poles with multiplicity at least  $k$  and 2 respectively. Let  $P$  be a polynomial as theorem 2. If, for each  $f, g \in \mathcal{F}$ ,  $P(f)(f^{(k)})^m$  and  $P(g)(g^{(k)})^m$  share  $b$  IM in  $D$ , then  $\mathcal{F}$  is normal in  $D$ .

Example 1. Let  $D = \{|z| < 1\}$  and take a non-zero complex number  $a$ . Fix three positive integers  $n \geq 2, k, m$ .

$$\mathcal{F} = \{f_d(z) = dz^{k-1} | d = 1, 2, \dots\}.$$

Obviously, for distinct positive integers,  $i, j$ , we have  $f_i^n(f_i^{(k)})^m$  and  $f_j^n(f_j^{(k)})^m$  share  $a$  IM. However, the family  $\mathcal{F}$  is not normal at  $z = 0$ .

Example 2. Set

$$\mathcal{F} = \{f_d(z) = dz - \frac{d}{3} + \frac{a}{d^m} | d = 1, 2, \dots\}.$$

For distinct positive integers,  $i, j$ , we have  $f_i(f_i')^m$  and  $g_j(g_j')^m$  share  $a$  IM. However, the family  $\mathcal{F}$  is not normal at  $z = \frac{1}{3}$ .

Example 3. Set

$$\mathcal{F} = \{e^{dz} | d = 1, 2, \dots\}.$$

Obviously, any  $f_d \in \text{mathcal{F}}$  has only zeros of multiplicity at least  $k$ . For distinct positive integers  $i, j$ , we have  $f_i^n(f_i^{(k)})^m$  and  $f_j^n(f_j^{(k)})^m$  share 0 IM. However, the families are not normal at  $z = 0$ .

REMARK 2. Example 1 shows that the condition that  $f$  has only zeros of multiplicity at least  $k$  is sharp in Theorem 1 and 2. Example 2 shows the conditions  $p = \deg(P) \geq k + 2$  in Theorem 1 and  $P$  is a polynomial with two distinct zeros at least in Theorem 2 seems not be omitted. Example 3 shows the condition  $a \neq 0$  in Theorem 1 is necessary.

REMARK 3. Obviously, Theorem 1 and Theorem 2 both extend the Theorem D1 and Theorem D2.

## 2. Some lemmas

In order to prove our theorem, we need the following Lemmas:

Lemma 2.1 [Pang-Zalcman’s lemma](see [4], [21], [31]) Let  $k$  be a positive integer, let  $\mathcal{F}$  be a family of meromorphic functions in the unit disc  $\Delta$  with the property that for each  $f \in \mathcal{F}$ , all zeros of multiplicity at least  $k$ . Suppose that there exists a number  $A \geq 1$  such that  $|f^{(k)}(z)| \leq A$  whenever  $f = 0$ . Suppose that  $\mathcal{F}$  is not normal at  $z_0$ , then for  $0 \leq \alpha \leq k$ , there exist

- a). points  $z_n \in \Delta$ ,  $z_n \rightarrow z_0$ ;
- b). functions  $f_n \in \mathcal{F}$ ;
- c). positive numbers  $\rho_n \rightarrow 0^+$ ;

such that  $\rho_n^{-\alpha} f_n(z_n + \rho_n \xi) = g_n(\xi) \rightarrow g(\xi)$  locally uniformly with respect to the spherical metric, where  $g(\xi)$  is a nonconstant meromorphic function on  $\mathbb{C}$ , all of whose zeros have multiplicity at least  $k$ , such that  $g^\sharp(\xi) \leq g^\sharp(0) = kA + 1$ . In particular,  $g$  has order at most 2.

Lemma 2.2([28]) Let  $n \geq 2$ ,  $k$  be a positive integer and  $n_k$  be a positive integer. If  $f$  is a transcendental meromorphic function, then  $f^n(f^{(k)})^{n_k}$  assume every finite non-zero complex value infinitely often.

Lemma 2.3 ([26]) Let  $f$  be a non-constant rational function and  $n, m$  be two positive integers. let  $k$  be a positive integer, and let  $b$  be a non-zero finite complex number. then  $f^n(f^{(k)})^m - b$  has one zero at least.

Lemma 2.4 Let  $f$  be a non-constant rational function in the plane  $\mathbb{C}$  and  $n \geq 2, m$  be two positive integers. let  $k$  be a positive integer, and let  $b$  be a non-zero finite complex number. then  $f^n(f^{(k)})^m - b$  has two distinct zeros at least.

*Proof.* By Lemma 2.3,  $f^n(f^{(k)})^m - b$  has one zero at least. On the contrary that  $f^n(f^{(k)})^m - b$  has one zero at most.

Suppose that  $f$  is a non-constant polynomial, we have

$$f^n(f^{(k)})^m(z) = A(z - z_0)^l + b, \tag{2.1}$$

where  $A$  is a non-zero constant and  $l \geq 2$  is a positive integer. The right hand side of (2.1) has only simple zeros, but the left has multiple zeros, a contradiction. Thus  $f$  is a non-polynomial rational function.

Set

$$f^n(f^{(k)})^m(z) = A \frac{(z - \alpha_1)^{m_1} (z - \alpha_2)^{m_2} \cdots (z - \alpha_s)^{m_s}}{(z - \beta_1)^{n_1} (z - \beta_2)^{n_2} \cdots (z - \beta_t)^{n_t}} = \frac{P}{Q}, \quad (2.2)$$

where  $A$  is a non-zero constant,  $P$  and  $Q$  are polynomials of degree  $M$  and  $N$  respectively. Also  $P$  and  $Q$  have no common factor. By the zeros of  $f$  are at least  $k$ , we obtain  $m_i \geq 2 (i = 1, 2, \dots, s)$ ,  $n_j \geq 2 (j = 1, 2, \dots, t)$ . Hence

$$m_1 + m_2 + \cdots + m_s = M \geq 2s, \quad (2.3)$$

$$n_1 + n_2 + \cdots + n_t = N \geq 2t. \quad (2.4)$$

From (2.2), we obtain

$$(f^n(f^{(k)})^m(z))' = \frac{(z - \alpha_1)^{m_1-1} (z - \alpha_2)^{m_2-1} \cdots (z - \alpha_s)^{m_s-1} g_1(z)}{(z - \beta_1)^{n_1+1} (z - \beta_2)^{n_2+1} \cdots (z - \beta_t)^{n_t+1}}, \quad (2.5)$$

where  $g_1$  is a polynomial of degree at most  $s + t - 1$ .

Since  $f^n(f^{(k)})^m - b$  has only one zero. So we may set

$$f^n(f^{(k)})^m(z) = b + \frac{B(z - z_0)^l}{(z - \beta_1)^{n_1} (z - \beta_2)^{n_2} \cdots (z - \beta_t)^{n_t}} = \frac{P}{Q}. \quad (2.6)$$

Note that  $b \neq 0$ , we obtain  $z_0 \neq \alpha_i (i = 1, \dots, s)$ , where  $B$  is a non-zero constant, and  $l \geq 2$  is a positive integer.

From (2.6), we obtain

$$[f^n(f^{(k)})^m(z)]' = \frac{(z - z_0)^{l-1} g_2(z)}{(z - \beta_1)^{n_1+1} \cdots (z - \beta_t)^{n_t+1}}. \quad (2.7)$$

Where  $g_2(z) = B(l - N)z^l + B_1 z^{l-1} \cdots + B_t$  is a polynomial ( $B_1, \dots, B_t$  are constants).

Case 1. If  $l \neq N$ , by (2.6), then we obtain the  $\deg(P) \geq \deg(Q)$ . So  $M \geq N$ . By (2.5) and (2.7), we obtain  $\sum_{i=1}^s (m_i - 1) \leq \deg(g_2) = t$ . So  $M - s - \deg(g_1) \leq t$ , and  $M \leq s + t + \deg(g_1) \leq 2(s + t) - 1 < 2(s + t)$ . By (2.3) and (2.4), we obtain

$$M < 2(s + t) \leq 2\left[\frac{M}{2} + \frac{N}{2}\right] \leq M.$$

So we deduce that  $M < M$ . This is impossible.

Case 2. If  $l = N$ , then we consider two subcases.

Case 2.1. If  $M \geq N$ , by (2.5) and (2.7), we obtain  $\sum_{i=1}^s (M_i - 1) \leq \deg(g_2) = t$ . So  $M - s - \deg(g_1) \leq t$ , and  $M \leq s + t + \deg(g_1) \leq 2(s + t) - k < 2(s + t)$ , by the same reasoning mentioned in the case 1. This is impossible.

Case 2.2. If  $M < N$ , by (2.5) and (2.7), we obtain  $l - 1 \leq \deg g_1 \leq s + t - 1$ , then

$$\begin{aligned} N = l &\leq \deg(g_1) + 1 \leq s + t - 1 + 1 < s + t \\ &\leq 2\left[\frac{M}{2} + \frac{N}{2}\right] \leq N. \end{aligned}$$

So we deduce that  $N < N$ . It is impossible.  $\square$

Lemma 2.5([27]) Let  $f(z)$  be a non-constant rational function, then  $f(z)$  has only one deficient value.

### 3. Proof of Theorem 2

*Proof.* Without loss of generality, we assume that  $P(z) = Q(z)z(z - 1)$ , where  $Q(z) \not\equiv 0$  is a polynomial. Suppose that  $\mathcal{F}$  is not normal in  $D$ . Then there exists at least one  $z_0$  such that  $\mathcal{F}$  is not normal at  $z_0$ , we assume that  $z_0 = 0$ . By Lemma 2.1, there exist points  $z_j \rightarrow 0$ ; a sequence  $\rho_j \rightarrow 0^+$  and functions  $f_j \in \mathcal{F}$  such that

$$g_j(\xi) = f_j(z_j + \rho_j \xi) \rightarrow g(\xi), \tag{3.1}$$

locally uniformly with respect to the spherical metric, where  $g$  is a non-constant meromorphic function in  $\mathbb{C}$ , all of whose zeros and poles are of multiplicity at least  $k$  and 2 respectively.

If  $Q(g)g(g - 1)(g^{(k)})^{qm} \equiv 0$ , then  $g$  is a constant, a contradiction.

If  $Q(g)g(g - 1)(g^{(k)})^{qm} \neq 0$ . Because of the zeros of  $g$  have multiplicity at least  $k$ , we obtain  $g \neq 0, 1$ .

We can claim that  $g$  is not a transcendental meromorphic function. In fact, if it is not true, we have

$$\begin{aligned} T(r, g) &\leq \bar{N}(r, g) + \bar{N}\left(r, \frac{1}{g}\right) + \bar{N}\left(r, \frac{1}{g - 1}\right) + S(r, g) \\ &\leq \frac{1}{2}N(r, g) + S(r, g) \\ &\leq \frac{1}{2}T(r, g) + S(r, g) \end{aligned}$$

Thus  $T(r, g) = S(r, g)$ , a contradiction. So  $g$  is a rational function. Since  $g \neq 0, 1$ , then  $g$  is a constant, a contradiction.

Thus  $Q(g)g(g-1)(g^{(k)})^{qm}$  is a non-constant meromorphic function and has one zero at least.

Next we will prove that  $Q(g)g(g-1)(g^{(k)})^{qm}$  has just a single zero. In fact, let  $\xi_0$  and  $\xi_0^*$  be two distinct solutions of  $Q(g)g(g-1)(g^{(k)})^{qm}$ . We choose a positive number  $\delta$  small enough such that  $g$  and  $g_j$  are holomorphic in  $D(\xi_0, \delta_1)D(\xi_0^*, \delta_1)$  and  $D(\xi_0, \delta_1) \cap D(\xi_0^*, \delta_1) = \emptyset$ .

From (3.1), we have

$$\begin{aligned} & \rho_j^k [Q(f_j(z_j + \rho_j \xi) f_j(z_j + \rho_j \xi) (f_j(z_j + \rho_j \xi) - 1) \cdot G^m(f_j(z_j + \rho_j \xi) - b)] \\ = & \rho_j^{mqk} [Q(g_j(\xi) g_j(\xi) (g_j(\xi) - 1) \cdot (\rho_j^{-qk} (g_j^{(k)}(\xi))^q + \cdots + \rho_j^{-k} b_1(z_j + \rho_j \xi) g_j^{(k)}(\xi)) \\ & + \sum_{j=1}^n a(z_j + \rho_j \xi) \rho_j^{\deg(M_j) - w(M_j)} M_i(g, g', \dots, g^{(k)}) - b]^{qm} \\ \rightarrow & Q(g(\xi)) g(\xi) (g(\xi) - 1) (g^{(k)}(\xi))^{qm}. \end{aligned} \quad (3.2)$$

By Hurwitz's theorem, there exist points  $\xi_j \in D(\xi_0, \delta)$ ,  $\xi_j^* \in D(\xi_0^*, \delta)$  such that for sufficiently large  $j$

$$\begin{aligned} Q(f_j(z_j + \rho_j \xi_j)) f_j(z_j + \rho_j \xi_j) (f_j(z_j + \rho_j \xi_j) - 1) G^m(f_j(z_j + \rho_j \xi_j)) &= b, \\ Q(f_j(z_j + \rho_j \xi_j^*)) f_j(z_j + \rho_j \xi_j^*) (f_j(z_j + \rho_j \xi_j^*) - 1) G^m(f_j(z_j + \rho_j \xi_j^*)) &= b. \end{aligned}$$

By the hypothesis that for each pair of functions  $f$  and  $g$  in  $\mathcal{F}$ ,  $P(f)G^m(f)$  and  $P(g)G^m(g)$  share 0 in  $D$ , we know that for any positive integer  $d$

$$\begin{aligned} Q(f_d(z_j + \rho_j \xi_j)) f_d(z_j + \rho_j \xi_j) (f_d(z_j + \rho_j \xi_j) - 1) G^m(f_d(z_j + \rho_j \xi_j)) &= b, \\ Q(f_d(z_j + \rho_j \xi_j^*)) f_d(z_j + \rho_j \xi_j^*) (f_d(z_j + \rho_j \xi_j^*) - 1) G^m(f_d(z_j + \rho_j \xi_j^*)) &= b. \end{aligned}$$

Fix  $d$ , take  $j \rightarrow \infty$ , and note  $z_j + \rho_j \xi_j \rightarrow 0$ ,  $z_j + \rho_j \xi_j^* \rightarrow 0$ , then

$$Q(f_d(0)) f_d(0) (f_d(0) - 1) G^m(f_d(0)) = b.$$

Since the zeros of  $P(f_d)G^m(f_d) - b$  has no accumulation point, so  $z_j + \rho_j \xi_j = 0$ ,  $z_j + \rho_j \xi_j^* = 0$ .

Hence

$$\xi_j = -\frac{z_j}{\rho_j}, \quad \xi_j^* = -\frac{z_j}{\rho_j}.$$

This contradicts with  $\xi_j \in D(\xi_0, \delta)$ ,  $\xi_j^* \in D(\xi_0^*, \delta)$  and  $D(\xi_0, \delta) \cap D(\xi_0^*, \delta) = \emptyset$ . So  $Q(g)g(g-1)(g^{(k)})^{qm}$  has just a single zero, which can be denoted by  $\xi_0$ .

Suppose that  $g$  is a transcendental meromorphic function. Since  $Q(g)g(g-1)(g^{(k)})^{qm}$  has only one zero, so  $g = 0$  and  $g = 1$  has only finite zeros. As the above argument, we obtain  $T(r, g) = S(r, g)$ , a contradiction.

Thus  $g$  is a rational function which is not a polynomial. Because  $Q(g)g(g-1)(g^{(k)})^{qm}$  has only one zero, we have  $g \neq 0$  or  $g \neq 1$ .



If  $g \neq 0$ , then  $g(\xi) = \frac{1}{H(\xi)}$  where  $H(\xi)$  is a non-constant polynomial. Since  $g(\xi) - 1 = \frac{1-H(\xi)}{H(\xi)}$  has just a single zero, so

$$1 - H(\xi) = A(\xi - B)^k, \tag{3.3}$$

where  $A \neq 0, B$  are constant,  $k \geq 2$  is a positive integer.

We claim  $H(\xi)$  has only simple zeros. Suppose, on the contrary, that  $H(z_0) = 0$  and  $z_0$  is multiple. Form (3.3), we arrive at  $0 = H'(z_0) = (1 - H(z_0))' = Ak(z_0 - B)^{(k-1)}$ , a contradiction, since  $z_0 \neq B$ .

Thus  $H(\xi)$  has just simple zeros, this contradicts that  $g$  has no simple pole. If  $g \neq 1$ , we can argue it in the same way. So  $\mathcal{F}$  is normal on  $D$ .  $\square$

### 4. Proof of Theorem 1

*Proof.* We may assume that  $D = \{|z| < 1\}$ . Suppose that  $\mathcal{F}$  is not normal in  $D$ . Without loss of generality, we assume that  $\mathcal{F}$  is not normal at  $z_0 = 0$ . Then, by Lemma 2.1, there exist a sequence  $z_j$  of complex numbers with  $z_j \rightarrow 0$  ( $j \rightarrow \infty$ ); a sequence  $f_j$  of  $\mathcal{F}$ ; and a sequence  $\rho_j \rightarrow 0^+$  such that

$$g_j(\xi) = \rho_j^{-\frac{mkq}{l+mq}} f_j(z_j + \rho_j \xi) \tag{3.4}$$

converges uniformly to a non-constant meromorphic functions  $g(\xi)$  in  $C$  with respect to the spherical metric. Moreover,  $g(\xi)$  is of order at most 2. Where  $q > l (\geq k + 1)$  is a constant. By Hurwitz's theorem, the zeros of  $g(\xi)$  have at least multiplicity  $k$ . Next we will distinguish two cases:

Case 1. When  $P(z)$  has two distinct zeros, then we can denote  $P(f) = f^l(f + 1)$  ( $q > l \geq k + 1$ ).

If  $g^l(g^{(k)})^{qm} \equiv b$ , then  $g$  has no zeros. Of course,  $g$  also has no poles. Since  $g$  is a non-constant meromorphic function of order at most 2, we obtain  $g(\xi) = e^{d\xi^2+h\xi+c}$  (where  $D, h, c$  are constants and  $dh \neq 0$ ). At this moment  $g^l(g^{(k)})^{qm} \neq b$ . Which is a contradiction.

If  $g^l(g^{(k)})^{qm} \neq b$ , then by Lemma 2.2, we obtain that  $g$  is a constant. This contradicts that  $g$  is a non-zero meromorphic function.

Thus  $g^l(g^{(k)})^{qm} - b$  is a non-constant meromorphic function and has one zero at least.

Next we will prove that  $g^l(g^{(k)})^{qm} - b$  has just a single zero. In fact, let  $\xi_0$  and  $\xi_0^*$  be two distinct solutions of  $g^l(g^{(k)})^{qm} - b$ . We choose a positive number  $\delta_1$  small enough such that  $g$  and  $g_j$  are holomorphic in  $\xi_j \in D(\xi_0, \delta), \xi_j^* \in D(\xi_0^*, \delta)$ .

From (3.4), we have

$$\begin{aligned} & [f_j^{l+1}(z_j + \rho_j \xi) + f_j^l(z_j + \rho_j \xi)] \cdot [P(f_j^{(k)}(z_j + \rho_j \xi)) + H(f, f', \dots, f^{(k)})]^m - b \\ = & [\rho_j^{-\frac{qk}{l+mq}} (g_j^{(k)}(\xi))^q + \dots + \sum_{j=1}^n a_j^* \rho_j^{\frac{kmq}{l+mq} + 1} \deg(M_j) - w(M_j) M_i(g, g', \dots, g^{(k)})]^m, \\ & [\rho_j^{\frac{kmq(l+1)}{l+mq}} g_j^{l+1}(\xi) + \rho_j^{\frac{mqk}{l+mq}} g_j^l(\xi)] - b \rightarrow g^l(\xi)(g^{(k)}(\xi))^{qm} - b. \end{aligned} \quad (3.5)$$

Choose  $\delta_2$  such that  $D(\xi_0, \delta_2) \cap D(\xi_0^*, \delta_2) = \emptyset$  and such that  $g^l(g^{(k)})^{qm} - b$  has no other zeros in  $D(\xi_0, \delta) \cup D(\xi_0^*, \delta)$ . By Hurwitz's theorem, there exist points  $\xi_j \in D(\xi_0, \delta)$ ,  $\xi_j^* \in D(\xi_0^*, \delta)$  such that for sufficiently large  $j$

$$\begin{aligned} & [f_j^{l+1}(z_j + \rho_j \xi_j) + f_j^l(z_j + \rho_j \xi_j)] G^m(f_j(z_j + \rho_j \xi_j)) - b = 0, \\ & [f_j^{l+1}(z_j + \rho_j \xi_j^*) + f_j^l(z_j + \rho_j \xi_j^*)] G^m(f_j(z_j + \rho_j \xi_j^*)) - b = 0. \end{aligned}$$

By the hypothesis that for each pair of functions  $f$  and  $g$  in  $\mathcal{F}$ ,  $P(f)G^m(f^{(k)})$  and  $P(g)G^m(g^{(k)})$  share  $b$  in  $D$ , we know that for any positive integer  $d$

$$\begin{aligned} & [f_d^{l+1}(z_j + \rho_j \xi_j) + f_d^l(z_j + \rho_j \xi_j)] G^m(f_d(z_j + \rho_j \xi_j)) - b = 0, \\ & [f_d^{l+1}(z_j + \rho_j \xi_j^*) + f_d^l(z_j + \rho_j \xi_j^*)] G^m(f_d(z_j + \rho_j \xi_j^*)) - b = 0. \end{aligned}$$

Fix  $d$ , take  $j \rightarrow \infty$ , and note  $z_j + \rho_j \xi_j \rightarrow 0$ ,  $z_j + \rho_j \xi_j^* \rightarrow 0$ , then

$$[f_d^{l+1}(0) + f_d^l(0)] G^m(f_d(0)) - b = 0.$$

Since the zeros of  $P(f)G^m(f) - b$  has no accumulation point, so  $z_j + \rho_j \xi_j = 0$ ,  $z_j + \rho_j \xi_j^* = 0$ .

Hence

$$\xi_j = -\frac{z_j}{\rho_j}, \quad \xi_j^* = -\frac{z_j}{\rho_j}.$$

This contradicts with  $\xi_j \in D(\xi_0, \delta)$ ,  $\xi_j^* \in D(\xi_0^*, \delta)$  and  $D(\xi_0, \delta) \cap D(\xi_0^*, \delta) = \emptyset$ . So  $g^l(g^{(k)})^{qm} - b$  has just a single zero, which can be denoted by  $\xi_0$ .

From the above, we know  $g^l(g^{(k)})^{qm} - b$  has just a unique zero. This contradicts Lemma 2.2 and Lemma 2.4.

Case 2. If  $P(z)$  has more than three distinct zeros, we can denote  $P(z) = Q(g)g(g-1)(g-a)$ .

Without loss of generality, we assume that  $\mathcal{F}$  is not normal at  $z_0 = 0$ . Then, by Lemma 2.1, there exist a sequence  $z_j$  of complex numbers with  $z_j \rightarrow 0$  ( $j \rightarrow \infty$ ); a sequence  $f_j$  of  $\mathcal{F}$ ; and a sequence  $\rho_j \rightarrow 0^+$  such that

$$g_j(\xi) = f_j(z_j + \rho_j \xi) \quad (3.6)$$

converges uniformly to a non-constant meromorphic functions  $g(\xi)$  in  $C$  with respect to the spherical metric.

Proceeding as in the proof of Theorem 2, we have  $Q(g)g(g-1)(g-a)G^m(g)$  only one zero.

Obviously,  $g$  is not a transcendental meromorphic function from Picard Theorem. Thus  $g$  is a non-constant rational function and  $g$  doesn't assume two complex number of  $\{0, 1, a\}$ , due to Lemma 2.5, a contradiction. So  $\mathcal{F}$  is normal in  $z_0$ .  $\square$

*Acknowledgement.* This work was supported by Applied Mathematical Academic Discipline Project of Shanghai Dianji University(16JCXK02), and Humanity and Social Science Youth foundation of Ministry of Education(18YJC630120).

## REFERENCES

- [1] W. BERGWELER, *Bloch's principle*, Comput Methods Funct. Theory., **6**, (2006), 77–108.
- [2] W. BERGWELER AND A. EREMENKO, *Complex dynamics and value distribution*, In International Conference of Complex Analysis, Nanjing, 1994.
- [3] M. CHAO, *Normal families and shared values of meromorphic functions*, Bull. Malays. Math. Sci. Soc., **2**, 31(2008), 1 85–90.
- [4] H. H. CHEN AND Y. X. GU, *Improvement of Maty's criterion and its application*, Sci China. Ser A, 36(1993) 647–681.
- [5] H. H. CHEN AND M. L. FANG, *On the value distribution of  $f^n f'$* , Sci China. Ser A, 38(1995), 789–798.
- [6] J. CLUNIE, *On a result of Hayman*, J. London Math Soc. 42(1967), 389–392.
- [7] M. L. FANG AND L. ZALCMAN, *A note on normality and shared values*, J. Aust. Math. Soc., 76(2004), 141–150.
- [8] Y. X. GU, *On normal families of meromorphic functions*, Sci China. Ser A, 36(1978), 373–384.
- [9] W. K. HAYMAN, *Meromorphic functions*, Clarendon Press, Oxford, 1964.
- [10] W. K. HAYMAN, *Picard value of meromorphic functions and their derivatives*, Ann. of Math. 70(1959), 9–42.
- [11] W. K. HAYMAN, *Research problem in function theory*, Athlone Press, University of London, 1967.
- [12] P. C. HU AND D. W. MENG, *Normality criteria of meromorphic functions with multiple zeros*, J. Math. Anal. Appl., 357(2009), 323–329.
- [13] C. L. LEI AND M. L. FANG, *Normality and shared values concerning differential polynomials*, Sci China. Ser A., 53(2010), 749–754.
- [14] L. C. LEI, M. L. FANG AND D. G. YANG, *Normal families and shared values of meromorphic functions*, Proc. Japan Acad.(Ser. A), 83(2007), 36–39.
- [15] S. Y. LI AND H. C. XIE, *On normal families of meromorphic functions*, Acta Math. Sinica, 29(1986), 468–476.
- [16] Y. T. LI AND Y. X. GU, *On normal families of meromorphic functions*, J. Math. Anal. Appl. 354(2009), 421–425.
- [17] E. MUES, *Über ein Problem von Hayman*, Math. Z. 164(1979), 239–259.
- [18] I. B. OSHKIN, *On a test for the normality of families of holomorphic functions*, Uspekhi Mat. Nauk, 37(2)(1982), 221–222.
- [19] X. C. PANG, *Bloch principle and normal criterion*, Sci China Ser A, 32(1989), 782–791.
- [20] X. C. PANG AND L. ZALCMAN, *Normal families and shared values*, Acta Math. 76(2000), 171–182.
- [21] X. C. PANG AND L. ZALCMAN, *Normal families and shared values*, Bull London Math Soc. 32(2000), 325–331.
- [22] J. M. QI, J. DING AND L. Z. YANG, *Normality Criteria for Families of Meromorphic Function Concerning Shared Values*, Bull. Malays. Math. Sci. Soc. **2**, 35 (2012), 449–457.
- [23] J. SCHIFF, *Normal Families*, Springer-Verlag, Berlin, 1993.
- [24] W. SCHWICK, *Sharing values and normality*, Arch. Math. 59(1992), 50–54.

- [25] J. Y. XIA AND Y. XU, *Normality criterion concerning sharing functions*, Bull. Malays. Math. Sci. Soc. **2**, 33(2010), 479–486.
- [26] J. F. XU AND W. S. CAO, *Some normality criteria of meromorphic functions*, J., Inequ., Appl., 2010, articles ID 926302, 10 pages, doi:10.1155/2010/926302.
- [27] C. C. YANG AND H. X. YI, *Uniqueness Theory of Meromorphic Functions*, Kluwer Academic Publishers 2003.
- [28] C. C. YANG AND P. C. HU, *On the value distribution of  $ff^{(k)}$* , Kodai Mathematical Journal, **2**, 19(1996), 157–167.
- [29] L. YANG, *Value distribution theory*, Springer-Verlag, Berlin, 1993.
- [30] L. YANG AND G. H. ZHANG, *Recherches sur la normalité des familles de fonction analytiques à des valeurs multiples, I. Un nouveau critère et quelques applications*, Sci China. Ser A, 14(1965), 1258-1271; II. Généralizations, Ibid. 15(1966), 433–453.
- [31] L. ZALCMAN, *Normal families New perspectives*, Bull. Amer. Math. Soc. 35(1998), 215–230.
- [32] Q. C. ZHANG, *On normal criterion of holomorphic functions*, Math. Practice and Theory. 36(2006), 283–286.
- [33] Q. C. ZHANG, *Some normality criteria of meromorphic functions*, Complex Var. Elliptic Equ. 53(2008), 791–795.
- [34] Z. L. ZHANG AND W. LI, *Picard exceptional values for two class differential polynomials*, Acta. Math. Sinica. 34(1994), 828–835.

(Received January 22, 2019)

*Jianming Qi*  
*School of Business*  
*Shanghai Dianji University*  
*Shanghai, 201306, P.R. China,*  
*e-mail: qijianmingsdju@163.com*