

GENERALIZED INTEGRAL INEQUALITIES FOR CONVEX FUNCTIONS

M. EMIN ÖZDEMİR AND ALPER EKINCI

(Communicated by S. Varošanec)

Abstract. In this paper, we prove some general inequalities for convex functions and give Ostrowski, Hadamard and Simpson type results for a special case of these inequalities.

1. Introduction

The function $f : [a, b] \rightarrow \mathbb{R}$, is said to be convex, if we have

$$f(tx + (1-t)y) \leq tf(x) + (1-t)f(y)$$

for all $x, y \in [a, b]$ and $t \in [0, 1]$. For more information see the papers [3], [1], [2].

Let $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a convex function and let $a, b \in I$, with $a < b$. The following double inequality:

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a)+f(b)}{2}$$

is known in the literature as Hadamard's inequality.

In 1928 Ostrowski proved the following famous inequality:

THEOREM 1. *Let $f : [a, b] \rightarrow \mathbb{R}$ be continuous on $[a, b]$ and differentiable on (a, b) and its derivative $f' : (a, b) \rightarrow \mathbb{R}$ be bounded on (a, b) , that is, $\|f'\|_\infty := \sup_{t \in (a, b)} |f'(t)| < \infty$. Then for any $x \in [a, b]$, the following inequality holds:*

$$\left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \left| \frac{1}{4} + \frac{(x - \frac{a+b}{2})^2}{(b-a)^2} \right| (b-a) \|f'\|_\infty.$$

The inequality is sharp in the sense that the constant $1/4$ cannot be replaced by a smaller one. In the rest of this section we list known results which we will generalize in the following section.

Sarikaya et al. obtained following Simpson type inequalities in [5].

Mathematics subject classification (2010): 26D15, 26D10.

Keywords and phrases: Convex functions, Hermite-Hadamard inequality, Simpson's inequality, power-mean inequality, Ostrowski's inequality; Hölder's inequality.

THEOREM 2. Let $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a differential mapping on I^o such that $f' \in L_1[a, b]$, where $a, b \in I$ with $a < b$. If $|f'|$ is convex on $[a, b]$, then the following inequality holds:

$$\left| \frac{1}{6} \left[f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right] - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq \frac{5(b-a)}{72} [|f'(a)| + |f'(b)|]. \quad (1)$$

THEOREM 3. Let $f : I^o \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a differential mapping on I^o such that $f' \in L_1[a, b]$, where $a, b \in I$ with $a < b$. If $|f'|^q$ is convex on $[a, b]$, $q \geq 1$, then the following inequality holds:

$$\begin{aligned} & \left| \frac{1}{6} \left[f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right] - \frac{1}{b-a} \int_a^b f(u) du \right| \quad (2) \\ & \leq \frac{(b-a)}{72} (5)^{1-\frac{1}{q}} \left\{ \left[\frac{61|f'(b)|^q + 29|f'(a)|^q}{18} \right]^{\frac{1}{q}} \right. \\ & \quad \left. + \left[\frac{61|f'(a)|^q + 29|f'(b)|^q}{18} \right]^{\frac{1}{q}} \right\}. \end{aligned}$$

Estimation for the difference between the middle and the leftmost term in the Hadamard inequality was proved by Kirmaci in [8].

THEOREM 4. Let $f : I^o \rightarrow \mathbb{R}$ be a differential mapping on I^o , $a, b \in I^o$ with $a < b$. If $|f'|$ is convex on $[a, b]$, then we have

$$\left| \frac{1}{b-a} \int_a^b f(x) dx - f\left(\frac{a+b}{2}\right) \right| \leq \frac{b-a}{4} \left(\frac{|f'(a)| + |f'(b)|}{2} \right). \quad (3)$$

Similarly, bound for the difference between the middle and the right most term in the Hadamard inequality was considered by Dragomir and Agarwal in [7] and has the following forms.

THEOREM 5. Let $f : I^o \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a differential mapping on I^o , $a, b \in I^o$ with $a < b$. If $|f'|$ is convex on $[a, b]$, then the following inequality holds:

$$\begin{aligned} & \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \quad (4) \\ & \leq \frac{(b-a)}{8} [|f'(a)| + |f'(b)|]. \end{aligned}$$

THEOREM 6. *Let $f : I^o \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a differential mapping on I^o , $a, b \in I^o$ with $a < b$ and let $p > 1$. If the new mapping $|f'|^{p/(p-1)}$ is convex on $[a, b]$, then the following inequality holds:*

$$\begin{aligned} & \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \tag{5} \\ & \leq \frac{(b-a)}{2(p+1)^{1/p}} \left[\frac{|f'(a)|^{p/p-1} + |f'(b)|^{p/p-1}}{2} \right]^{(p-1)/p} . \end{aligned}$$

In this paper we give inequalities involving m harmonic polynomials and a function f such that $|f^{(n)}|^q$ is convex for some $q \geq 1$ and $n \in \mathbb{N}$. After each general inequality we obtain a result related to particular case $n = 1, m = 2$ and point out that for some especial values of our variables h and c , these results become the known results given in the above text.

2. Main results

In the further text m and n are fixed integers, $\sigma := \{a = x_0 < x_1 < x_2 < \dots < x_m = b\}$ is a division of an interval $[a, b]$ with $m + 1$ nodes, $\sigma' := \{0 = s_0 < s_1 < \dots < s_m = 1\}$ is a corresponding division of the interval $[0, 1]$ connected with σ by relation $s_j = \frac{x_j - a}{b - a}$. Let $\{P_{jk}\}_{k \in \mathbb{N}}, j = 1, \dots, m$ be harmonic sequences of polynomials, i.e. $P'_{jk} = P_{j,k-1}, k \in \mathbb{N}, P_{j0} = 1, j = 1, \dots, m$. Let us define a kernel S_n as

$$S_n(t, \sigma) = \begin{cases} P_{1n}(t), & t \in [a, x_1] \\ P_{2n}(t), & t \in (x_1, x_2) \\ \vdots \\ P_{mn}(t), & t \in (x_{m-1}, b]. \end{cases}$$

In paper [4] Pečarić and Varošanec gave the following identity for n -times differentiable function f on $[a, b]$:

$$\begin{aligned} (-1)^n \int_a^b S_n(x, \sigma) f^{(n)}(x) dx &= \int_a^b f(t) dt + \sum_{k=1}^n (-1)^k [P_{mk}(b) f^{(k-1)}(b) \tag{6} \\ &+ \sum_{j=1}^{m-1} [P_{jk}(x_j) - P_{j+1,k}(x_j)] f^{(k-1)}(x_j) - P_{1k}(a) f^{(k-1)}(a)]. \end{aligned}$$

For further applications of this identity see [4], [6].

Let us denote the right-hand side of the above identity by I_m . Using substitution

$x = tb + (1 - t)a$ on the left handside of (6) we get

$$\begin{aligned} & (-1)^n \int_a^b S_n(x, \sigma) f^{(n)}(x) dx \\ &= (-1)^n (b - a) \int_0^1 S_n(tb + (1 - t)a, \sigma) f^{(n)}(tb + (1 - t)a) dt. \end{aligned}$$

Using notation $C_n(t, \sigma') = (b - a)S_n(tb + (1 - t)a)$ identity (6) becomes

$$(-1)^n \int_0^1 C_n(t, \sigma') f^{(n)}(tb + (1 - t)a) dt = I_m. \tag{7}$$

Also, in the further text we use abbreviations $K_{jn}(t) = (b - a)P_{jn}(tb + (1 - t)a)$, $j = 1, \dots, m$, *i.e.*

$$C_n(t, \sigma') = \begin{cases} K_{1n}(t), & t \in [0, s_1] \\ K_{2n}(t), & t \in (s_1, s_2] \\ \vdots \\ K_{mn}(t), & t \in (s_{m-1}, 1]. \end{cases}$$

THEOREM 7. *Let $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be an n -times differentiable function on I° , $a, b \in I^\circ$, $a < b$ and $|f^{(n)}|^q$ be convex on $[a, b]$ for some $q \geq 1$ such that $S_n(t, \sigma) f^{(n)}(t)$ is integrable on $[a, b]$. Then*

$$\begin{aligned} |I_m| &\leq \sum_{j=1}^m \left(\int_{s_{j-1}}^{s_j} |K_{jn}(t)| dt \right)^{1-\frac{1}{q}} \left\{ |f^{(n)}(b)|^q \left(\int_{s_{j-1}}^{s_j} t |K_{jn}(t)| dt \right) \right. \\ &\quad \left. + |f^{(n)}(a)|^q \left(\int_{s_{j-1}}^{s_j} (1-t) |K_{jn}(t)| dt \right) \right\}^{\frac{1}{q}}. \end{aligned} \tag{8}$$

Proof. Using (7) and the power-mean inequality for $q \geq 1$ we have

$$\begin{aligned} |I_m| &\leq \int_0^1 |C_n(t, \sigma')| |f^{(n)}(tb + (1 - t)a)| dt \\ &= \sum_{j=1}^m \int_{s_{j-1}}^{s_j} |K_{jn}(t)| |f^{(n)}(tb + (1 - t)a)| dt \\ &\leq \sum_{j=1}^m \left(\int_{s_{j-1}}^{s_j} |K_{jn}(t)| dt \right)^{1-\frac{1}{q}} \left(\int_{s_{j-1}}^{s_j} |K_{jn}(t)| |f^{(n)}(tb + (1 - t)a)|^q dt \right)^{\frac{1}{q}}. \end{aligned}$$

Since $|f^{(n)}|^q$ is convex we have

$$\begin{aligned}
 |I_m| &\leq \sum_{j=1}^m \left(\int_{s_{j-1}}^{s_j} |K_{jn}(t)| dt \right)^{1-\frac{1}{q}} \\
 &\quad \times \left(\int_{s_{j-1}}^{s_j} |K_{jn}(t)| \left(t |f^{(n)}(b)|^q + (1-t) |f^{(n)}(a)|^q \right) dt \right)^{\frac{1}{q}} \\
 &= \sum_{j=1}^m \left(\int_{s_{j-1}}^{s_j} |K_{jn}(t)| dt \right)^{1-\frac{1}{q}} \\
 &\quad \times \left(|f^{(n)}(b)|^q \int_{s_{j-1}}^{s_j} t |K_{jn}(t)| dt + |f^{(n)}(a)|^q \int_{s_{j-1}}^{s_j} (1-t) |K_{jn}(t)| dt \right)^{\frac{1}{q}}.
 \end{aligned}$$

So, we deduce the desired result. \square

The case when $q = 1$ is significant, so we write that result as the following corollary.

COROLLARY 1. *If f satisfies assumptions of Theorem 7 with $q = 1$ which also means $|f^{(n)}|$ is convex on $[a, b]$, then*

$$|I_m| \leq |f^{(n)}(b)| \sum_{j=1}^m \int_{s_{j-1}}^{s_j} t |K_{jn}(t)| dt + |f^{(n)}(a)| \sum_{j=1}^m \int_{s_{j-1}}^{s_j} (1-t) |K_{jn}(t)| dt. \tag{9}$$

COROLLARY 2. *If f satisfies assumptions of Theorem 7 with $n = 1, m = 2$, then*

$$\begin{aligned}
 &\left| h[f(a) + f(b)] + (1-2h)f(x) - \frac{1}{b-a} \int_a^b f(u) du \right| \tag{10} \\
 &\leq (b-a) \left\{ \left(\frac{c^2}{2} - hc + h^2 \right)^{1-\frac{1}{q}} [\lambda_1 |f'(b)|^q + \lambda_2 |f'(a)|^q]^{\frac{1}{q}} \right. \\
 &\quad \left. + \left(\frac{c^2+1}{2} - (c+h)(1-h) \right)^{1-\frac{1}{q}} [\lambda_3 |f'(b)|^q + \lambda_4 |f'(a)|^q]^{\frac{1}{q}} \right\},
 \end{aligned}$$

where $x \in [a, b]$, $h \in [0, \frac{1}{2}]$, $c = \frac{x-a}{b-a}$ such that $h \leq c \leq 1-h$ and

$$\begin{aligned} \lambda_1 &= \frac{1}{6} (2c^3 - 3c^2h + 2h^3) \\ \lambda_2 &= \frac{1}{6} (-6ch + 6h^2 - 2h^3 - 2c^3 + 3c^2 + 3c^2h) \\ \lambda_3 &= \frac{1}{6} (2c^3 + 3c^2h - 3c^2 - 2h^3 + 6h^2 - 3h + 1) \\ \lambda_4 &= \frac{1}{6} (2 - 2c^3 + 6c^2 - 3c^2h + 6ch - 6c + 2h^3 - 3h). \end{aligned}$$

Proof. Let us consider a division $\sigma = \{a \leq x \leq b\}$ of an interval $[a, b]$. Let us define the kernel $S_1(t, \sigma)$ as

$$S_1(t, \sigma) = \begin{cases} P_{11}(t) = t - (1-h)a - hb, & t \in [a, x] \\ P_{21}(t) = t - ha - (1-h)b, & t \in (x, b] \end{cases}.$$

Polynomials P_{11} and P_{21} are obviously harmonic. Denote by $c := \frac{x-a}{b-a}$. Then we consider a corresponding division $\sigma' = \{0 \leq c \leq 1\}$ of the interval $[0, 1]$ and the kernel $C_1(t, \sigma')$ is equal to

$$C_1(t, \sigma') = \begin{cases} K_{11}(t) = (b-a)^2(t-h), & t \in [0, c] \\ K_{21}(t) = (b-a)^2(t-1+h), & t \in (c, 1] \end{cases}.$$

Putting in Theorem 7 the kernel $C_1(t, \sigma')$ after simple calculation with taking into account the condition $h \leq c \leq 1-h$ that is,

$$\int_0^c |t-h| dt = \int_0^h (h-t) dt + \int_h^c (t-h) dt = \frac{c^2}{2} - ch + h^2$$

and

$$\int_c^1 |t-1+h| dt = \int_c^{1-h} (1-h-t) dt + \int_{1-h}^1 (t-1+h) dt = \frac{c^2+1}{2} - (c+h)(1-h),$$

we get the desired inequality. \square

REMARK 1. If we set $h = \frac{1}{6}$ and $c = \frac{1}{2}$ in (10) we obtain inequality (2).

REMARK 2. For different selections of parameters h and c in (10) we obtain the following Ostrowski, Simpson and Hadamard type inequalities.

(i) For the case $q = 1$, $h = 0$ we have

$$\begin{aligned} & \left| f(x) - \frac{1}{b-a} \int_a^b f(u) du \right| \\ & \leq \frac{b-a}{6} [(1-3c^2+4c^3) |f'(b)| + (2-6c+9c^2-4c^3) |f'(a)|] \end{aligned}$$

which is an Ostowski type inequality. Furthermore, if $c = \frac{1}{2}$ we get inequality (3).

(ii) For the case $q = 1, h = \frac{1}{6}$ we obtain the following Simpson type inequality

$$\left| \frac{1}{6} [f(a) + 4f(x) + f(b)] - \frac{1}{b-a} \int_a^b f(u) du \right| \leq \frac{(b-a)}{6} \left[\left(\frac{2}{3} - 3c^2 + 4c^3 \right) |f'(b)| + \left(\frac{5}{3} - 6c + 9c^2 - 4c^3 \right) |f'(a)| \right]$$

which for $c = \frac{1}{2}$ collapses to inequality (1).

(iii) If $q = 1, h = c = \frac{1}{2}$ we obtain the following Hadamard type inequality (4).

THEOREM 8. Let $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be an n -times differentiable mapping on I° , $a, b \in I^\circ, a < b$, and $|f^{(n)}|^q$ is convex on $[a, b]$ for some $q > 1, S_n(t, \sigma) f^{(n)}(t)$ is integrable on $[a, b]$. Then the following inequality holds:

$$|I_m| \leq \left(\sum_{j=1}^m \int_{s_{j-1}}^{s_j} |K_{jn}(t)|^p dt \right)^{\frac{1}{p}} \left(\frac{|f^{(n)}(a)|^q + |f^{(n)}(b)|^q}{2} \right)^{\frac{1}{q}}, \tag{11}$$

where $\frac{1}{p} + \frac{1}{q} = 1$.

Proof. From (7) and property of modulus we have

$$\begin{aligned} |I_m| &= \left| \int_0^1 C_n(t, \sigma') f^{(n)}(tb + (1-t)a) dt \right| \\ &\leq \int_0^1 |C_n(t, \sigma') f^{(n)}(tb + (1-t)a)| dt. \end{aligned}$$

By using Hölder inequality we have

$$\begin{aligned} &\left| \int_0^1 C_n(t, \sigma') f^{(n)}(tb + (1-t)a) dt \right| \\ &\leq \left(\int_0^1 |C_n(t, \sigma')|^p dt \right)^{\frac{1}{p}} \left(\int_0^1 |f^{(n)}(tb + (1-t)a)|^q dt \right)^{\frac{1}{q}}. \end{aligned}$$

Since $|f^{(n)}|^q$ is convex, by Hadamard inequality we get

$$\int_0^1 |f^{(n)}(tb + (1-t)a)|^q dt \leq \frac{|f^{(n)}(a)|^q + |f^{(n)}(b)|^q}{2}.$$

Also we have

$$\int_0^1 |C_n(t, \sigma')|^p dt = \sum_{j=1}^m \int_{s_{j-1}}^{s_j} |K_{jn}(t)|^p dt.$$

By combining these we deduce the desired result. \square

COROLLARY 3. *If f satisfies assumptions of Theorem 8 with $n = 1, m = 2$, then*

$$\begin{aligned} & \left| h[f(a) + f(b)] + (1-2h)f(x) - \frac{1}{b-a} \int_a^b f(u) du \right| \tag{12} \\ & \leq (b-a) \left(\frac{2h^{1+p} + (c-h)^{1+p} + (1-c-h)^{1+p}}{1+p} \right)^{\frac{1}{p}} \left(\frac{|f'(a)|^q + |f'(b)|^q}{2} \right)^{\frac{1}{q}}, \end{aligned}$$

where $x \in [a, b], h \in [0, \frac{1}{2}], c = \frac{x-a}{b-a}$ such that $h \leq c \leq 1-h$.

Proof. Putting in Theorem 8 the kernel $C_1(t, \sigma')$ defined as in the proof of Corollary 2, after simple calculation we get desired inequality. \square

REMARK 3. For different selections of the parameters h and c in (12) we obtain the following Ostrowski, Hadamard and Simpson type inequalities

(i) For the case $h = 0$ we have

$$\left| f(x) - \frac{1}{b-a} \int_a^b f(u) du \right| \leq (b-a) \left(\frac{c^{1+p} + (1-c)^{1+p}}{1+p} \right)^{\frac{1}{p}} \left(\frac{|f'(a)|^q + |f'(b)|^q}{2} \right)^{\frac{1}{q}}.$$

(ii) For the case $h = \frac{1}{2}$ and $c = \frac{1}{2}$ we obtain (5).

(iii) For $h = \frac{1}{6}$ and $c = \frac{1}{2}$ we have

$$\begin{aligned} & \left| \frac{1}{6} \left[f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right] - \frac{1}{b-a} \int_a^b f(u) du \right| \\ & \leq \frac{(b-a)}{6} \left(\frac{2^{1+p} + 1}{3(p+1)} \right)^{\frac{1}{p}} \left(\frac{|f'(a)|^q + |f'(b)|^q}{2} \right)^{\frac{1}{q}}. \end{aligned}$$

THEOREM 9. Let $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be an n -times differentiable mapping on I° , $a, b \in I^\circ$, $a < b$, and $|f^{(n)}|^q$ is convex on $[a, b]$ for some $q > 1$, $S_n(t, \sigma) f^{(n)}(t)$ is integrable on $[a, b]$. Then the following inequality holds:

$$|I_m| \leq \sum_{j=1}^m \left(\int_{s_{j-1}}^{s_j} |K_{jn}(t)|^p dt \right)^{\frac{1}{p}} \left(\frac{|f^{(n)}(x_j)|^q + |f^{(n)}(x_{j-1})|^q}{2} \right)^{\frac{1}{q}}, \tag{13}$$

where $\frac{1}{p} + \frac{1}{q} = 1$.

Proof. By using (7), property of modulus and Hölder Inequality we have

$$\begin{aligned} |I_m| &\leq \int_0^1 |C_n(t, \sigma') f^{(n)}(tb + (1-t)a)| dt \\ &\leq \sum_{j=1}^m \int_{s_{j-1}}^{s_j} |K_{jn}(t) f^{(n)}(tb + (1-t)a)| dt \\ &\leq \sum_{j=1}^m \left(\int_{s_{j-1}}^{s_j} |K_{jn}(t)|^p dt \right)^{\frac{1}{p}} \left(\int_{s_{j-1}}^{s_j} |f^{(n)}(tb + (1-t)a)|^q dt \right)^{\frac{1}{q}}. \end{aligned}$$

Since $|f^{(n)}|^q$ is convex, by using Hadamard inequality we get

$$\int_{s_{j-1}}^{s_j} |f^{(n)}(tb + (1-t)a)|^q dt \leq \frac{|f^{(n)}(x_j)|^q + |f^{(n)}(x_{j-1})|^q}{2}.$$

So, this implies (13). \square

COROLLARY 4. If f satisfies assumptions of Theorem 8 with $n = 1$, $m = 2$, then

$$\begin{aligned} &\left| h[f(a) + f(b)] + (1-2h)f(x) - \frac{1}{b-a} \int_a^b f(u) du \right| \tag{14} \\ &\leq (b-a) \left\{ \left(\frac{h^{1+p} + (c-h)^{1+p}}{1+p} \right)^{\frac{1}{p}} \left(\frac{|f'(a)|^q + |f'(x)|^q}{2} \right)^{\frac{1}{q}} \right. \\ &\quad \left. + \left(\frac{h^{1+p} + (1-c-h)^{1+p}}{1+p} \right)^{\frac{1}{p}} \left(\frac{|f'(b)|^q + |f'(x)|^q}{2} \right)^{\frac{1}{q}} \right\}, \end{aligned}$$

where $x \in [a, b]$, $h \in [0, \frac{1}{2}]$, $c = \frac{x-a}{b-a}$ such that $h \leq c \leq 1-h$.

Proof. Putting in Theorem 9 the kernel $C_n(t, \sigma')$ defined as in the proof of Corollary 2, after simple calculation we get desired inequality. \square

REMARK 4. For different selections of the parameters h and c in (14) we obtain the following Ostrowski, Hadamard and Simpson type inequalities

(i) For the case $h = 0$ we have

$$\begin{aligned} & \left| f(x) - \frac{1}{b-a} \int_a^b f(u) du \right| \\ & \leq (b-a) \left\{ \left(\frac{c^{1+p}}{1+p} \right)^{\frac{1}{p}} \left(\frac{|f'(a)|^q + |f'(x)|^q}{2} \right)^{\frac{1}{q}} \right. \\ & \quad \left. + \left(\frac{(1-c)^{1+p}}{1+p} \right)^{\frac{1}{p}} \left(\frac{|f'(b)|^q + |f'(x)|^q}{2} \right)^{\frac{1}{q}} \right\}. \end{aligned}$$

(ii) For the case $h = \frac{1}{2}$ and $c = \frac{1}{2}$ we get the following inequality

$$\begin{aligned} & \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(u) du \right| \\ & \leq \frac{(b-a)}{2} \left(\frac{1}{2(1+p)} \right)^{\frac{1}{p}} \left\{ \left(\frac{|f'(a)|^q + |f'(\frac{a+b}{2})|^q}{2} \right)^{\frac{1}{q}} \right. \\ & \quad \left. + \left(\frac{|f'(b)|^q + |f'(\frac{a+b}{2})|^q}{2} \right)^{\frac{1}{q}} \right\}. \end{aligned}$$

(iii) For $h = \frac{1}{6}$ and $c = \frac{1}{2}$ we have the following Simpson type inequality

$$\begin{aligned} & \left| \frac{1}{6} \left[f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right] - \frac{1}{b-a} \int_a^b f(u) du \right| \\ & \leq \frac{(b-a)}{6} \left(\frac{2^{1+p} + 1}{6(p+1)} \right)^{\frac{1}{p}} \left\{ \left(\frac{|f'(a)|^q + |f'(\frac{a+b}{2})|^q}{2} \right)^{\frac{1}{q}} \right. \\ & \quad \left. + \left(\frac{|f'(b)|^q + |f'(\frac{a+b}{2})|^q}{2} \right)^{\frac{1}{q}} \right\}. \end{aligned}$$

REFERENCES

- [1] B. G. PACHPATTE, *On some inequalities for convex functions*, RGMIA Research Report Collection **6**, E (1) (2003).
- [2] G. H. TOADER, *On a generalization of the convexity*, *Mathematica* **30**, 53 (1988), 83–87.
- [3] H. KAVURMACI, M. AVCI AND M. E. ÖZDEMİR, *New inequalities of Hermite-Hadamard type for convex functions with applications*, *Journal of Inequalities and Applications* **86**, 1 (2011).
- [4] J. PEČARIĆ, S. VAROŠANEC, *Harmonic polynomials and generalization of Ostrowski inequality with Applications in Numerical Integration*, *Nonlinear Analysis Series A: Theory, Methods and Applications* **47**, 1 (2011), 2365–2374.
- [5] M. Z. SARIKAYA, E. SET AND M. E. ÖZDEMİR, *On new inequalities of Simpson type for convex functions*, RGMIA Research Report Collection **13**, 2 (2) (2010).
- [6] S. KOVAČ, J. PEČARIĆ, *Weighted version of general integral formula of Euler type*, *Math. Inequal. Appl.* **13**, 3 (2010), 579–599.
- [7] S. S. DRAGOMIR, R. P. AGARWAL, *Two inequalities for differentiable mappings and applications*, *Applied Mathematics Letters* **11**, 5 (1998), 91–95.
- [8] U. S. KIRMACI, *Inequalities for differentiable mappings and applications to special means of real numbers and to midpoint formula*, *Appl. Math. Comp.* **147**, 1 (2004), 137–146.

(Received January 16, 2016)

M. Emin Özdemir
Uludağ University, Education Faculty
Bursa, Turkey
e-mail: eminozdemir@uludag.edu.tr

Alper Ekinci
Ağrı İbrahim Çeçen University
Faculty of Science and Letters
Department of Mathematics
04100 Ağrı, Turkey
e-mail: alperekinci@hotmail.com