

CONTINUITY AND APPROXIMATE DIFFERENTIABILITY OF MULTISUBLINEAR FRACTIONAL MAXIMAL FUNCTIONS

FENG LIU

(Communicated by J. Pečarić)

Abstract. In this note we investigate the continuity and approximate differentiability of the m -sublinear fractional maximal operator

$$\mathfrak{M}_\alpha(\vec{f})(x) = \sup_{r>0} |B(x,r)|^{\alpha/d-m} \prod_{i=1}^m \int_{B(x,r)} |f_i(y)| dy,$$

where $m \geq 1$, $0 \leq \alpha < md$ and $\vec{f} = (f_1, \dots, f_m)$ with each $f_j \in L^1_{\text{loc}}(\mathbb{R}^d)$. More precisely, we prove that \mathfrak{M}_α maps $W^{1,p_1}(\mathbb{R}^d) \times \dots \times W^{1,p_m}(\mathbb{R}^d)$ into $W^{1,q}(\mathbb{R}^d)$ continuously, provided that $1 < p_1, \dots, p_m < \infty$ and $0 < \sum_{i=1}^m 1/p_i - \alpha/d = 1/q \leq 1$. We also show that the multisublinear fractional maximal functions $\mathfrak{M}_\alpha(\vec{f})$ are approximately differentiable a.e. if $\vec{f} = (f_1, f_2, \dots, f_m)$ with each $f_j \in L^1(\mathbb{R}^d)$ being approximately differentiable a.e. As applications, the corresponding results for fractional maximal operators are established.

1. Introduction

Let $d \geq 1$ and M denote the centered Hardy-Littlewood maximal operator on \mathbb{R}^d , i.e. for $f \in L^1_{\text{loc}}(\mathbb{R}^d)$,

$$Mf(x) = \sup_{r>0} \frac{1}{|B(x,r)|} \int_{B(x,r)} |f(y)| dy$$

for any $x \in \mathbb{R}^d$, where $B(x,r)$ is the ball in \mathbb{R}^d centered at x with radius r and $|B(x,r)|$ denotes the volume of $B(x,r)$. One of the cornerstones of harmonic analysis is the celebrated theorem of Hardy-Littlewood-Wiener that asserts that $M : L^p(\mathbb{R}^d) \rightarrow L^p(\mathbb{R}^d)$ is bounded for $1 < p \leq \infty$. For $p = 1$ we have $M : L^1(\mathbb{R}^d) \rightarrow L^{1,\infty}(\mathbb{R}^d)$ bounded. During the last several years, a considerable amount of attention has been given to investigate the behavior of differentiability under a maximal operator. The first work in this direction is due to Kinnunen [15] who proved that M is bounded on $W^{1,p}(\mathbb{R}^d)$ for $1 < p \leq \infty$, where $W^{1,p}(\mathbb{R}^d)$ is the first order Sobolev space, which consists of

Mathematics subject classification (2010): 42B25, 46E35.

Keywords and phrases: Multisublinear fractional maximal operator, Sobolev space, continuity, approximate differentiability.

This research was supported by the NNSF of China (No. 11526122), Scientific Research Foundation of Shandong University of Science and Technology for Recruited Talents (No. 2015RCJJ053), Research Award Fund for Outstanding Young Scientists of Shandong Province (No. BS2015SF012) and the Support Program for Outstanding Young Scientific and Technological Top-notch Talents of College of Mathematics and Systems Science (No. Sxy2016K01).

functions $f \in L^p(\mathbb{R}^d)$, whose first weak partial derivatives $D_i f$, $i = 1, 2, \dots, d$, belong to $L^p(\mathbb{R}^d)$. We endow $W^{1,p}(\mathbb{R}^d)$ with the norm

$$\|f\|_{1,p} = \|f\|_{L^p(\mathbb{R}^d)} + \|\nabla f\|_{L^p(\mathbb{R}^d)},$$

where $\nabla f = (D_1 f, \dots, D_d f)$ is the weak gradient of f . Later on, this paradigm has been extended to a local version in [16], to a fractional version in [17], to a bilinear version in [11], to a multisublinear fractional version in [25] and to a one-sided version in [24]. Due to the lack of reflexivity of L^1 , results for $p = 1$ are subtler. A crucial question on whether the operator $f \mapsto |\nabla M f|$ is bounded from $W^{1,1}(\mathbb{R}^d)$ to $L^1(\mathbb{R}^d)$, posed by Hajlasz and Onninen in [14], has been restricted to dimension $d = 1$. For example, see [4, 23, 34] for the uncentered Hardy-Littlewood maximal operator, [18] for the centered Hardy-Littlewood maximal operator, [10] for the fractional maximal operators, [26] for the multisublinear fractional maximal operators. Other interesting works related to this topic are [3, 5, 6, 12, 13, 22, 28].

In general, bounded non-sublinear operators need not be continuous (see [7] for a famous example). Continuity of the maximal operator in $L^p(\mathbb{R}^d)$ follows easily from its sublinearity and boundedness. Since we do not have sublinearity for the weak derivatives of the Hardy-Littlewood maximal function, the result of Kinnunen now leads us to another question: is $M : W^{1,p}(\mathbb{R}^d) \rightarrow W^{1,p}(\mathbb{R}^d)$ continuous for $1 < p < \infty$? This question was posed in [14, Question 3] where it was attributed to T. Iwaniec and has been studied by many authors. In 2007, Luiro first proved that M is continuous on $W^{1,p}(\mathbb{R}^d)$ for $1 < p < \infty$. Subsequently, Carneiro and Moreira [11] extended above result to the bilinear case. Later on, Luiro [30] extended the result of [29] to the local case. Recently, Luiro and Nuutinen [31] established the continuity of a class of discrete maximal operators in Sobolev space $W^{1,p}(\mathbb{R}^d)$ under certain sufficient assumptions.

As well known, the multilinear Calderón-Zygmund theory and the boundedness of these maximal operators on various function spaces have been extensively studied, for example, see [8, 9, 19, 20, 21, 25] and therein references. In this paper we focus on the regularity of the multisublinear fractional maximal operator. More precisely, let $m \geq 1$ and $\vec{f} = (f_1, \dots, f_m)$ with each $f_j \in L^1_{\text{loc}}(\mathbb{R}^d)$. For $0 \leq \alpha < md$, we define the multisublinear fractional maximal operator \mathfrak{M}_α by

$$\mathfrak{M}_\alpha(\vec{f})(x) = \sup_{r>0} |B(x,r)|^{\alpha/d-m} \prod_{i=1}^m \int_{B(x,r)} |f_i(y)| dy,$$

for any $x \in \mathbb{R}^d$. For $\alpha = 0$, the operator \mathfrak{M}_α recovers the classical multisublinear Hardy-Littlewood maximal operator. Specially, the centered Hardy-Littlewood maximal operator M corresponds to the special case of \mathfrak{M}_α for $m = 1$ and $\alpha = 0$. For $m = 1$ and $0 < \alpha < d$, the operator \mathfrak{M}_α reduces to the classical fractional maximal operator denoted by M_α , which has extensive applications in potential theory and partial differential equations (see [1, 2, 32, 33] for example).

Recently, it was shown in [25] that

$$\mathfrak{M}_\alpha : W^{1,p_1}(\mathbb{R}^d) \times \dots \times W^{1,p_m}(\mathbb{R}^d) \rightarrow W^{1,q}(\mathbb{R}^d)$$

is bounded for $1/q = \sum_{i=1}^m 1/p_i - \alpha/d$ with $1 < p_1, \dots, p_m < \infty$ and $1 \leq q < \infty$. A question that arises naturally is

QUESTION A. *Is the operator \mathfrak{M}_α continuous from $W^{1,p_1}(\mathbb{R}^d) \times \dots \times W^{1,p_m}(\mathbb{R}^d)$ to $W^{1,q}(\mathbb{R}^d)$, when $1/q = \sum_{i=1}^m 1/p_i - \alpha/d$, $1 < p_1, \dots, p_m < \infty$ and $1 \leq q < \infty$?*

Question A is the main motivation for this work. This problem is solved by our next theorem.

THEOREM 1. *Let $0 \leq \alpha < md$. Then \mathfrak{M}_α maps $W^{1,p_1}(\mathbb{R}^d) \times \dots \times W^{1,p_m}(\mathbb{R}^d)$ into $W^{1,q}(\mathbb{R}^d)$ continuously, where $1/q = \sum_{i=1}^m 1/p_i - \alpha/d$, $1 < p_1, \dots, p_m < \infty$ and $1 \leq q < \infty$.*

REMARK 1. Theorem 1 extends the continuity result in [29], which corresponds to the case $m = 1$ and $\alpha = 0$. In [17], Kinnunen and Saksman observed that $M_\alpha : W^{1,p}(\mathbb{R}^d) \rightarrow W^{1,q}(\mathbb{R}^d)$ is bounded if $1/q = 1/p - \alpha/d$ with $1 < p < \infty$ and $0 \leq \alpha < d/p$. As an application of Theorem 1, it is known that $M_\alpha : W^{1,p}(\mathbb{R}^d) \rightarrow W^{1,q}(\mathbb{R}^d)$ is continuous if $1/q = 1/p - \alpha/d$ with $1 < p < \infty$ and $0 \leq \alpha < d/p$.

To the best of my knowledge Hajlasz and Onninen’s question remains open in dimension $d \geq 2$. Motivated by this challenging problem, Hajlasz and Maly [13] established the approximate differentiability of the Hardy-Littlewood maximal operator, which can be listed as follows.

THEOREM B. ([13]) *Let $f \in L^1(\mathbb{R}^d)$ be approximately differentiable a.e., then Mf is also approximately differentiable a.e.*

For the endpoint regularity of \mathfrak{M}_α , it was shown in [26, 27] that if $d \geq 2$, $1 \leq \alpha < m(d-1) + 1$ and $\vec{f} = (f_1, \dots, f_m)$ with each $f_i \in W^{1,1}(\mathbb{R}^d)$, then $\mathfrak{M}_\alpha(\vec{f})$ is weakly differentiable and

$$\|\nabla \mathfrak{M}_\alpha(\vec{f})\|_{L^{\frac{d}{m(d-1)-\alpha+1}}(\mathbb{R}^d)} \leq C \prod_{j=1}^m \|\nabla f_j\|_{L^1(\mathbb{R}^d)}.$$

A natural and interesting question is that what about the smoothness properties of the maximal function $\mathfrak{M}_\alpha(\vec{f})$ if $0 \leq \alpha < 1$ or $m(d-1) + 1 \leq \alpha < md$ and $\vec{f} = (f_1, \dots, f_m)$ with each $f_i \in W^{1,1}(\mathbb{R}^d)$. Motivated by Theorem B, we shall establish the following result.

THEOREM 2. *Let $0 \leq \alpha \leq md - 1$ and $\vec{f} = (f_1, \dots, f_m)$ with each $f_j \in L^1(\mathbb{R}^d)$ being approximately differentiable a.e., then $\mathfrak{M}_\alpha(\vec{f})$ is approximately differentiable a.e.*

REMARK 2. Clearly, Theorem 2 extends Theorem B, which corresponds to the case $m = 1$ and $\alpha = 0$. Since every function $f \in W^{1,1}(\mathbb{R}^d)$ is approximately differentiable a.e., Theorem 2 implies that if $0 \leq \alpha \leq md - 1$ and $\vec{f} = (f_1, \dots, f_m)$ with each $f_j \in W^{1,1}(\mathbb{R}^d)$, then $\mathfrak{M}_\alpha(\vec{f})$ is approximately differentiable a.e. Moreover, we know from Theorem 2 that the fractional maximal function $M_\alpha f$ is approximately differentiable a.e. if $f \in L^1(\mathbb{R}^d)$ is approximately differentiable a.e. and $0 \leq \alpha \leq d - 1$.

The rest of this paper is organized as follows. In Section 2 we prove Theorem 1, the proof of Theorem 2 will be given in Section 3. We remark that the main ideas in our proofs are greatly motivated by [13, 29], but our methods and techniques are more subtle and complex than that of [13, 29]. The main ingredient of our proof of Theorem 1 is to present an explicit formula for the derivative of the multisublinear fractional maximal functions (see Lemma 2). In what follows, we use the following conventions

$$\prod_{i \in \emptyset} a_i = 1 \quad \text{and} \quad \sum_{i \in \emptyset} a_i = 0.$$

2. Proof of Theorem 1

In this section we follow carefully the proof for the continuity in $W^{1,p}(\mathbb{R}^d)$ of the Hardy-Littlewood maximal operator in [29] and simply adjust the notation to our context.

For $R > 0$, we denote by B_R the ball of radius R centered at the origin. For $A \subset \mathbb{R}^d$ and $x \in \mathbb{R}^d$, we define

$$d(x, A) := \inf_{a \in A} |x - a| \quad \text{and} \quad A_{(\lambda)} := \{x \in \mathbb{R}^d; d(x, A) \leq \lambda\} \quad \text{for } \lambda \geq 0.$$

We denote by $\|f\|_{p,A}$ the L^p -norm of $f\chi_A$ for all measurable sets $A \subset \mathbb{R}^d$. Let $\vec{f} = (f_1, \dots, f_m) \in L^{p_1}(\mathbb{R}^d) \times \dots \times L^{p_m}(\mathbb{R}^d)$ with $1 < p_1, \dots, p_m < \infty$, $1 \leq q < \infty$ and $1/q = \sum_{i=1}^m 1/p_i - \alpha/d$. For a fixed point $x \in \mathbb{R}^d$, we define the set $\mathcal{R}(\vec{f})(x)$ by

$$\begin{aligned} & \mathcal{R}(\vec{f})(x) \\ & := \left\{ r \geq 0 : \mathfrak{M}_\alpha(\vec{f})(x) = \limsup_{r_k \rightarrow r} |B(x, r_k)|^{\alpha/d-m} \prod_{i=1}^m \int_{B(x, r_k)} |f_i(y)| dy \text{ for some } r_k > 0 \right\}. \end{aligned}$$

We also define the function $u_{x, \vec{f}} : [0, \infty) \rightarrow \mathbb{R}$ by

$$u_{x, \vec{f}}(0) = \begin{cases} \prod_{i=1}^m |f_i(x)|, & \text{if } \alpha = 0; \\ 0, & \text{if } 0 < \alpha < md, \end{cases}$$

$$u_{x, \vec{f}}(r) = |B(x, r)|^{\alpha/d-m} \prod_{i=1}^m \int_{B(x, r)} |f_i(y)| dy \quad \text{when } r \in (0, \infty).$$

We notice that the following facts are valid: (i) $u_{x, \vec{f}}$ is continuous on $(0, \infty)$ for all $x \in \mathbb{R}^d$ and at $r = 0$ for a.e. $x \in \mathbb{R}^d$; (ii) $\lim_{r \rightarrow \infty} u_{x, \vec{f}}(r) = 0$ since $u_{x, \vec{f}}(r) \leq \prod_{i=1}^m \|f_i\|_{L^{p_i}(\mathbb{R}^d)} |B(x, r)|^{-1/q}$ for any $r > 0$ and $x \in \mathbb{R}^d$; (iii) $\mathcal{R}(\vec{f})(x)$ is nonempty and closed for all $x \in \mathbb{R}^d$ and

$$\mathfrak{M}_\alpha(\vec{f})(x) = u_{x, \vec{f}}(r) \text{ if } 0 < r \in \mathcal{R}(\vec{f})(x), \quad \forall x \in \mathbb{R}^d,$$

$$\mathfrak{M}_\alpha(\vec{f})(x) = u_{x, \vec{f}}(0) \text{ for a.e. } x \in \mathbb{R}^d \text{ such that } 0 \in \mathcal{R}(\vec{f})(x).$$

Motivated by the idea in [29], we can get the following

LEMMA 1. Let $\vec{f}_j = (f_{1,j}, f_{2,j}, \dots, f_{m,j})$. Suppose that $f_{i,j} \rightarrow f_i$ in $L^{p_i}(\mathbb{R}^d)$ when $j \rightarrow \infty$ for all $i = 1, 2, \dots, m$, where $1 < p_1, \dots, p_m < \infty$, $1 \leq q < \infty$ and $1/q = \sum_{i=1}^m 1/p_i - \alpha/d$. Then for all $R > 0$ and $\lambda > 0$, we have

$$\lim_{j \rightarrow \infty} |\{x \in B_R; \mathcal{R}(\vec{f}_j)(x) \not\leq \mathcal{R}(\vec{f})(x)_{(\lambda)}\}| = 0. \quad (1)$$

Proof. Without loss of generality we may assume that all $f_{i,j} \geq 0$ and $f_i \geq 0$. By the similar argument as in the proof of [29, Lemma 2.2], we can conclude that the set $\{x \in \mathbb{R}^d; \mathcal{R}(\vec{f}_j)(x) \not\leq \mathcal{R}(\vec{f})(x)_{(\lambda)}\}$ is measurable for any $j \in \mathbb{Z}$. Let $\lambda > 0$, $R > 0$ and $\varepsilon \in (0, 1)$. We can claim that for a.e. $x \in B_R$, there exists $\gamma(x) \in \mathbb{N} \setminus \{0\}$ such that

$$u_{x, \vec{f}}(r) < \mathfrak{M}_\alpha(\vec{f})(x) - \frac{1}{\gamma(x)}, \quad \text{when } d(r, \mathcal{R}(\vec{f})(x)) > \lambda. \quad (2)$$

Otherwise, for a.e. $x \in B_R$, there exists a bounded sequence of radii $\{r_k\}_{k=1}^\infty$ such that

$$\lim_{k \rightarrow \infty} u_{x, \vec{f}}(r_k) = \mathfrak{M}_\alpha(\vec{f})(x) \quad \text{and } d(r_k, \mathcal{R}(\vec{f})(x)) > \lambda.$$

We can choose a subsequence $\{s_k\}_{k=1}^\infty$ of $\{r_k\}_{k=1}^\infty$ such that $s_k \rightarrow r$ as $k \rightarrow \infty$. It follows that $r \in \mathcal{R}(\vec{f})(x)$ and $d(r, \mathcal{R}(\vec{f})(x)) \geq \lambda$, which is a contradiction. Thus, (2) holds. From (2) we can conclude that there exists $\gamma = \gamma(R, \lambda, \varepsilon) \in \mathbb{N} \setminus \{0\}$ and a measurable set E with $|E| < \varepsilon$ such that

$$\begin{aligned} B_R &\subset \{x \in \mathbb{R}^d : u_{x, \vec{f}}(r) < \mathfrak{M}_\alpha(\vec{f})(x) - \gamma^{-1} \text{ if } d(r, \mathcal{R}(\vec{f})(x)) > \lambda\} \cup E \\ &\subset A_{1,j} \cup A_{2,j} \cup A_{3,j} \cup E, \end{aligned} \quad (3)$$

where

$$A_{1,j} := \{x \in \mathbb{R}^d : |\mathfrak{M}_\alpha(\vec{f}_j)(x) - \mathfrak{M}_\alpha(\vec{f})(x)| \geq (4\gamma)^{-1}\},$$

$$A_{2,j} := \{x \in \mathbb{R}^d : |u_{x, \vec{f}_j}(r) - u_{x, \vec{f}}(r)| \geq (2\gamma)^{-1} \text{ for some } r \text{ such that } d(r, \mathcal{R}(\vec{f})(x)) > \lambda\},$$

$$A_{3,j} := \{x \in \mathbb{R}^d : u_{x, \vec{f}_j}(r) < \mathfrak{M}_\alpha(\vec{f}_j)(x) - (4\gamma)^{-1} \text{ if } d(r, \mathcal{R}(\vec{f})(x)) > \lambda\}.$$

Let \bar{A} be the set of all points x such that x is a Lebesgue point of all f_j . Note that $|\mathbb{R}^d \setminus \bar{A}| = 0$. One can easily check that $A_{3,j} \cap \bar{A} \subset \cup \{x \in \mathbb{R}^d : \mathcal{R}(\vec{f}_j)(x) \subset \mathcal{R}(\vec{f})(x)_{(\lambda)}\}$. This together with (3) yields that

$$\{x \in B_R; \mathcal{R}(\vec{f}_j)(x) \not\leq \mathcal{R}(\vec{f})(x)_{(\lambda)}\} \subset A_{1,j} \cup A_{2,j} \cup E \cup (\mathbb{R}^d \setminus \bar{A}).$$

It follows that

$$|\{x \in B_R; \mathcal{R}(\vec{f}_j)(x) \not\leq \mathcal{R}(\vec{f})(x)_{(\lambda)}\}| \leq |A_{1,j}| + |A_{2,j}| + \varepsilon. \quad (4)$$

Since $f_{i,j} \rightarrow f_i$ in $L^{p_i}(\mathbb{R}^d)$ when $j \rightarrow \infty$ for all $i = 1, 2, \dots, m$, then for any fixed $\varepsilon \in (0, 1)$, there exists $N_0 = N_0(\varepsilon) \in \mathbb{N}$ such that

$$\|f_{i,j} - f_i\|_{L^{p_i}(\mathbb{R}^d)} < \frac{\varepsilon}{\gamma} \quad \text{and } \|f_{i,j}\|_{L^{p_i}(\mathbb{R}^d)} \leq \|f_i\|_{L^{p_i}(\mathbb{R}^d)} + 1 \quad (5)$$

for any $j \geq N_0$ and $i = 1, 2, \dots, m$. We can write

$$\begin{aligned}
& |\mathfrak{M}_\alpha(\vec{f}_j)(x) - \mathfrak{M}_\alpha(\vec{f})(x)| \\
& \leq \sup_{r>0} |B(x,r)|^{\alpha/d-m} \left| \prod_{i=1}^m \int_{B(x,r)} f_{i,j}(y) dy - \prod_{i=1}^m \int_{B(x,r)} f_i(y) dy \right| \\
& \leq \sum_{i=1}^m \sup_{r>0} |B(x,r)|^{\alpha/d-m} \prod_{\mu=1}^{i-1} \int_{B(x,r)} f_\mu(y) dy \prod_{v=i+1}^m \int_{B(x,r)} f_{v,j}(y) dy \\
& \quad \times \int_{B(x,r)} |f_{i,j}(y) - f_i(y)| dy \\
& = \sum_{i=1}^m \mathfrak{M}_\alpha(\vec{F}_j^i)(x)
\end{aligned} \tag{6}$$

for any $x \in \mathbb{R}^d$, where $\vec{F}_j^i = (f_1, \dots, f_{i-1}, f_{i,j} - f_i, f_{i+1,j}, \dots, f_{m,j})$. It follows from (5) and (6) that

$$\begin{aligned}
|A_{1,j}| & \leq \left| \left\{ x \in \mathbb{R}^d : \sum_{i=1}^m \mathfrak{M}_\alpha(\vec{F}_j^i)(x) \geq (4\gamma)^{-1} \right\} \right| \\
& \leq \sum_{i=1}^m |\{x \in \mathbb{R}^d : \mathfrak{M}_\alpha(\vec{F}_j^i)(x) \geq (4m\gamma)^{-1}\}| \\
& \leq (4m\gamma)^q \sum_{i=1}^m \|\mathfrak{M}_\alpha(\vec{F}_j^i)\|_{L^q(\mathbb{R}^d)}^q \\
& \leq C(m, \gamma, q, \alpha, d, p_1, \dots, p_m) \varepsilon
\end{aligned} \tag{7}$$

for any $j \geq N_0$. Similarly, we can get

$$|A_{2,j}| \leq C(m, \gamma, q, \alpha, d, p_1, \dots, p_m) \varepsilon \tag{8}$$

for any $j \geq N_0$. (1) follows from (4), (7) and (8). \square

Let $e_l = (0, \dots, 0, 1, 0, \dots, 0)$ be the canonical l -th base vector in \mathbb{R}^d for $l = 1, 2, \dots, d$. For any fixed $i = 1, 2, \dots, m$, $h > 0$ and $f_i \in L^p(\mathbb{R}^d)$ with $p \geq 1$, define

$$f_{i,h}^l(x) = \frac{f_{\tau(h)}^{i,l}(x) - f_i(x)}{h} \quad \text{and} \quad f_{\tau(h)}^{i,l}(x) = f_i(x + he_l).$$

It is well known that for $p \geq 1$, $f_{\tau(h)}^{i,l} \rightarrow f_i$ in $L^p(\mathbb{R}^d)$ when $h \rightarrow 0$, and if $f_i \in W^{1,p}(\mathbb{R}^d)$ we have $f_{i,h}^l \rightarrow D_l f_i$ in $L^p(\mathbb{R}^d)$ when $h \rightarrow 0$. Let A, B be two subsets of \mathbb{R}^d , we define the Hausdorff distance of A and B by

$$\pi(A, B) := \inf\{\delta > 0 : A \subset B_{(\delta)} \text{ and } B \subset A_{(\delta)}\}.$$

Applying Lemma 1 and the argument similar to that in the proof of [29, Corollary 2.3], we can get the following result. The details are omitted.

COROLLARY 1. *Let $\vec{f} = (f_1, \dots, f_m) \in L^{p_1}(\mathbb{R}^d) \times \dots \times L^{p_m}(\mathbb{R}^d)$ with $1 \leq q < \infty$, $1 < p_1, \dots, p_m < \infty$ and $1/q = \sum_{i=1}^m 1/p_i - \alpha/d$. Then for all $R > 0$, $\lambda > 0$ and $l = 1, 2, \dots, d$, we have*

$$|\{x \in B_R : \pi(\mathcal{R}(\vec{f})(x), \mathcal{R}(\vec{f})(x + he_l)) > \lambda\}| \rightarrow 0 \text{ when } h \rightarrow 0.$$

We now state several formulas for the derivative of the multisublinear maximal fractional functions, which play a key role in the proof of Theorem 1.

LEMMA 2. Let $\vec{f} = (f_1, f_2, \dots, f_m) \in W^{1,p_1}(\mathbb{R}^d) \times \dots \times W^{1,p_m}(\mathbb{R}^d)$ with $1 \leq q < \infty$, $1 < p_1, \dots, p_m < \infty$ and $1/q = \sum_{i=1}^m 1/p_i - \alpha/d$. Then for any $l = 1, 2, \dots, d$ and a.e. $x \in \mathbb{R}^d$, we have

$$D_l \mathfrak{M}_\alpha(\vec{f})(x) = \sum_{i=1}^m |B(x, r)|^{\alpha/d-m} \left(\prod_{1 \leq j \neq i \leq m} \int_{B(x, r)} |f_j(y)| dy \right) \\ \times \int_{B(x, r)} D_l(|f_i|)(y) dy \text{ for all } 0 < r \in \mathcal{R}(\vec{f})(x),$$

$$D_l \mathfrak{M}_\alpha(\vec{f})(x) = \begin{cases} \sum_{i=1}^m D_l |f_i|(x) \prod_{1 \leq j \neq i \leq m} |f_j(x)|, & \text{if } \alpha = 0 \text{ and } 0 \in \mathcal{R}(\vec{f})(x), \\ 0, & \text{if } 0 < \alpha < md \text{ and } 0 \in \mathcal{R}(\vec{f})(x). \end{cases}$$

Proof. We may assume without loss of generality that all $f_i \geq 0$, since $|f_i| \in W^{1,p_i}(\mathbb{R}^d)$ if $f_i \in W^{1,p_i}(\mathbb{R}^d)$. Let $R > 0$. Invoking Corollary 1, we can choose a sequence $\{s_k\}_{k=1}^\infty$, $s_k > 0$ and $s_k \rightarrow 0$ such that $\lim_{k \rightarrow \infty} \pi(\mathcal{R}(\vec{f})(x), \mathcal{R}(\vec{f})(x + s_k e_l)) = 0$ for a.e. $x \in B_R$. Then for any $i = 1, 2, \dots, m$ and $l = 1, 2, \dots, d$ we have

$$\|f_{\tau(s_k)}^{i,l} - f_i\|_{L^{p_i}(\mathbb{R}^d)} \rightarrow 0 \text{ as } k \rightarrow \infty,$$

$$\|f_{i,s_k}^l - D_l f_i\|_{L^{p_i}(\mathbb{R}^d)} \rightarrow 0 \text{ as } k \rightarrow \infty,$$

$$\|M(f_{\tau(s_k)}^{i,l} - f_i)\|_{L^{p_i}(\mathbb{R}^d)} \rightarrow 0 \text{ as } k \rightarrow \infty,$$

$$\|M(f_{i,s_k}^l - D_l f_i)\|_{L^{p_i}(\mathbb{R}^d)} \rightarrow 0 \text{ as } k \rightarrow \infty,$$

$$\|(\mathfrak{M}_\alpha(\vec{f}))_{s_k}^l - D_l \mathfrak{M}_\alpha(\vec{f})\|_{L^q(\mathbb{R}^d)} \rightarrow 0 \text{ as } k \rightarrow \infty.$$

Furthermore, there exists a subsequence $\{h_k\}_{k=1}^\infty$ of $\{s_k\}_{k=1}^\infty$ and a measurable set $A_1 \subset B_R$ such that $|B_R \setminus A_1| = 0$ and

(i) $f_{\tau(h_k)}^{i,l}(x) \rightarrow f_i(x)$, $f_{i,h_k}^l(x) \rightarrow D_l f_i(x)$, $M(f_{\tau(h_k)}^{i,l} - f_i)(x) \rightarrow 0$, $M(f_{i,h_k}^l - D_l f_i)(x) \rightarrow 0$ and $(\mathfrak{M}_\alpha(\vec{f}))_{h_k}^l(x) \rightarrow D_l \mathfrak{M}_\alpha(\vec{f})(x)$ when $k \rightarrow \infty$ for any $x \in A_1$, $i = 1, 2, \dots, m$ and $l = 1, \dots, d$;

(ii) $\lim_{k \rightarrow \infty} \pi(\mathcal{R}(\vec{f})(x), \mathcal{R}(\vec{f})(x + h_k e_l)) = 0$ for any $x \in A_1$.

Let

$$A_2 := \bigcap_{k=1}^\infty \{x \in \mathbb{R}^d : \mathfrak{M}_\alpha(\vec{f})(x + h_k e_l) \geq u_{x+h_k e_l, \vec{f}}(0)\},$$

$$A_3 := \{x \in \mathbb{R}^d : \mathfrak{M}_\alpha(\vec{f})(x) = u_{x, \vec{f}}(0) \text{ if } 0 \in \mathcal{R}(\vec{f})(x)\},$$

$$A_4 := \bigcap_{k=1}^\infty \{x \in \mathbb{R}^d : \mathfrak{M}_\alpha(\vec{f})(x + h_k e_l) = u_{x+h_k e_l, \vec{f}}(0) \text{ if } 0 \in \mathcal{R}(\vec{f})(x + h_k e_l)\}.$$

One can easily check that $|B_R \setminus A_i| = 0$ for any $i = 2, 3, 4$. Let $x \in A_1 \cap A_2 \cap A_3 \cap A_4$ be a Lebesgue point of all f_i and $D_l f_i$ and $r \in \mathcal{R}(\vec{f})(x)$, there exists radii $r_k \in \mathcal{R}(\vec{f})(x + h_k e_l)$ such that $\lim_{k \rightarrow \infty} r_k = r$. We now complete the rest of proof by considering the following two cases:

Case A ($r > 0$). Without loss of generality we may assume that all $r_k > 0$. We can write

$$\begin{aligned}
D_l \mathfrak{M}_\alpha(\vec{f})(x) &= \lim_{k \rightarrow \infty} \frac{1}{h_k} (\mathfrak{M}_\alpha(\vec{f})(x + h_k e_l) - \mathfrak{M}_\alpha(\vec{f})(x)) \\
&\leq \lim_{k \rightarrow \infty} \frac{1}{h_k} (u_{x+h_k e_l, \vec{f}}(r_k) - u_{x, \vec{f}}(r_k)) \\
&= \lim_{k \rightarrow \infty} \frac{1}{h_k} \sum_{j=1}^m |B(x, r_k)|^{\alpha/d-m} \int_{B(x, r_k)} (f_j(y + h_k e_l) - f_j(y)) dy \\
&\quad \times \prod_{\mu=1}^{j-1} \int_{B(x, r_k)} f_\mu(y + h_k e_l) dy \prod_{v=j+1}^m \int_{B(x, r_k)} f_v(y) dy \\
&= \sum_{j=1}^m \lim_{k \rightarrow \infty} |B(x, r_k)|^{\alpha/d-m} \int_{B(x, r_k)} f_{j, h_k}^l(y) dy \\
&\quad \times \prod_{\mu=1}^{j-1} \int_{B(x, r_k)} f_{\tau(h_k)}^{\mu, l}(y) dy \prod_{v=j+1}^m \int_{B(x, r_k)} f_v(y) dy.
\end{aligned} \tag{9}$$

Since $\lim_{k \rightarrow \infty} |B(x, r_k)| = |B_r|$, $f_{\tau(h_k)}^{i, l} \chi_{B(x, r_k)} \rightarrow f_i \chi_{B(x, r)}$ and $f_{j, h_k}^l \chi_{B(x, r_k)} \rightarrow D_l f_j \chi_{B(x, r)}$ in $L^1(\mathbb{R}^d)$ as $k \rightarrow \infty$. It follows that

$$D_l \mathfrak{M}_\alpha(\vec{f})(x) \leq \sum_{i=1}^m |B(x, r)|^{\alpha/d-m} \left(\prod_{1 \leq j \neq i \leq m} \int_{B(x, r)} f_j(y) dy \right) \int_{B(x, r)} D_l f_i(y) dy.$$

On the other hand, we have

$$\begin{aligned}
D_l \mathfrak{M}_\alpha(\vec{f})(x) &= \lim_{k \rightarrow \infty} \frac{1}{h_k} (\mathfrak{M}_\alpha(\vec{f})(x + h_k e_l) - \mathfrak{M}_\alpha(\vec{f})(x)) \\
&\geq \lim_{k \rightarrow \infty} \frac{1}{h_k} (u_{x+h_k e_l, \vec{f}}(r) - u_{x, \vec{f}}(r)) \\
&= \lim_{k \rightarrow \infty} \frac{1}{h_k} \sum_{j=1}^m |B(x, r)|^{\alpha/d-m} \int_{B(x, r)} (f_j(y + h_k e_l) - f_j(y)) dy \\
&\quad \times \prod_{\mu=1}^{j-1} \int_{B(x, r)} f_\mu(y + h_k e_l) dy \prod_{v=j+1}^m \int_{B(x, r)} f_v(y) dy \\
&= \sum_{j=1}^m |B(x, r)|^{\alpha/d-m} \lim_{k \rightarrow \infty} \int_{B(x, r)} f_{j, h_k}^l(y) dy \\
&\quad \times \prod_{\mu=1}^{j-1} \int_{B(x, r)} f_{\tau(h_k)}^{\mu, l}(y) dy \prod_{v=j+1}^m \int_{B(x, r)} f_v(y) dy \\
&= \sum_{i=1}^m |B(x, r)|^{\alpha/d-m} \int_{B(x, r)} D_l f_i(y) dy \left(\prod_{1 \leq j \neq i \leq m} \int_{B(x, r)} f_j(y) dy \right).
\end{aligned}$$

Case B ($r = 0$). When $0 < \alpha < md$. Since $\mathfrak{M}_\alpha(\vec{f})(x) = 0$ and then $\prod_{i=1}^m f_i(y) = 0$ for a.e. $y \in \mathbb{R}^d$. Thus $\mathfrak{M}_\alpha(\vec{f}) \equiv 0$ and $D_l \mathfrak{M}_\alpha(\vec{f})(x) = 0$. We now consider the case $\alpha = 0$. Let us begin with estimating the lower bound of $D_l \mathfrak{M}_\alpha(\vec{f})(x)$. We can write

$$\begin{aligned}
 D_l \mathfrak{M}_\alpha(\vec{f})(x) &= \lim_{k \rightarrow \infty} \frac{1}{h_k} (\mathfrak{M}_\alpha(\vec{f})(x + h_k e_l) - \mathfrak{M}_\alpha(\vec{f})(x)) \\
 &\geq \lim_{k \rightarrow \infty} \frac{1}{h_k} \left(\prod_{i=1}^m f_i(x + h_k e_l) - \prod_{i=1}^m f_i(x) \right) \\
 &= \lim_{k \rightarrow \infty} \frac{1}{h_k} \sum_{j=1}^m (f_j(x + h_k e_l) - f_j(x)) \prod_{i=1}^{j-1} f_i(x + h_k e_l) \prod_{i=j+1}^m f_i(x) \\
 &= \sum_{i=1}^m D_l f_i(x) \prod_{1 \leq j \neq i \leq m} f_j(x).
 \end{aligned} \tag{10}$$

Below we estimate the upper bound of $D_l \mathfrak{M}_\alpha(\vec{f})(x)$. If we have $r_k = 0$ for infinitely many k , then

$$\begin{aligned}
 D_l \mathfrak{M}_\alpha(\vec{f})(x) &= \lim_{k \rightarrow \infty} \frac{1}{h_k} (\mathfrak{M}_\alpha(\vec{f})(x + h_k e_l) - \mathfrak{M}_\alpha(\vec{f})(x)) \\
 &= \lim_{k \rightarrow \infty} \frac{1}{h_k} \left(\prod_{i=1}^m f_i(x + h_k e_l) - \prod_{i=1}^m f_i(x) \right) \\
 &= \lim_{k \rightarrow \infty} \frac{1}{h_k} \sum_{j=1}^m (f_j(x + h_k e_l) - f_j(x)) \prod_{\mu=1}^{j-1} f_\mu(x + h_k e_l) \prod_{\nu=j+1}^m f_\nu(x) \\
 &= \sum_{i=1}^m D_l f_i(x) \prod_{1 \leq j \neq i \leq m} f_j(x).
 \end{aligned}$$

If there exists $k_0 \in \mathbb{N}$ such that $r_k > 0$ when $k \geq k_0$. We get from (9) that

$$\begin{aligned}
 D_l \mathfrak{M}_\alpha(\vec{f})(x) &\leq \sum_{j=1}^m \lim_{k \rightarrow \infty} |B(x, r_k)|^{-m} \int_{B(x, r_k)} f_{j, h_k}^l(y) dy \\
 &\quad \times \prod_{\mu=1}^{j-1} \int_{B(x, r_k)} f_{\tau(h_k)}^{\mu, l}(y) dy \prod_{\nu=j+1}^m \int_{B(x, r_k)} f_\nu(y) dy.
 \end{aligned} \tag{11}$$

Since

$$\begin{aligned}
 &\lim_{k \rightarrow \infty} \left| \frac{1}{|B(x, r_k)|} \int_{B(x, r_k)} f_{j, h_k}^l(y) dy - D_l f_j(x) \right| \\
 &= \lim_{k \rightarrow \infty} \left| \frac{1}{|B(x, r_k)|} \int_{B(x, r_k)} (f_{j, h_k}^l(y) - D_l f_j(y)) dy \right| \\
 &\leq \lim_{k \rightarrow \infty} M(f_{j, h_k}^l - D_l f_j)(x) = 0.
 \end{aligned}$$

It follows that

$$\lim_{k \rightarrow \infty} \frac{1}{|B(x, r_k)|} \int_{B(x, r_k)} f_{j, h_k}^l(y) dy = D_l f_j(x). \tag{12}$$

By argument similar to those used to derive (11),

$$\lim_{k \rightarrow \infty} \frac{1}{|B(x, r_k)|} \int_{B(x, r_k)} f_{\tau(h_k)}^{\mu, l}(y) dy = f_{\mu}(x). \quad (13)$$

It follows from (11)–(13) that

$$D_l \mathfrak{M}_{\alpha}(\vec{f})(x) \leq \sum_{i=1}^m D_l f_i(x) \prod_{1 \leq j \neq i \leq m} f_j(x),$$

which together with (10) yields

$$D_l \mathfrak{M}_{\alpha}(\vec{f})(x) = \sum_{i=1}^m D_l f_i(x) \prod_{1 \leq j \neq i \leq m} f_j(x).$$

Since R was arbitrary and $|B_R \setminus (A_1 \cap A_2 \cap A_3 \cap A_4)| = 0$. This proves Lemma 2. \square

We are now in the position of proving Theorem 1.

Proof of Theorem 1. Let $p_1, \dots, p_m, q, \alpha$ be given as in Theorem 1 and $\vec{f} = (f_1, \dots, f_m) \in W^{1, p_1}(\mathbb{R}^d) \times \dots \times W^{1, p_m}(\mathbb{R}^d)$. For $i = 1, 2, \dots, m$, let $\vec{f}_j = (f_{1,j}, \dots, f_{m,j})$ and $f_{i,j} \rightarrow f_i$ in $W^{1, p_i}(\mathbb{R}^d)$ when $j \rightarrow \infty$. We get from (6) that

$$\begin{aligned} \|\mathfrak{M}_{\alpha}(\vec{f}_j) - \mathfrak{M}_{\alpha}(\vec{f})\|_{L^q(\mathbb{R}^d)} &\leq \sum_{i=1}^m \|\mathfrak{M}_{\alpha}(\vec{F}_j^i)\|_{L^q(\mathbb{R}^d)} \\ &\leq \sum_{i=1}^m \|f_{i,j} - f_i\|_{L^{p_i}(\mathbb{R}^d)} \prod_{\mu=1}^{i-1} \|f_{\mu}\|_{L^{p_{\mu}}(\mathbb{R}^d)} \prod_{\nu=i+1}^m \|f_{\nu,j}\|_{L^{p_{\nu}}(\mathbb{R}^d)}, \end{aligned}$$

where \vec{F}_j^i is given as in (6). It follows that

$$\|\mathfrak{M}_{\alpha}(\vec{f}_j) - \mathfrak{M}_{\alpha}(\vec{f})\|_{L^q(\mathbb{R}^d)} \rightarrow 0 \quad \text{when } j \rightarrow \infty.$$

It suffices to show that

$$\|D_l \mathfrak{M}_{\alpha}(\vec{f}_j) - D_l \mathfrak{M}_{\alpha}(\vec{f})\|_{L^q(\mathbb{R}^d)} \rightarrow 0 \quad \text{when } j \rightarrow \infty \quad (14)$$

for any $l = 1, 2, \dots, d$. We only prove (14) for $l = d$ (since other cases are analogous). We may assume without loss of generality that all $f_i \geq 0$ and $f_{i,j} \geq 0$.

For any $i = 1, 2, \dots, m$, let $\vec{f}^i = (f_1, \dots, f_{i-1}, D_d f_i, f_{i+1}, \dots, f_m)$. Given $\varepsilon > 0$, there exists $R > 0$ such that $\sum_{i=1}^m \|\mathfrak{M}_{\alpha}(\vec{f}^i)\|_{q, G_1} < \varepsilon$ with $G_1 = \mathbb{R}^d \setminus B_R$. By absolute continuity, there exists $\eta > 0$ such that $\sum_{i=1}^m \|\mathfrak{M}_{\alpha}(\vec{f}^i)\|_{q, A} < \varepsilon$ whenever $|A| < \eta$ and A is a measurable subset of B_R . As we already observed, for a.e. $x \in \mathbb{R}^d$ and any $1 \leq i \leq m$, the function u_{x, \vec{f}^i} is uniformly continuous on $[0, \infty)$. Thus, for a.e. $x \in \mathbb{R}^d$, the function $\sum_{i=1}^m u_{x, \vec{f}^i}$ is uniformly continuous on $[0, \infty)$ and we can find $\delta_x > 0$ such that

$$\left| \sum_{i=1}^m u_{x, \vec{f}^i}(r_1) - \sum_{i=1}^m u_{x, \vec{f}^i}(r_2) \right| < |B_R|^{-1/q} \varepsilon \quad \text{whenever } |r_1 - r_2| < \delta_x.$$

It follows that there exists $\delta > 0$ such that

$$\begin{aligned} & \left| \left\{ x \in B_R : \left| \sum_{i=1}^m u_{x, \vec{f}_i}(r_1) - \sum_{i=1}^m u_{x, \vec{f}_i}(r_2) \right| \right. \right. \\ & \left. \left. \geq |B_R|^{-1/q} \varepsilon \text{ for some } r_1, r_2 \text{ with } |r_1 - r_2| < \delta \right\} \right| \\ & =: |G_2| < \frac{\eta}{2}. \end{aligned}$$

Applying Lemma 1, there exists $j_1 \in \mathbb{N}$ such that

$$|\{x \in B_R; \mathcal{R}(\vec{f}_j)(x) \not\subseteq \mathcal{R}(\vec{f})(x)_{(\delta)}\}| =: |B^j| < \frac{\eta}{2} \text{ when } j \geq j_1.$$

Let $\vec{f}_{i,j} = (f_{1,j}, \dots, f_{i-1,j}, D_d f_{i,j}, f_{i+1,j}, \dots, f_{m,j})$ and $r \in \mathcal{R}(\vec{f}_j)(x)$. We consider the following two cases: (i) $r > 0$. We can write

$$\begin{aligned} & |u_{x, \vec{f}_{i,j}}(r) - u_{x, \vec{f}_i}(r)| \\ & = \left| |B(x,r)|^{\alpha/d-m} \left(\prod_{1 \leq \mu \neq i \leq m} \int_{B(x,r)} f_{\mu,j}(y) dy \right) \int_{B(x,r)} D_d f_{i,j}(y) dy \right. \\ & \quad \left. - |B(x,r)|^{\alpha/d-m} \left(\prod_{1 \leq \mu \neq i \leq m} \int_{B(x,r)} f_{\mu}(y) dy \right) \int_{B(x,r)} D_d f_i(y) dy \right| \\ & \leq \sum_{\mu=1}^{i-1} \mathfrak{M}_{\alpha}(F_{\mu,j}^{\vec{}})(x) + \sum_{v=i+1}^m \mathfrak{M}_{\alpha}(G_{v,j}^{\vec{}})(x) + \mathfrak{M}_{\alpha}(H_{i,j}^{\vec{}})(x) =: \mathcal{G}_{i,j}(x), \end{aligned}$$

where

$$F_{\mu,j}^{\vec{}} = (f_1, \dots, f_{\mu-1}, f_{\mu,j} - f_{\mu}, f_{\mu+1,j}, \dots, f_{i-1,j}, D_d f_{i,j}, f_{i+1,j}, \dots, f_{m,j}),$$

$$G_{v,j}^{\vec{}} = (f_1, \dots, f_{i-1}, D_d f_i, f_{i+1}, \dots, f_{v-1}, f_{v,j} - f_v, f_{v+1,j}, \dots, f_{m,j}),$$

$$H_{i,j}^{\vec{}} = (f_1, \dots, f_{i-1}, D_d f_{i,j} - D_d f_i, f_{i+1,j}, \dots, f_{m,j}).$$

(ii) $r = 0$. If $0 < \alpha < md$, then $|u_{x, \vec{f}_{i,j}}(r) - u_{x, \vec{f}_i}(r)| = 0$. If $\alpha = 0$, we have

$$\begin{aligned} & |u_{x, \vec{f}_{i,j}}(r) - u_{x, \vec{f}_i}(r)| \\ & \leq \sum_{\mu=1}^{i-1} \left(\prod_{l_1=1}^{\mu-1} f_{l_1}(x) \right) (f_{\mu,j}(x) - f_{\mu}(x)) \left(\prod_{l_2=\mu+1}^{i-1} f_{l_2,j}(x) \right) |D_d f_{i,j}(x)| \left(\prod_{l_3=i+1}^m f_{l_3,j}(x) \right) \\ & \quad + \sum_{v=i+1}^m \left(\prod_{l_1=1}^{i-1} f_{l_1}(x) \right) |D_d f_i(x)| \left(\prod_{l_2=i+1}^{v-1} f_{l_2}(x) \right) |f_{v,j}(x) - f_v(x)| \left(\prod_{l_3=v+1}^m f_{l_3,j}(x) \right) \\ & \quad + \left(\prod_{l_1=1}^{i-1} f_{l_1}(x) \right) |D_d f_{i,j}(x) - D_d f_i(x)| \left(\prod_{l_2=i+1}^m f_{l_2,j}(x) \right). \end{aligned}$$

From the above, the Lebesgue differential theorem and Lemma 2, we have that for a.e.

$x \in \mathbb{R}^d$,

$$\begin{aligned}
& \left| D_d \mathfrak{M}_\alpha(\vec{f}_j)(x) - D_d \mathfrak{M}_\alpha(\vec{f})(x) \right| \\
&= \left| \sum_{i=1}^m u_{x, \vec{f}_i}(\vec{r}_1) - \sum_{i=1}^m u_{x, \vec{f}_i}(\vec{r}_2) \right| \\
&\leq \left| \sum_{i=1}^m u_{x, \vec{f}_i}(\vec{r}_1) - \sum_{i=1}^m u_{x, \vec{f}_i}(\vec{r}_1) \right| + \left| \sum_{i=1}^m u_{x, \vec{f}_i}(\vec{r}_1) - \sum_{i=1}^m u_{x, \vec{f}_i}(\vec{r}_2) \right| \\
&\leq \sum_{i=1}^m \mathcal{G}_{i,j}(x) + \left| \sum_{i=1}^m u_{x, \vec{f}_i}(\vec{r}_1) - \sum_{i=1}^m u_{x, \vec{f}_i}(\vec{r}_2) \right|
\end{aligned} \tag{15}$$

for any $r_1 \in \mathcal{R}(\vec{f}_j)(x)$ and $r_2 \in \mathcal{R}(\vec{f})(x)$. One can easily check that

$$\lim_{j \rightarrow \infty} \|\mathcal{G}_{i,j}\|_{L^q(\mathbb{R}^d)} = 0$$

for any $i = 1, 2, \dots, m$. It follows that there exists $j_2 \in \mathbb{N}$ such that $\|\mathcal{G}_{i,j}\|_{L^q(\mathbb{R}^d)} < \varepsilon$ for any $i = 1, 2, \dots, m$ and $j \geq j_2$.

If $x \notin G_1 \cup G_2 \cup B^j$ we can choose $r_1 \in \mathcal{R}(\vec{f}_j)(x)$ and $r_2 \in \mathcal{R}(\vec{f})(x)$ such that $|r_1 - r_2| < \delta$ and

$$\left| \sum_{i=1}^m u_{x, \vec{f}_i}(\vec{r}_1) - \sum_{i=1}^m u_{x, \vec{f}_i}(\vec{r}_2) \right| < |B_R|^{-1/q} \varepsilon.$$

On the other hand, we have that for any $i = 1, 2, \dots, m$, $r_1 \in \mathcal{R}(\vec{f}_j)(x)$ and $r_2 \in \mathcal{R}(\vec{f})(x)$,

$$\left| \sum_{i=1}^m u_{x, \vec{f}_i}(\vec{r}_1) - \sum_{i=1}^m u_{x, \vec{f}_i}(\vec{r}_2) \right| \leq \sum_{i=1}^m |u_{x, \vec{f}_i}(\vec{r}_1) - u_{x, \vec{f}_i}(\vec{r}_2)| \leq 2 \sum_{i=1}^m \mathfrak{M}_\alpha(\vec{f}_i)(x).$$

Note that $|G_2 \cup B^j| < \eta$ for $j \geq j_1$. Thus we get from (15) that

$$\begin{aligned}
& \|D_d \mathfrak{M}_\alpha(\vec{f}_j) - D_d \mathfrak{M}_\alpha(\vec{f})\|_{L^q(\mathbb{R}^d)} \\
&\leq \left\| \sum_{i=1}^m \mathcal{G}_{i,j} \right\|_{L^q(\mathbb{R}^d)} + \| |B_R|^{-1/q} \varepsilon \|_{q, B_R} + 2 \left\| \sum_{i=1}^m \mathfrak{M}_\alpha(\vec{f}_i) \right\|_{q, G_1 \cup G_2 \cup B^j} \leq C\varepsilon,
\end{aligned}$$

for any $j \geq \max\{j_1, j_2\}$, which gives

$$\lim_{j \rightarrow \infty} \|D_d \mathfrak{M}_\alpha(\vec{f}_j) - D_d \mathfrak{M}_\alpha(\vec{f})\|_{L^q(\mathbb{R}^d)} = 0.$$

This completes the proof of Theorem 1. \square

3. Proof of Theorem 2

This section is devoted to proving Theorem 2. We now recall the definition of approximate differentiability. Let f be a real-valued function defined on a set $E \subset$

\mathbb{R}^d . We say that f is approximately differentiable at $x_0 \in E$ if there is a vector $L = (L_1, L_2, \dots, L_d) \in \mathbb{R}^d$ such that for any $\varepsilon > 0$ the set

$$A_\varepsilon = \left\{ x \in \mathbb{R}^d : \frac{|f(x) - f(x_0) - L(x - x_0)|}{|x - x_0|} < \varepsilon \right\}$$

has x_0 as a density point. If this is the case, then x_0 is a density point of E and L is uniquely determined. The vector L is called the approximate differential of f at x_0 and is denoted by $\nabla f(x_0)$. Note that every function $f \in W^{1,1}(\mathbb{R}^d)$ is approximately differentiable a.e. It follows from Theorem A that Mf is approximately differentiable a.e. under the assumption that $f \in W^{1,1}(\mathbb{R}^d)$. However, it is unknown that whether $f \in W^{1,1}(\mathbb{R}^d)$ implies the weak differentiability of Mf when $d \geq 2$. For now, the relationship of approximate differentiability and weak differentiability is also not clear.

To prove Theorem 2, we need the following lemma followed from [35], which provides several characterizations of a.e. approximate differentiability of a function.

LEMMA 3. ([35]) *Let $f : E \rightarrow \mathbb{R}$ be measurable, $E \subset \mathbb{R}^d$. Then the following conditions are equivalent:*

- (i) *f is approximately differentiable a.e.*
- (ii) *For any $\varepsilon > 0$, there is a closed set $F \subset E$ and a locally Lipschitz function $g : \mathbb{R}^d \rightarrow \mathbb{R}$ such that $f = g$ on $x \in F$ and $|E \setminus F| < \varepsilon$.*
- (iii) *For any $\varepsilon > 0$, there is a closed set $F \subset E$ and a function $g \in \mathcal{C}^1(\mathbb{R}^d)$ such that $f = g$ on $x \in F$ and $|E \setminus F| < \varepsilon$.*

LEMMA 4. *Let $0 \leq \alpha \leq md - 1$ and $\vec{f} = (f_1, \dots, f_m)$ with each $f_j \in L^1(\mathbb{R}^d)$. For $\varepsilon > 0$, we define the truncated maximal operator $\mathfrak{M}_\alpha^\varepsilon$ by*

$$\mathfrak{M}_\alpha^\varepsilon(\vec{f})(x) = \sup_{r \geq \varepsilon} |B(x, r)|^{\alpha/d-m} \prod_{i=1}^m \int_{B(x, r)} |f_i(y)| dy.$$

Then $\mathfrak{M}_\alpha^\varepsilon(\vec{f})$ is Lipschitz continuous for every $\varepsilon > 0$.

Proof. For any $r \geq \varepsilon$ and $\delta \geq 1$, we have

$$\left(\frac{r}{r + |x - y|} \right)^\delta \geq \left(\frac{\varepsilon}{\varepsilon + |x - y|} \right)^\delta \geq 1 - \delta \frac{|x - y|/\varepsilon}{1 + |x - y|/\varepsilon} \geq 1 - \frac{\delta}{\varepsilon} |x - y|. \quad (16)$$

Fix $x, y \in \mathbb{R}^d$ and $r \geq \varepsilon$ we have $B(y, r) \subset B(x, r + |x - y|)$. This together with (16) yields that

$$\begin{aligned} \mathfrak{M}_\alpha^\varepsilon(\vec{f})(x) &\geq |B(x, r + |x - y|)|^{\alpha/d-m} \prod_{i=1}^m \int_{B(x, r + |x - y|)} |f_i(y)| dy \\ &\geq \left(\frac{r}{r + |x - y|} \right)^{md-\alpha} |B(y, r)|^{\alpha/d-m} \prod_{i=1}^m \int_{B(y, r)} |f_i(y)| dy \\ &\geq \left(1 - \frac{md-\alpha}{\varepsilon} |x - y| \right) |B(y, r)|^{\alpha/d-m} \prod_{i=1}^m \int_{B(y, r)} |f_i(y)| dy. \end{aligned}$$

whenever $\alpha \leq md - 1$. It follows that

$$\mathfrak{M}_\alpha^\varepsilon(\vec{f})(y) - \mathfrak{M}_\alpha^\varepsilon(\vec{f})(x) \leq \frac{md - \alpha}{\varepsilon} |x - y| \mathfrak{M}_\alpha^\varepsilon(\vec{f})(y).$$

Similarly,

$$\mathfrak{M}_\alpha^\varepsilon(\vec{f})(x) - \mathfrak{M}_\alpha^\varepsilon(\vec{f})(y) \leq \frac{md - \alpha}{\varepsilon} |x - y| \mathfrak{M}_\alpha^\varepsilon(\vec{f})(x).$$

Thus we have

$$\begin{aligned} |\mathfrak{M}_\alpha^\varepsilon(\vec{f})(x) - \mathfrak{M}_\alpha^\varepsilon(\vec{f})(y)| &\leq \frac{md - \alpha}{\varepsilon} |x - y| (\mathfrak{M}_\alpha^\varepsilon(\vec{f})(x) + \mathfrak{M}_\alpha^\varepsilon(\vec{f})(y)) \\ &\leq 2 \frac{md - \alpha}{\varepsilon} |x - y| \left(\frac{1}{\frac{2\pi^{d/2}}{d\Gamma(d/2)} \varepsilon^d} \right)^{m-\alpha/d} \prod_{j=1}^m \|f_j\|_{L^1(\mathbb{R}^d)} \\ &\leq 2 \frac{md - \alpha}{\varepsilon^{md-\alpha+1}} \left(\frac{d\Gamma(d/2)}{2\pi^{d/2}} \right)^{m-\alpha/d} \prod_{j=1}^m \|f_j\|_{L^1(\mathbb{R}^d)} |x - y|. \end{aligned}$$

Then Lemma 4 is proved. \square

We now turn to prove Theorem 2.

Proof of Theorem 2. Let Z_j be the set of all Lebesgue points of f_j and $u_{x,\vec{f}}(r)$ as in Section 2. We set $F = \mathbb{R}^d \setminus (\bigcap_{j=1}^m Z_j)$. Let $x \in \bigcap_{j=1}^m Z_j$ such that $\mathfrak{M}_\alpha(\vec{f})(x) > u_{x,\vec{f}}(0)$. Since $f_j \in L^1(\mathbb{R}^d)$ and $\mathfrak{M}_\alpha(\vec{f})(x) > 0$, there exists a sequence of positive bounded numbers $\{r_k\}_{k \geq 1}$ such that

$$u_{x,\vec{f}}(r_k) \rightarrow \mathfrak{M}_\alpha(\vec{f})(x) \text{ when } k \rightarrow \infty.$$

Hence there exists a subsequence $\{s_k\}_{k \geq 1} \subset \{r_k\}_{k \geq 1}$ such that $\lim_{k \rightarrow \infty} s_k = r > 0$. It follows that

$$\mathfrak{M}_\alpha(\vec{f})(x) = u_{x,\vec{f}}(r).$$

This yields that

$$\mathbb{R}^d = F \cup \{x \in \mathbb{R}^d : \mathfrak{M}_\alpha(\vec{f})(x) = u_{x,\vec{f}}(0)\} \cup E,$$

where $E = \bigcup_{k=1}^\infty E_k$ and $E_k = \{x \in \mathbb{R}^d : \mathfrak{M}_\alpha(\vec{f})(x) = \mathfrak{M}_\alpha^{1/k}(\vec{f})(x)\}$. Obviously, $E_k \subset E_{k+1}$. By Lemma 3, we know that $\prod_{j=1}^m |f_j|$ is approximately differentiable a.e. Then $\mathfrak{M}_\alpha(\vec{f})$ is approximately differentiable a.e. in the set $\{x \in \mathbb{R}^d : \mathfrak{M}_\alpha(\vec{f})(x) = u_{x,\vec{f}}(0)\}$. By Lemma 4 we have that $\mathfrak{M}_\alpha^{1/k}(\vec{f})$ is Lipschitz continuous for any $k \geq 1$. It follows that $\mathfrak{M}_\alpha(\vec{f})\chi_{E_{k+1} \setminus E_k}$ is approximately differentiable a.e. for all $k \geq 1$. Using Lemma 3 again we have that $\mathfrak{M}_\alpha(\vec{f})\chi_E = \mathfrak{M}_\alpha(\vec{f})\chi_{E_k} + \sum_{k=1}^\infty \mathfrak{M}_\alpha(\vec{f})\chi_{E_{k+1} \setminus E_k}$ is approximately differentiable a.e. Note that $|F| = 0$. Therefore, $\mathfrak{M}_\alpha(\vec{f})$ is approximately differentiable a.e. \square

REFERENCES

- [1] D. R. ADAMS, *A note on Riesz potentials*, Duke Math. J., **42**, (1975), 765–778.
- [2] D. R. ADAMS AND L. I. HEDBERG, *Function Spaces and Potential Theory*, Springer-Verlag, Berlin Heidelberg, 1996.
- [3] J. M. ALDAZ, L. COLZANI AND J. PÉREZ LÁZARO, *Optimal bounds on the modulus of continuity of the uncentered Hardy-Littlewood maximal function*, J. Geom. Anal., **22**, 1 (2012), 132–167.
- [4] J. M. ALDAZ AND J. PÉREZ LÁZARO, *Functions of bounded variation, the derivative of the one dimensional maximal function, and applications to inequalities*, Trans. Amer. Math. Soc., **359**, 5 (2007), 2443–2461.
- [5] J. M. ALDAZ AND J. PÉREZ LÁZARO, *Boundedness and unboundedness results for some maximal operators on functions of bounded variation*, J. Math. Anal. Appl., **337**, 1 (2008), 130–143.
- [6] J. M. ALDAZ AND J. PÉREZ LÁZARO, *Regularity of the Hardy-Littlewood maximal operator on block decreasing functions*, Studia Math., **194**, (2009), 253–277.
- [7] F. J. ALMGREN AND E. H. LIEB, *Symmetric decreasing rearrangement is sometimes continuous*, J. Amer. Math. Soc., **2**, (1989), 683–773.
- [8] M. CAO AND Q. XUE, *Characterization of two-weighted inequalities for multilinear fractional maximal operator*, Non. Anal. TMA, **130**, (2016), 214–228.
- [9] M. CAO AND Q. XUE, *Two-weight entropy boundedness of multilinear fractional type operators*, J. Math. Inequa., **11**, 2 (2017), 441–453.
- [10] E. CARNEIRO AND J. MADRID, *Derivative bounds for fractional maximal functions*, Trans. Amer. Math. Soc., to appear.
- [11] E. CARNEIRO AND D. MOREIRA, *On the regularity of maximal operators*, Proc. Amer. Math. Soc., **136**, 12 (2008), 4395–4404.
- [12] E. CARNEIRO AND B. F. SVAITER, *On the variation of maximal operators of convolution type*, J. Funct. Anal., **265**, (2013), 837–865.
- [13] P. HAJŁASZ AND J. MALY, *On approximate differentiability of the maximal function*, Proc. Amer. Math. Soc., **138**, 1 (2010), 165–174.
- [14] P. HAJŁASZ AND J. ONNINEN, *On boundedness of maximal functions in Sobolev spaces*, Ann. Acad. Sci. Fenn. Math., **29**, 1 (2004), 167–176.
- [15] J. KINNUNEN, *The Hardy-Littlewood maximal function of a Sobolev function*, Israel J. Math., **100**, (1997), 117–124.
- [16] J. KINNUNEN AND P. LINDQVIST, *The derivative of the maximal function*, J. Reine. Angew. Math., **503**, (1998), 161–167.
- [17] J. KINNUNEN AND E. SAKSMAN, *Regularity of the fractional maximal function*, Bull. London Math. Soc., **35**, 4 (2003), 529–535.
- [18] O. KURKA, *On the variation of the Hardy-Littlewood maximal function*, Ann. Acad. Sci. Fenn. Math., **40**, (2015), 109–133.
- [19] A. K. LERNER, S. OMBROSI, C. PÉREZ, R. H. TORRES AND R. TRUJILLO-GONZÁLEZ, *New maximal functions and multiple weightes for the multilinear Calderón-Zygmund theory*, Adv. Math., **220**, (2009), 1222–1264.
- [20] K. LI, K. MONE AND W. SUN, *Sharp weighted inequalities for multilinear fractional maximal operator and fractional integrals*, Math. Nachr., **288**, (5-6) (2015), 619–632.
- [21] K. LI AND W. SUN, *Characterization of a two weight inequality for multilinear fractional maximal operators*, Houston J. Math., **42**, 3 (2016), 977–990.
- [22] F. LIU, *A remark on the regularity of the discrete maximal operator*, Bull. Austral. Math. Soc., **95**, (2017), 108–120.
- [23] F. LIU, T. CHEN AND H. WU, *A note on the endpoint regularity of the Hardy-Littlewood maximal functions*, Bull. Austral. Math. Soc., **94**, (2016), 121–130.
- [24] F. LIU AND S. MAO, *On the regularity of the one-sided Hardy-Littlewood maximal functions*, Czech. Math. J., **67**, 142 (2017), 219–234.
- [25] F. LIU AND H. WU, *On the regularity of the multisublinear maximal functions*, Canad. Math. Bull., **58**, 4 (2015), 808–817.
- [26] F. LIU AND H. WU, *Endpoint regularity of multisublinear fractional maximal functions*, Canad. Math. Bull., to appear. DOI:10.4153/CMB-2016-044-9.

- [27] F. LIU AND H. WU, *Regularity of discrete multisublinear fractional maximal functions*, Sci. China Math., to appear. DOI:10.1007/s11425-016-9011-2.
- [28] F. LIU AND H. WU, *On the regularity of maximal operators supported by submanifolds*, J. Math. Anal. Appl., **453**, (2017), 144–158.
- [29] H. LUIRO, *Continuity of the maximal operator in Sobolev spaces*, Proc. Amer. Math. Soc., **135**, 1 (2007), 243–251.
- [30] H. LUIRO, *On the regularity of the Hardy-Littlewood maximal operator on subdomains of \mathbb{R}^n* , Proc. Edinburgh Math. Soc., **53**, 1 (2010), 211–237.
- [31] H. LUIRO AND J. NUUTINEN, *On the continuity of discrete maximal operators in Sobolev spaces*, Ann. Acad. Sci. Fenn. Math., **39**, (2014), 175–185.
- [32] B. MUCKENHOUT AND R. L. WHEEDEN, *Weighted norm inequalities for fractional integrals*, Trans. Amer. Math. Soc., **192**, (1974), 261–274.
- [33] C. PÉREZ AND R. WHEEDEN, *Potential operators, maximal functions, and generalizations of A_∞* , Pot. Anal., **19**, 1 (2003), 1–33.
- [34] H. TANAKA, *A remark on the derivative of the one-dimensional Hardy-Littlewood maximal function*, Bull. Austr. Math. Soc., **65**, 2 (2002), 253–258.
- [35] H. WHITNEY, *On totally differentiable and smooth functions*, Pacific. J. Math., **1**, (1951), 143–159.

(Received February 16, 2017)

Feng Liu
College of Mathematics and Systems Science
Shandong University of Science and Technology
Qingdao, Shandong 266590, P. R. China
e-mail: liufeng860314@163.com