

## A NONEXISTENCE RESULT FOR DISCRETE SYSTEMS RELATED TO THE REVERSED HARDY–LITTLEWOOD–SOBOLEV INEQUALITY

TING TANG

(Communicated by S. Varošanec)

*Abstract.* In this paper, we study the Euler-Lagrange system related to the extremal sequences of the discrete reversed Hardy-Littlewood-Sobolev inequality. This system comes into play in estimating bounds of the Coulomb energy and is associated with the study of conformal geometry. By estimating the infinite series, we prove that the nonexistence of super-solutions of the Euler-Lagrange system satisfied by the extremal sequences of this discrete reversed Hardy-Littlewood-Sobolev inequality.

### 1. Introduction

The Hardy-Littlewood-Sobolev (HLS) inequality is read as (cf. [17])

$$\left| \int_{R^n} \int_{R^n} \frac{f(x)g(y)dxdy}{|x-y|^\lambda} \right| \leq C \|f\|_{L^r} \|g\|_{L^s}, \quad \forall (f, g) \in L^r(R^n) \times L^s(R^n), \quad (1)$$

where  $0 < \lambda < n$ ,  $\min\{r, s\} > 1$  and  $\frac{1}{r} + \frac{1}{s} + \frac{\lambda}{n} = 2$ . In 1983, Lieb proved the existence of extremal functions of (1), and obtained the Euler-Lagrange system (cf. [15])

$$\begin{cases} u(x) = \int_{R^n} \frac{v^q(y)dy}{|x-y|^\lambda}, & u > 0 \text{ in } R^n, \\ v(x) = \int_{R^n} \frac{u^p(y)dy}{|x-y|^\lambda}, & v > 0 \text{ in } R^n, \end{cases} \quad (2)$$

where

$$u = c_1 f^{r-1}, v = c_2 g^{s-1}, p = \frac{1}{r-1}, q = \frac{1}{s-1}, \quad (3)$$

and  $c_1, c_2$  are proper constants. Now,  $\frac{1}{r} + \frac{1}{s} + \frac{\lambda}{n} = 2$  becomes the critical condition of Sobolev type

$$\frac{1}{p+1} + \frac{1}{q+1} = \frac{\lambda}{n}. \quad (4)$$

*Mathematics subject classification* (2010): 26D15, 40B05, 47J20.

*Keywords and phrases:* Discrete reversed Hardy-Littlewood-Sobolev inequality, Euler-Lagrange equation, nonexistence of positive solution.

In 2015, Dou and Zhu [6] proved a reversed Hardy-Littlewood-Sobolev (RHLS) inequality (see also [16])

$$\left| \int_{R^n} \int_{R^n} \frac{f(x)g(y)dxdy}{|x-y|^\lambda} \right| \geq C \|f\|_{L^r} \|g\|_{L^s}, \quad \forall (f, g) \in L^r(R^n) \times L^s(R^n), \quad (5)$$

and obtained the existence of extremal functions, where  $n \geq 1, \lambda < 0, \frac{n}{n-\lambda} < r, s < 1$  and  $\frac{1}{r} + \frac{1}{s} + \frac{\lambda}{n} = 2$ . The Euler-Lagrange system which the extremal functions satisfy is still (2) with (3).

When  $f \equiv g$  and  $r = s$  in (1), Lieb classified the extremal functions and thus obtained the best constant (cf. [15]). Meanwhile, when  $p = q$  and  $u \equiv v$ , (2) is reduced to

$$u(x) = \int_{R^n} \frac{u^p(y)dy}{|x-y|^\lambda}, \quad u > 0 \quad \text{in } R^n. \quad (6)$$

This equation is related to the study of the conformal geometry and the nonlinear elliptic PDEs. Lieb posed a problem in [15] - whether the solutions of the integral equation (6) still has the classification result. When  $0 < \lambda < n$  and  $p = 2n/\lambda - 1$ , Chen, Li, Ou [4] and Li [13] classified the positive solution  $u \in L^{2n/\lambda}_{loc}(R^n)$  and thus answered independently the question. Namely,  $u$  must be of the form

$$u(x) = a(b^2 + |x - x_0|^2)^{-\lambda/2} \quad (7)$$

with  $a, b > 0$  and  $x_0 \in R^n$ . When  $0 < \lambda < n$  and  $p > 0$ , Chen, Li and Ou obtained the results of the existence/nonexistence, the radial symmetry, the integrability of the solutions (cf. [2]), and the symmetry of the components (cf. [1]). When  $\lambda < 0$ , Li studied (6) with the negative exponent  $\frac{2n-\lambda}{\lambda} \leq p < 0$  (cf. [13]), and proved that  $p = \frac{2n-\lambda}{\lambda}$  and  $u$  is classified as the form (7). A problem posed by Li is whether or not does (6) admit any positive (regular) solutions for all  $n \geq 1, \lambda < 0$  and  $p < (2n - \lambda)/\lambda$ . Xu answered this question and obtained the following results (cf. [20]).

(Ri) Let  $\lambda < 0$  and  $p < 0$ . Eq. (6) has a positive solution if and only if  $2n - \lambda = p\lambda$ . Now,  $u$  is given by (7).

(Rii) If  $0 < \lambda < n$  and  $p < 0$ , then (6) has no positive solution.

Let  $(u, v) \in L^{p+1}(R^n) \times L^{q+1}(R^n)$  be a pair of positive solutions of (2),  $\lambda \in (0, n)$  and  $\min\{p, q\} > 1$  satisfy (4). Chen, Li and Ou proved  $u$  and  $v$  are radially symmetric and decreasing about some point by the method of moving planes (cf. [3]). Afterwards, the optimal integrability intervals of  $u$  and  $v$  are obtained (cf. [8]). Based on these results, the decay rates of  $u$  and  $v$  were estimated in [11] and [18].

When  $\lambda < 0$  and  $\min\{p, q\} < (n - \lambda)/\lambda$ , paper [9] studied (2). The author proved that both  $u^{-1}$  and  $v^{-1}$  decay to zero with rate  $\lambda$ , and  $u^{-1}, v^{-1} \in L^s(R^n)$  for all  $s > n/\lambda$ . Moreover, [9] pointed out that (4) is a necessary condition for the existence of positive solutions of (2). Dou and Zhu [6] obtained the existence of extremal functions of (5), which implies that the (4) is still the sufficient condition for the existence of positive solutions of (2). In addition, they also obtained the classification result of (Ri) by the method of moving spheres developed by Li and Zhu in [14].

Eq. (4) can also be viewed as the necessary and sufficient condition ensuring the conformal invariant. Namely, the system (2) and the energy functionals  $\|u\|_{p+1}$

and  $\|v\|_{q+1}$  are invariant under the scaling transformation (see [10] for  $\lambda \in (0, n)$  and  $\min\{p, q\} > 0$ , and [9] for  $\lambda < 0$  and  $\max\{p, q\} < 0$ ).

In 2015, Huang, Li and Yin employed (1) to prove the discrete Hardy-Littlewood-Sobolev (DHLS) inequality (cf. [7])

$$\left| \sum_{i,j,i \neq j} \frac{f_i g_j}{|i-j|^\lambda} \right| \leq C \|f\|_{l^r} \|g\|_{l^s}, \quad \forall (f, g) \in l^r(\mathbb{Z}^n) \times l^s(\mathbb{Z}^n), \tag{8}$$

where  $f = (f_i)_{i \in \mathbb{Z}^n}$ ,  $g = (g_j)_{j \in \mathbb{Z}^n}$ ,  $n \geq 1$ ,  $0 < \lambda \leq n$ ,  $\min\{r, s\} > 1$  and  $\frac{1}{r} + \frac{1}{s} + \frac{\lambda}{n} \geq 2$ . When  $n = \lambda = 1$ , it is the classical Hardy-Littlewood-Polya inequality. They pointed out in [7] that there exist extremal sequences when  $\frac{1}{r} + \frac{1}{s} + \frac{\lambda}{n} > 2$ , and conjectured that it is also true when  $\frac{1}{r} + \frac{1}{s} + \frac{\lambda}{n} = 2$ . More results on the best constants and the extremal sequences of the inequality with finite elements can be seen in [5], [12] and [19].

The following discrete reversed Hardy-Littlewood-Sobolev inequality can be established as the same argument in [7]

$$\|f\|_{l^r} \|g\|_{l^s} \leq C \left( \sum_{i,j \in \mathbb{Z}^n} \frac{|f_i| |g_j|}{|i-j|^\lambda} + \sum_{j \in \mathbb{Z}^n} |f_j| |g_j| \right), \quad \forall (f, g) \in l^r(\mathbb{Z}^n) \times l^s(\mathbb{Z}^n), \tag{9}$$

where  $f = (f_i)_{i \in \mathbb{Z}^n}$ ,  $g = (g_j)_{j \in \mathbb{Z}^n}$ ,

$$\lambda < 0, \quad \frac{n}{n-\lambda} < r, s < 1, \quad \frac{1}{r} + \frac{1}{s} + \frac{\lambda}{n} \leq 2. \tag{10}$$

The Euler-Lagrange system satisfied by the extremal sequences is

$$\begin{cases} u_j = \sum_{k \in \mathbb{Z}^n} \frac{v_k^q}{|k-j|^\lambda} + v_j^q, \\ v_j = \sum_{k \in \mathbb{Z}^n} \frac{u_k^p}{|k-j|^\lambda} + u_j^p, \end{cases} \tag{11}$$

where  $u_j, v_j > 0$  ( $j \in \mathbb{Z}^n$ ), and  $\lambda < 0$ ,  $\max\{p, q\} < 0$ .

In this paper, we are concerned with nonexistence of positive solution of the following system

$$\begin{cases} u_j = c_1(j) \left( \sum_{k \in \mathbb{Z}^n, k \neq j} \frac{v_k^q}{|k-j|^\lambda} + v_j^q \right), \\ v_j = c_2(j) \left( \sum_{k \in \mathbb{Z}^n, k \neq j} \frac{u_k^p}{|k-j|^\lambda} + u_j^p \right), \end{cases} \tag{12}$$

with double bounded sequences  $c_i(j)$  ( $i = 1, 2$  and  $j \in \mathbb{Z}^n$ ). A sequence  $c(j)$  is called double bounded, if there exists  $C > 1$  such that  $C^{-1} \leq c(j) \leq C$  for all  $j \in \mathbb{Z}^n$ . Clearly, the nonexistence of positive solution of (12) implies the nonexistence of positive solution of (11).

The following theorem is the corresponding result of (Rii) to some extent.

**THEOREM 1.** *When  $0 < \lambda < n$  and  $\min\{p, q\} < 0$ , then for any double bounded sequences  $c_1(j)$  and  $c_2(j)$ , there does not exist solution of (12) satisfying  $u_j \simeq |j|^{\theta_1}$  and  $v_j \simeq |j|^{\theta_2}$  for all  $\theta_1, \theta_2 \in \mathbb{R}$ .*

Here  $u_j \simeq |j|^\theta$  means that there exists  $C > 0$  such that

$$\frac{|j|^\theta}{C} \leq u_j \leq C|j|^\theta$$

when  $|j| \rightarrow \infty$ .

### 2. Proof of Theorem 1

*Proof.* Without loss of generality, we assume  $q \leq p$ . Thus,  $q < 0$ . We prove Theorem 1 by contradiction.

Suppose that there exist  $\theta_1, \theta_2 \in R$  such that the solutions  $u, v$  of (12) satisfy

$$u(x) \simeq |x|^{\theta_1}, \quad v(x) \simeq |x|^{\theta_2}. \tag{13}$$

Therefore, from (12) it follows

$$u_j \geq c \sum_{|k| \geq 2|j|} \frac{v_k^q}{|k-j|^\lambda} \geq c \sum_{|k| \geq 2|j|} |k|^{-\lambda+q\theta_2},$$

which implies

$$n - \lambda + q\theta_2 < 0. \tag{14}$$

Similarly, we can also obtain

$$n - \lambda + p\theta_1 < 0. \tag{15}$$

Noting  $\lambda \in (0, n)$  and  $q < 0$ , we see from (14) and (15) that

$$\theta_2 > 0, \tag{16}$$

and

$$p\theta_1 < 0. \tag{17}$$

Clearly, from (12) there holds

$$v_j := \sum_{m=1}^5 T_m,$$

where

$$\begin{aligned} T_1 &:= \sum_{|k-j| \leq |j|/2} \frac{u_k^p}{|k-j|^\lambda} \\ T_2 &:= \sum_{|k| \geq 2|j|} \frac{u_k^p}{|k-j|^\lambda} \\ T_3 &:= \sum_{|k-j| \geq |j|/2, 2 \leq |k| \leq 2|j|} \frac{u_k^p}{|k-j|^\lambda} \\ T_4 &:= \sum_{|k| \leq 2} \frac{u_k^p}{|k-j|^\lambda} \\ T_5 &:= u_j^p. \end{aligned}$$

When  $|k - j| \leq |j|/2$ , there holds  $|j|/2 \leq |k| \leq 3|j|/2$ . Thus by (13),

$$T_1 \leq C|j|^{p\theta} \sum_{|k-j| \leq |j|/2} \frac{1}{|k|^\lambda} \leq C|j|^{n-\lambda+p\theta_1}.$$

When  $|k| \geq 2|j|$ , there holds  $|k - j| \geq |k|/2$ . Thus by (13) and (15), we have

$$T_2 \leq C \sum_{|k| \geq 2|j|} \frac{1}{|k|^{\lambda-p\theta_1}} \leq C|j|^{n-\lambda+p\theta_1}.$$

When  $|k - j| \geq |j|/2$  and  $2 \leq |k| \leq 2|j|$ , we have

$$T_3 \leq \frac{C}{|j|^\lambda} \sum_{|k-j| \geq |j|/2, 2 \leq |k| \leq 2|j|} |k|^{p\theta_1}.$$

Therefore,

$$T_3 \leq \begin{cases} C|j|^{n-\lambda+p\theta_1}, & \text{when } n + p\theta_1 > 0; \\ C|j|^{-\lambda} \log |j|, & \text{when } n + p\theta_1 = 0; \\ C|j|^{-\lambda}, & \text{when } n + p\theta_1 < 0. \end{cases}$$

When  $|k| \leq 2$ ,

$$T_4 \leq C|j|^{-\lambda}.$$

At last, by (13),

$$T_5 \leq C|j|^{p\theta_1}.$$

Therefore,

$$v_j \leq C \left( |j|^{n-\lambda+p\theta_1} + |j|^{-\lambda} \log |j| + |j|^{-\lambda} + |j|^{p\theta_1} \right).$$

Comparing this with (13), we get from (15) and (17) that

$$\theta_2 \leq \max\{n - \lambda + p\theta_1, -\lambda + \varepsilon, p\theta_1\} < 0.$$

Here  $\varepsilon > 0$  is sufficiently small. This contradicts with (16). Theorem 1 is proved.

*Acknowledgements.* This research was supported by NNSF (11871278) of China and Natural Science Foundation of Jiangsu Education Commission (19KJB110016).

#### REFERENCES

- [1] W. CHEN, C. LI, *Regularity of solutions for a system of integral equations*, Commun. Pure Appl. Anal., **4** (2005), 1–8.
- [2] W. CHEN, C. LI, B. OU, *Qualitative properties of solutions for an integral equation*, Discrete Contin. Dyn. Syst., **12** (2005), 347–354.
- [3] W. CHEN, C. LI, B. OU, *Classification of solutions for a system of integral equations*, Comm. Partial Differential Equations, **30** (2005), 59–65.

- [4] W. CHEN, C. LI, B. OU, *Classification of solutions for an integral equation*, Comm. Pure Appl. Math., **59** (2006), 330–343.
- [5] Z. CHENG, C. LI, *An extended discrete Hardy-Littlewood-Sobolev inequality*, Discrete Contin. Dyn. Syst., **34** (2014), 1951–1959.
- [6] J. DOU, M. ZHU, *Reversed Hardy-Littlewood-Sobolev inequality*, Internat. Math. Res. Notices, (2015), no.19, 9696–9726.
- [7] G. HUANG, C. LI, X. YIN, *Existence of the maximizing pair for the discrete Hardy-Littlewood-Sobolev inequality*, Discrete Contin. Dyn. Syst., **35** (2015), 935–942.
- [8] C. JIN, C. LI, *Qualitative analysis of some systems of integral equations*, Calc. Var. Partial Differential Equations, **26** (2006), 447–457.
- [9] Y. LEI, *On the integral systems with negative exponents*, Discrete Contin. Dyn. Syst., **35** (2015), 1039–1057.
- [10] Y. LEI, C. LI, *Sharp criteria of Liouville type for some nonlinear systems*, Discrete Contin. Dyn. Syst., **36** (2016), 3277–3315.
- [11] Y. LEI, C. LI, C. MA, *Asymptotic radial symmetry and growth estimates of positive solutions to weighted Hardy-Littlewood-Sobolev system*, Calc. Var. Partial Differential Equations, **45** (2012), 43–61.
- [12] C. LI, J. VILLAVERTE, *An extension of the Hardy-Littlewood-Polya inequality*, Acta Math. Scientia, **31(B)** (2011), 2285–2288.
- [13] Y. LI, *Remark on some conformally invariant integral equations: the method of moving spheres*, J. Eur. Math. Soc., **6** (2004), 153–180.
- [14] Y. LI AND M. ZHU, *Uniqueness theorems through the method of moving spheres*, Duke Math. J., **80** (1995), 383–417.
- [15] E. LIEB, *Sharp constants in the Hardy-Littlewood-Sobolev and related inequalities*, Ann. of Math., **118** (1983), 349–374.
- [16] Q. NGO, V. NGUYEN, *Sharp reversed Hardy-Littlewood-Sobolev inequality on  $R^n$* , Israel J. Math., **220** (2017), 1–35.
- [17] E. M. STEIN, *Singular integrals and differentiability properties of function*, Princeton Math. Series, Vol. 30, Princeton University Press, Princeton, NJ, 1970.
- [18] S. SUN, Y. LEI, *Fast decay estimates for integrable solutions of the Lane-Emden type integral systems involving the Wolff potentials*, J. Funct. Anal., **263** (2012), 3857–3882.
- [19] J. XU, Z. CHENG, Y. FANG, *An extension of discrete weighted Hardy-Littlewood-Sobolev inequality in space dimension one*, Sci. Sin. Math., **45** (2015), 129–140.
- [20] X. XU, *Uniqueness theorem for integral equations and its application*, J. Funct. Anal., **247** (2007), 95–109.

(Received June 19, 2019)

Ting Tang  
 Institute of Mathematics, School of Mathematical Sciences  
 Nanjing Normal University  
 Nanjing, 210023, China  
 e-mail: 941059734@qq.com