

EXTENSION FORMULAS AND NORM INEQUALITIES IN SOBOLEV HILBERT SPACES

SABUROU SAITOH

(Communicated by I. Perić)

Abstract. In this note, we shall consider extension formulas and norm inequalities for some typical Sobolev Hilbert spaces. We see many related open problems.

1. Introduction

In order to give the background for the results in this note, we first recall the fundamental general property of reproducing kernel Hilbert spaces based on [5].

We consider a positive definite quadratic form function $K : E \times E \rightarrow \mathbb{C}$ and its restriction of K to $E_0 \times E_0$, where E_0 is a subset of E . Of course, the restriction is again a positive definite quadratic form function on the subset $E_0 \times E_0$. The relation between two reproducing kernel Hilbert spaces derived from the positive definite quadratic form functions is given by the following theorem.

THEOREM A. ([5], page 78) *Let E_0 be a subset of E . Then the reproducing kernel Hilbert space that $K|_{E_0 \times E_0} : E_0 \times E_0 \rightarrow \mathbb{C}$ defines is given by:*

$$H_{K|_{E_0 \times E_0}}(E_0) = \{f \in \mathcal{F}(E_0) : f = \tilde{f}|_{E_0} \text{ for some } \tilde{f} \in H_K(E)\}. \quad (1)$$

Furthermore, the norm is expressed in terms of the one of $H_K(E)$:

$$\|f\|_{H_{K|_{E_0 \times E_0}}(E_0)} = \min\{\|\tilde{f}\|_{H_K(E)} : \tilde{f} \in H_K(E), f = \tilde{f}|_{E_0}\}. \quad (2)$$

In Theorem A, note that the inequality, for any function $f \in H_K(E)$

$$\|f\|_{H_{K|_{E_0 \times E_0}}(E_0)} \leq \|f\|_{H_K(E)} \quad (3)$$

holds, that is, the restriction map is a bounded linear operator.

At first, we shall consider the simplest Sobolev Hilbert space.

Mathematics subject classification (2020): 30C40, 44A05, 35A22, 46E22, 45A05.

Keywords and phrases: Extension and restriction of a function, Sobolev space, Fourier transform, reproducing kernel, bounded linear operator, norm estimate, norm inequality.

The author wishes to express his deep thanks Professors Y. Sawano, M. Seto and A. Yamada for their great contributions to the meeting: RIMS Workshop: Research on Real, Complex and Functional Analysis from the Perspective of Function Spaces, (Oct. 18th.–20th., 2023) and the paper. This meeting and paper were supported by the RIMS. Furthermore, the author expresses deeply the referee for very careful readings and valuable comments.

The space $H_S(\mathbb{R})$ is comprising of absolutely continuous functions f on \mathbb{R} with the norm

$$\|f\|_{H_S(\mathbb{R})} \equiv \sqrt{\int_{\mathbb{R}} (|f(x)|^2 + |f'(x)|^2) dx}. \quad (4)$$

The Hilbert space $H_S(\mathbb{R})$ admits the reproducing kernel

$$K(x, y) \equiv \frac{1}{2\pi} \int_{\mathbb{R}} \frac{1}{1 + \xi^2} \exp(i(x - y)\xi) d\xi = \frac{1}{2} e^{-|x-y|} \quad (x, y \in \mathbb{R}). \quad (5)$$

Its restriction to the closed interval $[a, b]$ is the reproducing kernel Hilbert space $H_S[a, b] = W^{1,2}[a, b]$ as a set of functions, and the norm is given by

$$\|f\|_{H_S[a,b]} \equiv \sqrt{\left(\int_a^b (|f(x)|^2 + |f'(x)|^2) dx \right) + |f(a)|^2 + |f(b)|^2} \quad (6)$$

([5], pages 10–16).

The representation (5) means that the functions $f(x)$ of $H_S(\mathbb{R})$ are represented in the form

$$f(x) = \frac{1}{2\pi} \int_{\mathbb{R}} \frac{1}{1 + \xi^2} \exp(ix\xi) F(\xi) d\xi$$

with the functions $F(\xi)$ satisfying

$$\frac{1}{2\pi} \int_{\mathbb{R}} \frac{1}{1 + \xi^2} |F(\xi)|^2 d\xi < \infty$$

and the norm is represented by

$$\|f\|_{H_S(\mathbb{R})} = \sqrt{\frac{1}{2\pi} \int_{\mathbb{R}} \frac{1}{1 + \xi^2} |F(\xi)|^2 d\xi}.$$

The restriction mapping L from the space $H_S(\mathbb{R})$ to the space $H_S[a, b]$ is, of course, not injective and so, in particular, we obtain the norm inequality

$$\|f\|_{H_S(\mathbb{R})} \geq \|f\|_{H_S[a,b]};$$

that is,

$$\int_{\mathbb{R}} (|f(x)|^2 + |f'(x)|^2) dx \geq \left(\int_a^b (|f(x)|^2 + |f'(x)|^2) dx \right) + |f(a)|^2 + |f(b)|^2. \quad (7)$$

By our general theory, we can give the precise correspondence of the two spaces; that is,

$$f|_{[a,b]}(x) = (f(\xi), K(\xi, x))_{H_S(\mathbb{R})} \quad (8)$$

and

$$f(x) = (f|_{[a,b]}(\xi), K(\xi, x))_{H_S[a,b]}, \quad (9)$$

with the minimum extension f of $f_{[a,b]}$ in $H_S([a, b])$ to $H_S(\mathbb{R})$. Indeed, we can derive directly the identity (9) for the minimum extension f of $f_{[a,b]}$ in $H_S[a, b]$ to $H_S(\mathbb{R})$. See the following proof of Theorem 1 for the space $W^{2,2}(\mathbb{R})$.

However, for the minimum extension formula we have the general formula in Theorem A,

$$f(p) = (f|E_0(\cdot), K(\cdot, p))_{H_{K|E_0 \times E_0}(E_0)},$$

for the minimum extension f of $f|E_0$. See the proof of Proposition 2.5 in [5] (pages 79-80), in particular, (2.48). There, we see some general and abstract proof, however, for the proof of Theorem 1 for the space $W^{2,2}(\mathbb{R})$, we can see the constructive proof for the minimum norm function.

We obtained several realizations of restricted reproducing kernel Hilbert spaces as in (6), however, they are, in general, involved. See [4], [5]. The formula (6) is a simple result, however, the realization of the restricted reproducing kernel spaces is, in general, complicated in this sense.

OPEN PROBLEM. Let $m > \frac{n}{2}$ be an integer. Denote by ${}_m C_\nu$ the binomial coefficient and by $W^{m,2}(\mathbb{R}^n)$ the Sobolev space whose norm is given by

$$\|F\|_{W^{m,2}(\mathbb{R}^n)} = \sqrt{\sum_{\nu=0}^m {}_m C_\nu \left(\sum_{\alpha \in \mathbb{Z}_+^n, |\alpha|=\nu} \frac{\nu!}{\alpha!} \int_{\mathbb{R}^n} \left| \frac{\partial^\alpha F(x)}{\partial x^\alpha} \right|^2 dx \right)}. \tag{10}$$

Then, the reproducing kernel K is given by

$$K(x, y) \equiv \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} \frac{\exp(i(x-y) \cdot \xi)}{(1 + |\xi|^2)^m} d\xi \quad (x, y \in \mathbb{R}^n) \tag{11}$$

[5], page 22). How will be the realization of the norm for the restricted reproducing kernel Hilbert space to some nontrivial subset (the typical case is a sphere $\{r < a\}$) of \mathbb{R}^n as in the case (6) of one dimensional way?

2. The typical case for the space $W^{2,2}(\mathbb{R})$

For the Sobolev Hilbert space $W^{2,2}(\mathbb{R})$ defined to be the completion of $C_c^\infty(\mathbb{R})$ with respect to the norm:

$$\|f\|_{W^{2,2}(\mathbb{R})} = \sqrt{\|f''\|_{L^2(\mathbb{R})}^2 + 2\|f'\|_{L^2(\mathbb{R})}^2 + \|f\|_{L^2(\mathbb{R})}^2},$$

we have the reproducing kernel

$$G(s, t) \equiv \frac{1}{4} e^{-|s-t|} (1 + |s-t|) \quad (s, t \in \mathbb{R})$$

[5], page 21–22).

For simplicity, in this section, we shall consider functions in real valued functions.

In order to look for the reproducing kernel Hilbert space $W_S([a, b])$, ($a < b$) admitting the restricted reproducing kernel $G(s, t)$ to the interval $[a, b]$, we calculate the integral, for any function $f \in W^{2,2}(\mathbb{R})$

$$(f(s), G(s, t))_{W^{2,2}([a, b])}.$$

By setting $G_t(s) = G(s, t)$, we note that

$$\begin{aligned} G_t(s) &= \frac{t-s+1}{4} \exp(s-t)\chi_{(a,t)}(s) + \frac{s-t+1}{4} \exp(t-s)\chi_{[t,b)}(s), \\ \frac{dG_t}{ds}(s) &= \frac{t-s}{4} \exp(s-t)\chi_{(a,t)}(s) - \frac{s-t}{4} \exp(t-s)\chi_{[t,b)}(s), \\ \frac{d^2G_t}{ds^2}(s) &= \frac{t-s-1}{4} \exp(s-t)\chi_{(a,t)}(s) + \frac{s-t-1}{4} \exp(t-s)\chi_{[t,b)}(s). \end{aligned} \tag{12}$$

Then, by integration by parts repeatedly, we have

$$\begin{aligned} &(f(s), G(s, t))_{W^{2,2}([a, b])} \\ &= f(t) + f(a) \frac{-t+a-2}{4} \exp(a-t) - f'(a) \frac{t-a-1}{4} \exp(a-t) \\ &\quad - f(b) \frac{b-t+2}{4} \exp(t-b) + f'(b) \frac{b-t-1}{4} \exp(t-b). \end{aligned} \tag{13}$$

That is

$$\begin{aligned} &(f(s), G(s, t))_{W^{2,2}([a, b])} \\ &= f(t) + f(a)G(a, t) \left(-1 + \frac{-1}{1+t-a} \right) + f'(a)G'(a, t) \left(\frac{1}{t-a} - 1 \right) \\ &\quad + f(b)G(b, t) \left(-1 + \frac{-1}{1+b-t} \right) + f'(b)G'(b, t) \left(\frac{1}{b-t} - 1 \right). \end{aligned} \tag{14}$$

We thus have the desired identity admitting the restricted reproducing kernel of $G(s, t)$ to the interval $[a, b]$

$$\begin{aligned} (f, G(\cdot, t))_{W_S([a, b])} &= (f, G(\cdot, t))_{W^{2,2}(\mathbf{R})} - f(a) \frac{-t+a-2}{4} \exp(a-t) \\ &\quad + f'(a) \frac{t-a-1}{4} \exp(a-t) + f(b) \frac{b-t+2}{4} \exp(t-b) \\ &\quad - f'(b) \frac{b-t-1}{4} \exp(t-b). \end{aligned} \tag{15}$$

We can see that this identity is right, indeed, we shall give another natural method in order to see it.

In order to look for the norm admitting the restricted reproducing kernel of $G(s, t)$ to the interval $[a, b]$, note that the integral

$$\int_{-\infty}^a (f''(x)^2 + 2f'(x)^2 + f(x)^2) dx \tag{16}$$

is identical with its integral of the function

$$f(x) = 4f(a)G_a(x) - 4f'(a)G'_a(x) \tag{17}$$

that is the minimum norm square over $(-\infty, a)$ of the functions $W^{2,2}(\mathbf{R})$ taking the values $f(a)$ and $f'(a)$.

Here, following Theorem A, we wish to look for the function with the minimum norm; on the interval $[a, b]$ the functions are the same and so the minimum should be considered on the intervals $(-\infty, a]$ and $(b, \infty]$. On the interval $(-\infty, a]$ case, it is given by the function (17) in term of the reproducing kernel. The function (17) is given by

$$f(x) = [(A + B)(a - x) + A] \exp(x - a)$$

with

$$A = f(a), \quad B = -f'(a).$$

Then, by direct calculations, we have

$$\begin{aligned} \int_{-\infty}^a (f''(x))^2 + 2f'(x)^2 + f(x)^2 dx &= 2(A^2 + AB + B^2) \\ &= 2(f(a)^2 - f(a)f'(a) + f'(a)^2). \end{aligned} \tag{18}$$

From this result, we see that the corresponding inner product over $(-\infty, a)$ is represented by

$$\begin{aligned} (f_1, f_2)_{W^{2,2}(-\infty, a)} &= 2(f_1(a)f_2(a) + f'_1(a)f'_2(a)) \\ &\quad + \frac{1}{2}(f_1(a) - f'_1(a))(f_2(a) - f'_2(a)) \\ &\quad - \frac{1}{2}(f_1(a) + f'_1(a))(f_2(a) + f'_2(a)). \end{aligned} \tag{19}$$

The situation for the integrals over $(b, +\infty)$ is similar and so we obtain the desired isometric identity

$$\begin{aligned} \|f\|_{W^{2,2}(\mathbf{R})}^2 &= \|f\|_{W^{2,2}([a,b])}^2 + 2(f(a)^2 - f(a)f'(a) + f'(a)^2) \\ &\quad + 2(f(b)^2 - f(b)f'(b) + f'(b)^2). \end{aligned} \tag{20}$$

Therefore, the inner product relation is given by

$$\begin{aligned} (f_1, f_2)_{W^{2,2}(\mathbf{R})} &= (f_1, f_2)_{W^{2,2}([a,b])} + 2(f_1(a)f_2(a) + f'_1(a)f'_2(a)) \\ &\quad + \frac{1}{2}(f_1(a) - f'_1(a))(f_2(a) - f'_2(a)) \\ &\quad - \frac{1}{2}(f_1(a) + f'_1(a))(f_2(a) + f'_2(a)) \\ &\quad + 2(f_1(b)f_2(b) + f'_1(b)f'_2(b)) + \frac{1}{2}(f_1(b) - f'_1(b))(f_2(b) - f'_2(b)) \\ &\quad - \frac{1}{2}(f_1(b) + f'_1(b))(f_2(b) + f'_2(b)). \end{aligned}$$

We can confirm that this identity and (15) are consistent, directly. Indeed,

$$2(f(a)G_t(a) + f'(a)G'_t(a)) + \frac{1}{2}(f(a) - f'(a))(G_t(a) - G'_t(a)) - \frac{1}{2}(f(a) + f'(a))(G_t(a) + G'_t(a))$$

is identical with

$$+f(a)\frac{-t+a-2}{4}\exp(a-t) + f'(a)\frac{t-a-1}{4}\exp(a-t).$$

For the point b , the result is similar.

Of course, the isometrical identity and the inner product forms identity are valid with the minimum extension functions in Theorem A.

In particular, we have

THEOREM 1. *The extension of the functions f in $W_S([a, b])$ to $W^{2,2}(\mathbf{R})$ with the minimum norm is given by*

$$f(t) = (f, G(\cdot, t))_{W^{2,2}([a, b])} + 2(f(a)G_t(a) + f'(a)G'_t(a)) + \frac{1}{2}(f(a) - f'(a))(G_t(a) - G'_t(a)) - \frac{1}{2}(f(a) + f'(a))(G_t(a) + G'_t(a)) + 2(f(b)G_t(b) + f'(b)G'_t(b)) + \frac{1}{2}(f(b) - f'(b))(G_t(b) - G'_t(b)) - \frac{1}{2}(f(b) + f'(b))(G_t(b) + G'_t(b)).$$

COROLLARY 1. *We obtain the inequality for real valued functions f of $W^{2,2}(\mathbf{R})$*

$$\begin{aligned} \|f\|_{W^{2,2}(\mathbf{R})}^2 &= \|f''\|_{L^2(\mathbf{R})}^2 + 2\|f'\|_{L^2(\mathbf{R})}^2 + \|f\|_{L^2(\mathbf{R})}^2 \\ &\geq \|f''\|_{L^2([a, b])}^2 + 2\|f'\|_{L^2([a, b])}^2 + \|f\|_{L^2([a, b])}^2 \\ &\quad + 2(f(a)^2 - f(a)f'(a) + f'(a)^2) + 2(f(b)^2 - f(b)f'(b) + f'(b)^2). \end{aligned}$$

Equality holds for the minimum extension stated in Theorem 1.

Related versions

By the similar method or directly we have the following results.

Let

$$K(s, t) \equiv \int_0^\infty \frac{\cos(su)\cos(tu)}{u^2 + 1} du = \frac{\pi}{4} (\exp(-|s - t|) + \exp(-s - t)) \tag{21}$$

for $s, t > 0$. Then $H_K(0, \infty) = W^{1,2}(0, \infty)$ as a set of functions and the norm is given by:

$$\|f\|_{H_K(0, \infty)} = \sqrt{\frac{2}{\pi} \int_0^\infty (|f'(u)|^2 + |f(u)|^2) du} \tag{22}$$

[5], page 12–13). From the restriction of the kernel $K(s, t)$ to $[a, b]$, $a > 0$, we have the norm inequality

$$\begin{aligned} \|f\|_{H_K(0, \infty)}^2 &\geq \frac{2}{\pi} \frac{1 - \exp(-2a)}{1 + \exp(-2a)} |f(a)|^2 \\ &\quad + \frac{2}{\pi} \int_a^b (|f'(u)|^2 + |f(u)|^2) du + \frac{2}{\pi} |f(b)|^2. \end{aligned} \tag{23}$$

Indeed, the minimum norm function $f_a(x)$ having $f(a)$ over $[0, a]$ is given by

$$f_a(x) = \frac{f(a)}{1 + \exp(-2a)} (\exp(x - a) + \exp(-x - a))$$

and its norm square over $[0, a]$ is given

$$\frac{2}{\pi} \frac{1 - \exp(-2a)}{1 + \exp(-2a)} |f(a)|^2.$$

Meanwhile, the minimum norm function $f_b(x)$ having $f(b)$ over $[b, \infty)$ is given by

$$f_b(x) = \frac{f(b)}{1 + \exp(-2b)} (\exp(b - x) + \exp(-x - b))$$

and its norm square over $[b, \infty)$ is given by

$$\frac{2}{\pi} |f(b)|^2.$$

Let

$$K(s, t) \equiv \int_0^\infty \frac{\sin(su) \sin(tu)}{u^2 + 1} du = \frac{\pi}{4} (\exp(-|s - t|) - \exp(-s - t)) \tag{24}$$

for $s, t > 0$. Then we have

$$H_K(0, \infty) = \{f \in AC(0, \infty) : f(0) = 0\} \tag{25}$$

as a set of functions and the norm is given by

$$\|f\|_{H_K(0, \infty)} = \sqrt{\frac{2}{\pi} \int_0^\infty (|f'(u)|^2 + |f(u)|^2) du} \tag{26}$$

[5], 13–14). From the restriction of the kernel $K(s, t)$ to $[a, b]$, $a > 0$, we have the norm inequality, similarly

$$\begin{aligned} \|f\|_{H_K(0, \infty)}^2 &\geq \frac{2}{\pi} \frac{1 + \exp(-2a)}{1 - \exp(-2a)} |f(a)|^2 \\ &\quad + \frac{2}{\pi} \int_a^b (|f'(u)|^2 + |f(u)|^2) du + \frac{2}{\pi} |f(b)|^2. \end{aligned} \tag{27}$$

Let

$$K(s, t) \equiv \min(s, t) \quad (s, t > 0). \quad (28)$$

Then we have

$$H_K(0, \infty) = \left\{ f \in W^{1,2}(0, \infty) : \lim_{\varepsilon \downarrow 0} f(\varepsilon) = 0 \right\} \quad (29)$$

as a set of functions and the norm is given by

$$\|f\|_{H_K(0, \infty)} = \sqrt{\int_0^\infty |f'(u)|^2 du} \quad (30)$$

[5], 14–15). From the restriction of the kernel $K(s, t)$ to $[a, b]$, $a > 0$, we have the norm inequality

$$\|f\|_{H_K(0, \infty)}^2 \geq \frac{1}{a} |f(a)|^2 + \int_a^b |f'(u)|^2 du. \quad (31)$$

Indeed, the minimum norm function $f_a(x)$ having $f(a)$ over $[0, a]$ is given by

$$f_a(x) = \frac{x}{a} f(a)$$

and its norm square over $[0, a]$ is given by

$$\frac{1}{a} |f(a)|^2.$$

Meanwhile, the minimum norm function $f_b(x)$ having $f(b)$ over $[b, \infty)$ is given by

$$f_b(x) = b$$

and its norm square over $[b, \infty)$ is given by 0.

We have many type Sobolev Hilbert spaces. For example, for $\omega^2 = \gamma^2 - \alpha^2 > 0$, the kernel

$$K(s, t) = \frac{\exp(-\alpha|s-t|)}{4\alpha\gamma^2} \left\{ \cos(\omega|s-t|) + \frac{\alpha}{\omega} \sin(\omega|s-t|) \right\}$$

is the reproducing kernel for the Sobolev Hilbert space admitting the norm

$$\|u\|^2 = 4\alpha\gamma^2 u(a)^2 + 4\alpha u'(a)^2 + \int_a^b (u''(t) + 2\alpha u'(t) + \gamma^2 u(t))^2 dt$$

(E. Parzen, [2]).

See also [1] and the recent paper A. Yamada ([6]).

3. Basic applications of the realization of the restricted reproducing kernel Hilbert space

Theorem 1 and other derived identities show that the extension of the function with the minimum norm to the whole space (the half space) from a closed interval $[a, b]$ is given simply. This means that in the related Fourier transform, the inversion that corresponds to the function with the minimum norm may be calculated in terms of the values on the interval $[a, b]$.

REFERENCES

- [1] A. BERLINET AND C. THOMAS-AGNAN, *Reproducing Kernel Hilbert Spaces in Probability and Statistics*, Kluwer Academic Publishers, Boston/Dordrecht/London, 2004.
- [2] E. PARZEN, *An approach to time series analysis*, Ann. Math. Statist., **32** (1961), 951–989.
- [3] S. SAITOH, *Theory of reproducing kernels and its applications*, Pitman Research Notes in Mathematics Series, 189, Longman Scientific & Technical, Harlow; copublished in the United States with John Wiley & Sons, Inc., New York, 1988. x+157 pp. ISBN: 0-582-03564-3.
- [4] S. SAITOH, *Integral Transforms, Reproducing Kernels and their Applications*, Pitman Res. Notes in Math. Series **369** (1997), Addison Wesley Longman Ltd, UK, CRC Press, Taylor & Francis Group, Boca Raton London, New York (in hard cover).
- [5] S. SAITOH AND Y. SAWANO, *Theory of Reproducing Kernels and Applications*, Developments in Mathematics **44**, 2016 (Springer).
- [6] A. YAMADA, *The reproducing kernel of the Sobolev space on an interval*, Proc. Amer. Math. Soc., **151** (2023), 4435–4446, In *Bergman, Szegő and Sobolev kernels*, Surikaiseikikenkyu-sho Kokyuroku, **2256** (2023), 52–73.

(Received November 26, 2023)

Saburou Saitoh
Kawauchi-cho 5-1648-16, Kiryu 376-0041, Japan
e-mail: saburou.saitoh@gmail.com