

## LIMITED APPROXIMATION OF NUMERICAL RANGE OF NORMAL MATRIX

MARIA ADAM AND JOHN MAROULAS

(Communicated by C.-K. Li)

*Abstract.* Let  $A$  be an  $n \times n$  normal matrix, whose numerical range  $NR[A]$  is a  $k$ -polygon. If a unit vector  $v \in W \subseteq \mathbb{C}^n$ , with  $\dim W = k$  and the point  $v^*Av \in \text{Int}NR[A]$ , then  $NR[A]$  is circumscribed to  $NR[P^*AP]$ , where  $P$  is an  $n \times (k-1)$  isometry of  $\{\text{span}\{v\}\}_W^\perp \rightarrow \mathbb{C}^n$ , [1]. In this paper, we investigate an internal approximation of  $NR[A]$  by an increasing sequence of  $NR[C_s]$  of compressed matrices  $C_s = R_s^*AR_s$ , with  $R_s^*R_s = I_{k+s-1}$ ,  $s = 1, 2, \dots, n-k$  and additionally  $NR[A]$  is expressed as limit of numerical ranges of  $k$ -compressions of  $A$ .

### 1. Introduction and preliminaries

Let  $\mathcal{M}_n$  denote the algebra of all  $n \times n$  complex matrices. The *numerical range* of  $A \in \mathcal{M}_n$  is the well known set

$$NR[A] = \{x^*Ax \in \mathbb{C} : x \in \mathbb{C}^n \text{ with } \|x\|_2 = 1\},$$

which is a nonempty compact and convex subset of  $\mathbb{C}$  that contains the spectrum  $\sigma(A)$  of  $A$  (see [5, Chapter 1]). We recall that the numerical ranges of unitarily similar matrices are identified and if  $A = MDM^*$ ;  $D = \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_n)$  is the unitary diagonalising form of a normal matrix  $A$ , then  $NR[A] = \text{Co}\{\sigma(A)\}$ , where  $\text{Co}\{\cdot\}$  denotes the convex hull of the set.

Given two matrices  $A \in \mathcal{M}_n$  and  $C \in \mathcal{M}_k$  with  $1 \leq k < n$ , the matrix  $C$  is a *k-compression* of  $A$ , if there exists an  $n \times k$  orthonormal matrix  $P$  (i.e.,  $P^*P = I_k$ ) such that  $C = P^*AP$ . Clearly,

$$NR[C] = NR[P^*AP] \subseteq NR[A], \tag{1}$$

and the equality holds only for  $k = n$ .

Moreover, we have

$$NR[C] \subseteq NR[PCP^*],$$

since  $NR[C] = NR[P^*(PCP^*)P] \subseteq NR[PCP^*]$ .

The numerical range of compressions of normal matrices have attracted attention and several results have been published in [1, 2, 3, 4]. The inclusion relation of  $NR[A]$

---

*Mathematics subject classification* (2000): 15A18, 15A60, 47A20.

*Keywords and phrases:* Compression, eigenvalue, numerical range.

in (1) has been presented in details in [1], where the investigation leads to a structure of  $P$  such that the boundary of  $NR[P^*AP]$  is supported by the edges of the boundary of  $NR[A]$ .

To explain, consider for a normal matrix  $A \in \mathcal{M}_n$  the convex polygon  $\mathcal{P} = \langle \lambda_1, \lambda_2, \dots, \lambda_k \rangle = \text{Co}\{\sigma(D)\} = NR[A]$ , where the eigenvalues  $\lambda_i, i = 1, \dots, k$ , are simple. If  $W = \text{span}\{e_1, \dots, e_k\}$  and  $e_i, i = 1, \dots, k$ , are vectors of the standard basis of  $\mathbb{C}^n$ , then for every unit vector

$$v = \sum_{i=1}^k v_i e_i \in W \quad ; \quad v_i \in \mathbb{C} \setminus \{0\}, \quad i = 1, 2, \dots, k, \tag{2}$$

the point  $v^*Dv$  lies inside of the polygon  $\mathcal{P}$ . Denoting by  $E_W^\perp(v)$  the orthogonal complement of  $\text{span}\{v\}$  with respect to the subspace  $W$ , clearly for the vector  $\gamma = \gamma_1 e_1 + \dots + \gamma_k e_k \in E_W^\perp(v)$ , we have  $\gamma \circ v = \sum_{i=1}^k \bar{v}_i \gamma_i = 0$  and further we take:

$$\gamma = \gamma_1 \left( e_1 - \frac{\bar{v}_1}{v_j} e_j \right) + \gamma_2 \left( e_2 - \frac{\bar{v}_2}{v_j} e_j \right) + \dots + \gamma_k \left( e_k - \frac{\bar{v}_k}{v_j} e_j \right), \tag{3}$$

for an index  $j$ . Therefore, by the vectors

$$\begin{aligned} b_1 &= e_1 - \frac{\bar{v}_1}{v_j} e_j, \dots, \quad b_{j-1} = e_{j-1} - \frac{\bar{v}_{j-1}}{v_j} e_j, \\ b_j &= e_{j+1} - \frac{\bar{v}_{j+1}}{v_j} e_j, \dots, \quad b_{k-1} = e_k - \frac{\bar{v}_k}{v_j} e_j, \end{aligned}$$

an orthonormal basis  $\{w_1, w_2, \dots, w_{k-1}\}$  of  $E_W^\perp(v)$  is constructed. Defining the  $n \times (k-1)$  matrix

$$P = [w_1 \ w_2 \ \dots \ w_{k-1}], \tag{4}$$

and  $C = P^*DP$  the corresponding  $(k-1)$ -compression of  $D = \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_n)$ , we conclude

$$\begin{aligned} NR[C] &= \{(Pz)^*D(Pz) : z \in \mathbb{C}^{k-1}, \|z\|_2 = 1\} = \{x^*Dx : x = Pz \in E_W^\perp(v), \|x\|_2 = 1\} \\ &\subseteq \{x^*Dx : x \in W, \|x\|_2 = 1\} = \langle \lambda_1, \lambda_2, \dots, \lambda_k \rangle. \end{aligned}$$

Moreover, the following unit vectors of  $\mathbb{C}^n \cap E_W^\perp(v)$

$$y_i = \frac{\bar{v}_{i+1}}{\sqrt{|v_i|^2 + |v_{i+1}|^2}} e_i - \frac{\bar{v}_i}{\sqrt{|v_i|^2 + |v_{i+1}|^2}} e_{i+1} \quad ; \quad i = 1, 2, \dots, k, \tag{5}$$

where in (5)  $e_{k+1}$  is substituted by  $e_1$  and  $v_{k+1}$  by  $v_1$ , correspond to the points

$$c_i = y_i^*Dy_i = \frac{|v_{i+1}|^2 \lambda_i + |v_i|^2 \lambda_{i+1}}{|v_i|^2 + |v_{i+1}|^2} \quad ; \quad i = 1, 2, \dots, k \quad ; \quad \lambda_{k+1} = \lambda_1 \tag{6}$$

which belong to the line segment  $\langle \lambda_i, \lambda_{i+1} \rangle \subset \partial NR[D]$ . Obviously, the points  $c_i$  depend on the unit vector  $v$  and by Theorem 1 in [1] we have  $\partial NR[D] \cap \partial NR[C] = \{c_1, \dots, c_k\}$ .

In the next section, we construct a sequence  $C_s$ ,  $s = 1, 2, \dots, n - k$  of compressions of a normal matrix  $A$  such that the area of  $NR[C_s]$  is increasing and is close enough to the polygon  $\mathcal{P}$ . Also, for  $i = 1, 2, \dots, k$ , a sequence of points  $t_{i,m} \in NR[C_{1,m}] \cap \langle \lambda_i, \lambda_r \rangle$  is constructed, where the matrix  $C_{1,m}$  is a  $k$ -compression of  $A$ , depending on a vector  $\zeta_m$ , with  $\|\zeta_m\|_2 \rightarrow \infty$  and  $\lambda_r$  is an interior eigenvalue of the polygon, finding out that  $\lim_{m \rightarrow \infty} t_{i,m} = \lambda_i$ . By this statement we are led to  $\lim_{m \rightarrow \infty} NR[C_{1,m}] = NR[A]$ . Analogue results are obtained for subpolygons of  $\mathcal{P}$ .

### 2. An interior approximation of $NR[A]$

The interior approximation of the boundary of  $NR[A]$  can be further elaborated, using a *compression* of a normal matrix  $A$  by a sequence of numerical ranges of suitable matrices.

**PROPOSITION 1.** *Let  $A$  be an  $n \times n$  normal matrix, where its numerical range is a  $k$ -polygon  $\mathcal{P}$ . Then there exists a finite sequence of compressions  $C_s = R_s^*DR_s$ , with  $R_s^*R_s = I_{k+s-1}$ ,  $s = 1, 2, \dots, n - k$ , such that*

$$NR[C] \subseteq NR[C_1] \subseteq NR[C_2] \subseteq \dots \subseteq NR[C_{n-k}] \subseteq NR[A], \tag{7}$$

and for every  $s$ , we have  $\{c_1, \dots, c_k\} \subseteq NR[C_s] \cap \mathcal{P}$ , with  $c_i$  in (6).

*Proof.* Consider the unit vector  $v \in W$  in (2) and let

$$\xi_1 = v + \pi_{k+1}e_{k+1} = \sum_{i=1}^k v_i e_i + \pi_{k+1}e_{k+1}.$$

If  $W_1 = \text{span}\{W, e_{k+1}\}$  and  $\gamma = (\gamma_1, \dots, \gamma_k, \gamma_{k+1}) \in E_{W_1}^\perp(\xi_1)$ , then,  $\gamma \circ \xi_1 = \sum_{i=1}^k \overline{v}_i \gamma_i + \overline{\pi}_{k+1} \gamma_{k+1} = 0$  and for the same index  $j$  as in (3), we have:

$$\gamma = \gamma_1 \left( e_1 - \frac{\overline{v}_1}{v_j} e_j \right) + \dots + \gamma_k \left( e_k - \frac{\overline{v}_k}{v_j} e_j \right) + \gamma_{k+1} \left( e_{k+1} - \frac{\overline{\pi}_{k+1}}{v_j} e_j \right).$$

Thus, the orthonormal basis  $\{w_1, \dots, w_{k-1}, r_1\}$  is constructed by the vectors

$$b_1 = e_1 - \frac{\overline{v}_1}{v_j} e_j, \dots, \quad b_{j-1} = e_{j-1} - \frac{\overline{v}_{j-1}}{v_j} e_j,$$

$$b_j = e_{j+1} - \frac{\overline{v}_{j+1}}{v_j} e_j, \dots, \quad b_{k-1} = e_k - \frac{\overline{v}_k}{v_j} e_j, \quad b_k = e_{k+1} - \frac{\overline{\pi}_{k+1}}{v_j} e_j.$$

Denoting by  $R_1 = [P \ r_1]$ , where  $P$  is the matrix in (4), clearly  $R_1^*R_1 = I_k$  and the equation

$$C_1 = R_1^*DR_1 = \begin{bmatrix} P^*DP & P^*Dr_1 \\ r_1^*DP & r_1^*Dr_1 \end{bmatrix} \tag{8}$$

yields the inclusion

$$NR[C] = NR[P^*DP] \subseteq NR[C_1].$$

If  $W_2 = \text{span}\{W_1, e_{k+2}\} = \text{span}\{W, e_{k+1}, e_{k+2}\}$  and  $\xi_2 = \xi_1 + \pi_{k+2}e_{k+2}$ , similarly we define the orthonormal basis  $\{w_1, \dots, w_{k-1}, r_1, r_2\}$  of  $E_{W_2}^\perp(\xi_2)$  and the matrix  $R_2 = \begin{bmatrix} P & r_1 & r_2 \end{bmatrix} = \begin{bmatrix} R_1 & r_2 \end{bmatrix}$ . Thus,

$$C_2 = R_2^*DR_2 = \begin{bmatrix} R_1^*DR_1 & R_1^*Dr_2 \\ r_2^*DR_1 & r_2^*Dr_2 \end{bmatrix},$$

concluding that  $NR[C_1] = NR[R_1^*DR_1] \subseteq NR[C_2]$ . Continuing in the same way, we consider the vector

$$\xi_{n-k} = v + \pi_{k+1}e_{k+1} + \pi_{k+2}e_{k+2} + \dots + \pi_n e_n$$

of subspace  $W_{n-k} = \text{span}\{W, e_{k+1}, \dots, e_n\}$  and for the same index  $j$  as in (3), we receive the orthonormal basis  $\{w_1, \dots, w_{k-1}, r_1, r_2, \dots, r_{n-k}\}$  of  $E_{W_{n-k}}^\perp(\xi_{n-k})$ . If

$$C_{n-k} = R_{n-k}^*DR_{n-k},$$

where  $R_{n-k} = \begin{bmatrix} R_{n-k-1} & r_{n-k} \end{bmatrix} = \dots = \begin{bmatrix} P & r_1 & r_2 & \dots & r_{n-k} \end{bmatrix}_{n \times (n-1)}$ , clearly

$$NR[C_{n-k-1}] \subseteq NR[C_{n-k}] \subseteq NR[A].$$

Furthermore, by the inclusions in (7), we have that the tangential points  $c_i$  in (6) of  $NR[C]$  and the polygon  $\mathcal{P} = NR[A]$ , belong also to  $NR[C_s]$ , for  $s = 1, \dots, n - k$ . Note that the vectors  $y_i$  in (5) belong to the subspaces  $E_{W_s}^\perp(\xi_s)$ , with  $\xi_s = v + \pi_{k+1}e_{k+1} + \pi_{k+2}e_{k+2} + \dots + \pi_{k+s}e_{k+s}$ ,  $s = 1, 2, \dots, n - k$  and for the unit vectors  $g_{i,s}$ ,  $i = 1, \dots, k$ , defined by the equation  $R_s g_{i,s} = y_i$ , we have:

$$c_i = \frac{|v_i|^2 \lambda_{i+1} + |v_{i+1}|^2 \lambda_i}{|v_i|^2 + |v_{i+1}|^2} = y_i^* D y_i = g_{i,s}^* (R_s^* D R_s) g_{i,s} = g_{i,s}^* C_s g_{i,s},$$

with  $\|g_{i,s}\|_2 = 1$ .  $\square$

If, instead of  $v$  in (2), we consider the vector

$$u = \sum_{j=k+1}^n u_j e_j,$$

where  $e_{k+1}, \dots, e_n$  are the remaining vectors of the standard basis of  $\mathbb{C}^n$  and simultaneously the eigenvectors of  $D$  corresponding to the eigenvalues in the interior of polygon  $\mathcal{P}$ , then  $E_W^\perp(u) = W$ . Thus for  $\tilde{P} = \begin{bmatrix} e_1 & e_2 & \dots & e_k \end{bmatrix}_{n \times k} = \begin{bmatrix} I_k \\ \mathbb{O}_{n-k} \end{bmatrix}$ , we obtain:

$$\begin{aligned} NR[\tilde{P}^*D\tilde{P}] &= \{(\tilde{P}z)^* D (\tilde{P}z) : z \in \mathbb{C}^k, \|z\|_2 = 1\} \\ &= \{z^* \text{diag}(\lambda_1, \dots, \lambda_k) z : z \in \mathbb{C}^k, \|z\|_2 = 1\} = \mathcal{P}. \end{aligned}$$

Regarding a vector,

$$\beta_{i,\tau} = u + \sum_{j=i}^{\tau} \rho_j e_j; \text{ for } 1 \leq i \leq \tau \leq k, \tag{9}$$

along similar lines as in Proposition 1, we conclude the following proposition.

PROPOSITION 2. *Let  $A$  be an  $n \times n$  normal matrix, whose the numerical range is a  $k$ -polygon  $\mathcal{P}$ . Let also a vector  $\beta_{i,\tau}$  as in (9). Then there exists a  $(n - 1)$ -compression  $\tilde{C}_{i,\tau}$  of  $D$  such that*

$$NR[\tilde{C}_{i,\tau}] = Co\{\langle \lambda_1, \dots, \lambda_{i-1} \rangle \cup \langle \lambda_{\tau+1}, \dots, \lambda_k \rangle \cup NR[B_{i,\tau}]\}, \tag{10}$$

where  $B_{i,\tau}$  is an  $(n - k + \tau - i)$ -compression of  $D$ .

*Proof.* Let a vector  $\gamma = (\gamma_1, \gamma_2, \dots, \gamma_n) \in \text{span}\{\beta_{i,\tau}\}^\perp$ , then

$$\gamma \circ \beta_{i,\tau} = \sum_{j=k+1}^n \bar{u}_j \gamma_j + \sum_{j=i}^{\tau} \bar{\rho}_j \gamma_j = 0.$$

Thus, for an index  $\ell$  with  $k + 1 \leq \ell \leq n$ , we have:

$$\begin{aligned} \gamma = & \gamma_1 e_1 + \dots + \gamma_i \left( e_i - \frac{\bar{\rho}_i}{\bar{u}_\ell} e_\ell \right) + \dots + \gamma_\tau \left( e_\tau - \frac{\bar{\rho}_\tau}{\bar{u}_\ell} e_\ell \right) + \gamma_{\tau+1} e_{\tau+1} + \dots + \gamma_k e_k \\ & + \gamma_{k+1} \left( e_{k+1} - \frac{\bar{u}_{k+1}}{\bar{u}_\ell} e_\ell \right) + \dots + \gamma_n \left( e_n - \frac{\bar{u}_n}{\bar{u}_\ell} e_\ell \right) \end{aligned}$$

By the vectors  $\omega_j = e_j - \frac{\bar{\rho}_j}{\bar{u}_\ell} e_\ell$  ( $j = i, \dots, \tau$ ) and  $\phi_{k+1} = e_{k+1} - \frac{\bar{u}_{k+1}}{\bar{u}_\ell} e_\ell, \dots, \phi_{\ell-1} = e_{\ell-1} - \frac{\bar{u}_{\ell-1}}{\bar{u}_\ell} e_\ell, \phi_\ell = e_{\ell+1} - \frac{\bar{u}_{\ell+1}}{\bar{u}_\ell} e_\ell, \dots, \phi_{n-1} = e_n - \frac{\bar{u}_n}{\bar{u}_\ell} e_\ell$  an orthonormal basis

$$\{e_1, \dots, e_{i-1}, \hat{\omega}_i, \dots, \hat{\omega}_\tau, e_{\tau+1}, \dots, e_k, \hat{\phi}_{k+1}, \dots, \hat{\phi}_{n-1}\}$$

is constructed and the  $n \times (n - 1)$  matrix

$$\tilde{P}_{i,\tau} = [Q_1 \quad Q_2 \quad \Omega \quad \Phi], \tag{11}$$

where  $Q_1 = [e_1 \ e_2 \ \dots \ e_{i-1}]$ ,  $Q_2 = [e_{\tau+1} \ e_{\tau+2} \ \dots \ e_k]_{n \times (k-\tau)}$ ,  $\Omega = [\hat{\omega}_i \ \dots \ \hat{\omega}_\tau]_{n \times (\tau-i+1)}$ ,  $\Phi = [\hat{\phi}_{k+1} \ \dots \ \hat{\phi}_{n-1}]_{n \times (n-k-1)}$ , is an isometry. Hence by the  $(n - 1)$ -compression of  $D$

$$\begin{aligned} \tilde{C}_{i,\tau} = \tilde{P}_{i,\tau}^* D \tilde{P}_{i,\tau} &= \begin{bmatrix} Q_1^* D Q_1 & \mathbb{O} & \mathbb{O} & \mathbb{O} \\ \mathbb{O} & Q_2^* D Q_2 & \mathbb{O} & \mathbb{O} \\ \mathbb{O} & \mathbb{O} & \Omega^* D \Omega & \Omega^* D \Phi \\ \mathbb{O} & \mathbb{O} & \Phi^* D \Omega & \Phi^* D \Phi \end{bmatrix} \\ &= \begin{bmatrix} \text{diag}(\lambda_1, \dots, \lambda_{i-1}) & \mathbb{O} & \mathbb{O} & \mathbb{O} \\ \mathbb{O} & \text{diag}(\lambda_{\tau+1}, \dots, \lambda_k) & \mathbb{O} & \mathbb{O} \\ \mathbb{O} & \mathbb{O} & \mathbb{O} & \Omega^* D \Omega \ \Omega^* D \Phi \\ \mathbb{O} & \mathbb{O} & \mathbb{O} & \Phi^* D \Omega \ \Phi^* D \Phi \end{bmatrix} \end{aligned} \tag{12}$$

we are led to the relation

$$NR[\tilde{C}_{i,\tau}] = \text{Co}\{\langle \lambda_1, \dots, \lambda_{i-1} \rangle \cup \langle \lambda_{\tau+1}, \dots, \lambda_k \rangle \cup NR[B_{i,\tau}]\},$$

where  $B_{i,\tau} = \begin{bmatrix} \Omega^* D \Omega & \Omega^* D \Phi \\ \Phi^* D \Omega & \Phi^* D \Phi \end{bmatrix}$  is an  $(n - k + \tau - i)$ -compression of  $D$ .  $\square$

Considering the vector

$$\beta_1 = u + \rho_i e_i; \quad i \in \{1, 2, \dots, k\},$$

as in (9), by (12) we may construct the corresponding compression

$$\tilde{C}_1 = \text{diag}(\text{diag}(\lambda_1, \dots, \lambda_{i-1}), \text{diag}(\lambda_{i+1}, \dots, \lambda_k), B_1), \tag{13}$$

where  $B_1 = \begin{bmatrix} \hat{\omega}_i^* D \hat{\omega}_i & \hat{\omega}_i^* D \Phi \\ \Phi^* D \hat{\omega}_i & \Phi^* D \Phi \end{bmatrix}$  is  $(n - k)$ -compression of  $D$ . Due to the construction of the orthonormal basis  $\{\hat{\omega}_i, \Phi\}$ ,  $\partial NR[B_1]$  is inscribed to the polygon  $\langle \lambda_i, \lambda_{k+1}, \dots, \lambda_n \rangle$ .

**PROPOSITION 3.** *Let  $A$  be an  $n \times n$  normal matrix and the polygon  $\mathcal{P} = \langle \lambda_1, \dots, \lambda_k \rangle = NR[A]$ .*

**I.** *If we consider a sequence of vectors  $\zeta_m = v + q_m e_j$ , where  $e_j$  is eigenvector of  $D$  corresponding to the interior eigenvalue  $\lambda_j$  of  $\mathcal{P}$ , such that  $\lim_{m \rightarrow \infty} \|\zeta_m\|_2 = \infty$ , and the matrices  $C_{1,m} \in \mathcal{M}_k$  are the  $k$ -compressions of  $D$  in (8) defined by  $\zeta_m$ , then there exists a sequence of points  $t_{i,m} \in NR[C_{1,m}] \cap \langle \lambda_i, \lambda_j \rangle$ , such that  $\lim_{m \rightarrow \infty} t_{i,m} = \lambda_i$ , for  $i \in \{1, 2, \dots, k\}$ .*

**II.** *Let  $\beta_m = \sum_{j=k+1}^n u_{j,m} e_j + \rho_i e_i$  be a sequence of vectors, with  $i \in \{1, 2, \dots, k\}$ .*

*If  $\lim_{m \rightarrow \infty} |u_{j,m}| = \infty$  holds for a prefixed  $j$ , and  $\tilde{C}_{1,m}$  is the corresponding  $(n - 1)$ -compression of  $D$  in (13), then there exists a sequence of points  $c_{i,m} \in NR[\tilde{C}_{1,m}] \cap \langle \lambda_i, \lambda_j \rangle$ , such that  $\lim_{m \rightarrow \infty} c_{i,m} = \lambda_i$ .*

*Proof.* **I.** Consider the unit vectors of  $\mathbb{C}^n$

$$z_{i,m} = \frac{\bar{q}_m}{\sqrt{|v_i|^2 + |q_m|^2}} e_i - \frac{\bar{v}_i}{\sqrt{|v_i|^2 + |q_m|^2}} e_j, \quad i = 1, 2, \dots, k,$$

and the vectors  $f_{i,m} \in \mathbb{C}^k$  defined by the equations  $\tilde{R}_{1,m} f_{i,m} = z_{i,m}$ , where the  $n \times k$  matrix  $\tilde{R}_{1,m}$  is constructed by  $\zeta_m$ , as  $R_1$  in Proposition 1. Obviously,  $\|f_{i,m}\|_2 = 1$  and the points

$$\begin{aligned} t_{i,m} &= f_{i,m}^* C_{1,m} f_{i,m} = f_{i,m}^* \tilde{R}_{1,m}^* D \tilde{R}_{1,m} f_{i,m} = z_{i,m}^* D z_{i,m} = \frac{|v_i|^2 \lambda_j + |q_m|^2 \lambda_i}{|v_i|^2 + |q_m|^2} \\ &= \lambda_i + \frac{|v_i|^2}{|v_i|^2 + |q_m|^2} (\lambda_j - \lambda_i), \quad i = 1, 2, \dots, k, \end{aligned} \tag{14}$$

belong to  $NR[C_{1,m}] \cap \langle \lambda_i, \lambda_j \rangle$ . Moreover, due to  $\lim_{m \rightarrow \infty} \|\zeta_m\|_2 = \infty$  by (14) we have

$$\lim_{m \rightarrow \infty} t_{i,m} = \lim_{m \rightarrow \infty} \left( \lambda_i + \frac{|v_i|^2}{|v_i|^2 + |q_m|^2} (\lambda_j - \lambda_i) \right) = \lambda_i.$$

**II.** Consider the unit vectors of  $\mathbb{C}^n$

$$\psi_{j,m} = \frac{\bar{\rho}_i}{\sqrt{|u_{j,m}|^2 + |\rho_i|^2}} e_j - \frac{\bar{u}_{j,m}}{\sqrt{|u_{j,m}|^2 + |\rho_i|^2}} e_i,$$

and the vectors  $h_{j,m} \in \mathbb{C}^{n-1}$  defined by the equations  $\tilde{P}_{1,m} h_{j,m} = \psi_{j,m}$ , where  $\tilde{P}_{1,m}$  is constructed as in (11). Then, the point

$$c_{i,m} = h_{j,m}^* \tilde{C}_{1,m} h_{j,m} = h_{j,m}^* \tilde{P}_{1,m}^* D \tilde{P}_{1,m} h_{j,m} = \psi_{j,m}^* D \psi_{j,m} = \frac{|u_{j,m}|^2 \lambda_i + |\rho_i|^2 \lambda_j}{|u_{j,m}|^2 + |\rho_i|^2} \quad (15)$$

belongs to  $NR[\tilde{C}_{1,m}] \cap \langle \lambda_i, \lambda_j \rangle$ , and due to  $\lim_{m \rightarrow \infty} |u_{j,m}| = \infty$ , the equality in (15) yields

$$\lim_{m \rightarrow \infty} c_{i,m} = \lim_{m \rightarrow \infty} \left( \lambda_i + \frac{|\rho_i|^2}{|u_{j,m}|^2 + |\rho_i|^2} (\lambda_j - \lambda_i) \right) = \lambda_i. \quad \square$$

Clearly, by Proposition 3 **I**,  $\lim_{m \rightarrow \infty} |t_{i,m}| = |\lambda_i|$ . If the point  $\ell_{i,m}$  lies on  $\partial NR[C_{1,m}] \cap \langle \lambda_i, \lambda_j \rangle$ , we have

$$|t_{i,m}| \leq |\ell_{i,m}| \leq |\lambda_i|,$$

i.e.,

$$\lim_{m \rightarrow \infty} |\ell_{i,m}| = |\lambda_i|.$$

Therefore, there exist an index  $m_0(i) \in \mathbb{N}$  and small enough  $\varepsilon > 0$ , such that for  $m \geq m_0(i)$ , the distance  $d(t_{i,m}, \partial NR[C_{1,m}]) < \varepsilon$ .

Numerically, we may assume that the equality  $|t_{i,m_0(i)}| \approx |\ell_{i,m_0(i)}|$  holds and the equality in (14) leads to

$$|t_{i,m}| = \frac{|v_i|^2}{|v_i|^2 + |q_m|^2} |\lambda_j| + \frac{|q_m|^2}{|v_i|^2 + |q_m|^2} |\lambda_i|,$$

whereby we derive

$$|q_{m_0(i)}|^2 \approx |v_i|^2 \frac{|\ell_{i,m_0(i)}| - |\lambda_j|}{|\lambda_i| - |\ell_{i,m_0(i)}|}.$$

Moreover, for  $m_1 < m_2$ , due to  $\lim_{m \rightarrow \infty} \|\zeta_m\|_2 = \infty$  by (14) we have

$$|t_{i,m_1} - \lambda_j| = \frac{|q_{m_1}|^2}{|v_i|^2 + |q_{m_1}|^2} |\lambda_i - \lambda_j| \leq \frac{|q_{m_2}|^2}{|v_i|^2 + |q_{m_2}|^2} |\lambda_i - \lambda_j| = |t_{i,m_2} - \lambda_j|,$$

yielding

$$NR[C_{1,m_1}] \subseteq NR[C_{1,m_2}]. \tag{16}$$

Since the sequence  $|q_m|$  is increasing, by (16) for  $m \geq m_0(i)$ , we conclude

$$|q_m|^2 = |v_i|^2 \frac{|t_{i,m}| - |\lambda_j|}{|\lambda_i| - |t_{i,m}|} \geq |q_{m_0(i)}|^2 = |v_i|^2 \frac{|\ell_{i,m_0(i)}| - |\lambda_j|}{|\lambda_i| - |\ell_{i,m_0(i)}|},$$

i.e.,  $t_{i,m}$  has to be nearly a boundary point of  $NR[C_{1,m}]$ . Hence, for  $m = m_0(i)$  we can write

$$t_{i,m_0(i)} \approx f_{i,m_0(i)}^* C_{1,m_0(i)} f_{i,m_0(i)},$$

where  $f_{i,m_0(i)}$  is the eigenvector of  $H(e^{-i\theta_i} C_{1,m_0(i)})$  corresponding to the largest eigenvalue,  $\lambda_{\max}(H(e^{-i\theta_i} C_{1,m_0(i)}))$ , of hermitian part of matrix  $e^{-i\theta_i} C_{1,m_0(i)}$ , and  $\theta_i \in [0, 2\pi)$  is the argument of  $t_{i,m_0(i)}$ , (see [5, p. 35, Theorem 1.5.11]).

**THEOREM 4.** *For any normal matrix  $A$ , whose  $NR[A]$  is a  $k$ -polygon, there exists a sequence of  $k$ -compressions  $C_{1,m}$  of  $D$  in (8) such that  $NR[C_{1,m}]$  is inscribed to the polygon for every  $m$  and  $\lim_{m \rightarrow \infty} NR[C_{1,m}] = NR[A]$ .*

*Proof.* Let  $\mathcal{Q}_m = \text{Co}\{t_{1,m(1)}, \dots, t_{k,m(k)}\}$ . If  $m_0 = \max\{m_0(1), m_0(2), \dots, m_0(k)\}$ , then by Proposition 3 I, for  $m > m_0$  and small enough  $\varepsilon > 0$ , we estimate that

$$|\mathcal{Q}_m| \leq |NR[C_{1,m}]| \leq |\mathcal{P}|,$$

where  $|\cdot|$  denotes the area of a convex set. Since  $\lim_{m \rightarrow \infty} \mathcal{Q}_m = \mathcal{P}$ , obviously we have the convergence of area of  $NR[C_{1,m}]$  to the area contained in  $NR[A]$ .  $\square$

**COROLLARY 5.** *For any normal matrix  $A$ , whose  $NR[A]$  is a  $k$ -polygon, there exists a sequence of vectors  $\beta_m = \sum_{j=k+1}^n u_{j,m} e_j + \rho_i e_i; i \in \{1, 2, \dots, k\}$ , and the associated sequence  $\tilde{C}_{1,m}$  of  $(n - 1)$ -compressions of  $D$  in (13), such that*

$$\lim_{m \rightarrow \infty} NR[\tilde{C}_{1,m}] = NR[A],$$

when  $\lim_{m \rightarrow \infty} |u_{j,m}| = \infty$ .

*Proof.* Since, by (13) the compression matrix

$$\tilde{C}_{1,m} = \text{diag}(\text{diag}(\lambda_1, \dots, \lambda_{i-1}), \text{diag}(\lambda_{i+1}, \dots, \lambda_k), B_{1,m}),$$

clearly

$$NR[B_{1,m}] \subseteq \text{Co}\{\lambda_i, \lambda_{k+1}, \dots, \lambda_n\},$$



and due to  $\lim_{m \rightarrow \infty} c_{i,m} = \lambda_i$ , we obtain

$$\lim_{m \rightarrow \infty} NR[B_{1,m}] = \langle \lambda_i, \lambda_{k+1}, \dots, \lambda_n \rangle.$$

Hence, by (10) we have

$$\begin{aligned} \lim_{m \rightarrow \infty} NR[\tilde{C}_{1,m}] &= \text{Co}\{\langle \lambda_1, \dots, \lambda_{i-1} \rangle \cup \langle \lambda_{i+1}, \dots, \lambda_k \rangle \cup \lim_{m \rightarrow \infty} NR[B_{1,m}]\} \\ &= \text{Co}\{\langle \lambda_1, \dots, \lambda_{i-1} \rangle \cup \langle \lambda_{i+1}, \dots, \lambda_k \rangle \cup \langle \lambda_i, \lambda_{k+1}, \dots, \lambda_n \rangle\} = \langle \lambda_1, \dots, \lambda_k \rangle. \end{aligned}$$

□

The next example illustrates Proposition 3 I and indirectly Theorem 4.

EXAMPLE. Let the  $6 \times 6$  normal matrix  $A = \text{diag}(4\mathbf{i}, -2, -3\mathbf{i}, 5, 0, 1 + \mathbf{i})$ , where  $NR[A] = \text{Co}\{4\mathbf{i}, -2, -3\mathbf{i}, 5\}$ , i.e., 0 and  $1 + \mathbf{i}$  belong to  $\text{Int}NR[A]$ . For the unit vector  $v = \frac{1}{\sqrt{15}}e_1 + \frac{2}{\sqrt{15}}e_2 + \frac{1}{\sqrt{15}}e_3 + \frac{3}{\sqrt{15}}e_4$ , we have the matrix in (4),

$$P = \begin{bmatrix} -0.8944 & -0.1826 & -0.3162 \\ 0.4472 & -0.3651 & -0.6325 \\ 0 & 0.9129 & -0.3162 \\ 0 & 0 & 0.6325 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

and the tangent points in (6) of  $\partial NR[A] \cap \partial NR[P^*AP]$

$$c_1 = \frac{-2 + 16\mathbf{i}}{5}, \quad c_2 = \frac{-2 - 12\mathbf{i}}{5}, \quad c_3 = \frac{5 - 27\mathbf{i}}{10}, \quad c_4 = \frac{5 + 36\mathbf{i}}{10}.$$

If  $\zeta_1 = v + \frac{4}{\sqrt{15}}e_5$ , we obtain  $\tilde{R}_{1,1} = \begin{bmatrix} -0.8944 & -0.1826 & -0.3162 & -0.1855 \\ 0.4472 & -0.3651 & -0.6325 & -0.3710 \\ 0 & 0.9129 & -0.3162 & -0.1855 \\ 0 & 0 & 0.6325 & -0.5565 \\ 0 & 0 & 0 & 0.6956 \\ 0 & 0 & 0 & 0 \end{bmatrix}$  and

the matrix  $C_{1,1} = \tilde{R}_{1,1}^* A \tilde{R}_{1,1}$  as in (8). By (14), for  $\lambda_5 = 0$ , we take:  $t_{1,1} = \frac{64\mathbf{i}}{17} = 3.7647\mathbf{i} \in \langle \lambda_1, \lambda_5 \rangle$ . Also,  $\theta_1 = \pi/2$  and  $|t_{1,1}| = 3.7647 \neq 3.8691 = \lambda_{\max}(H(e^{-i\theta_1} C_{1,1}))$ , i.e.,  $t_{1,1}$  is interior point of  $NR[C_{1,1}]$ .

Similarly, if  $\zeta_2 = v + \frac{20}{\sqrt{15}}e_5$ , we have  $\tilde{R}_{1,2} = \begin{bmatrix} -0.8944 & -0.1826 & -0.3162 & -0.2535 \\ 0.4472 & -0.3651 & -0.6325 & -0.5070 \\ 0 & 0.9129 & -0.3162 & -0.2535 \\ 0 & 0 & 0.6325 & -0.7605 \\ 0 & 0 & 0 & 0.1901 \\ 0 & 0 & 0 & 0 \end{bmatrix}$

and the matrix  $C_{1,2} = \tilde{R}_{1,2}^* A \tilde{R}_{1,2}$  as in (8). By (14) we take:  $t_{1,2} = \frac{1600\mathbf{i}}{401} = 3.9900\mathbf{i} \in \langle \lambda_1, \lambda_5 \rangle$  and  $|t_{1,2}| = 3.9900 \neq 3.9904 = \lambda_{\max}(H(e^{-i\theta_1} C_{1,2}))$ , i.e.,  $t_{1,2}$  is interior point of  $NR[C_{1,2}]$ .


If  $\zeta_3 = v + \frac{100}{\sqrt{15}}e_5$ , we have  $\tilde{R}_{1,3} = \begin{bmatrix} -0.8944 & -0.1826 & -0.3162 & -0.2580 \\ 0.4472 & -0.3651 & -0.6325 & -0.5160 \\ 0 & 0.9129 & -0.3162 & -0.2580 \\ 0 & 0 & 0.6325 & -0.7740 \\ 0 & 0 & 0 & 0.0387 \\ 0 & 0 & 0 & 0 \end{bmatrix}$  and

$C_{1,3} = \tilde{R}_{1,3}^* A \tilde{R}_{1,3}$ . By (14) we take:  $t_{1,3} = \frac{40000i}{10001} = 3.99960004i \in \langle \lambda_1, \lambda_5 \rangle$  and  $|t_{1,3}| = 3.99960004 \approx \lambda_{\max}(H(e^{-i\theta_1} C_{1,3})) = 3.9996006$ . The point  $t_{1,3}$  almost lies on the  $\partial NR[C_{1,3}]$ , i.e.,  $t_{1,3} \approx f_{1,3}^* C_{1,3} f_{1,3} = 0.00000036896091 + 3.99960058200810i \in \partial NR[C_{1,3}]$ , where  $f_{1,3} = [0.8945 \ 0.1825 \ 0.3163 \ 0.2580]^T$  is the eigenvector of  $H(e^{-i\theta_1} C_{1,3})$  corresponding to  $\lambda_{\max}(H(e^{-i\theta_1} C_{1,3}))$ .

If  $\zeta_4 = v + \frac{120}{\sqrt{15}}e_5$ , we have  $\tilde{R}_{1,4} = \begin{bmatrix} -0.8944 & -0.1826 & -0.3162 & -0.2581 \\ 0.4472 & -0.3651 & -0.6325 & -0.5161 \\ 0 & 0.9129 & -0.3162 & -0.2581 \\ 0 & 0 & 0.6325 & -0.7742 \\ 0 & 0 & 0 & 0.0323 \\ 0 & 0 & 0 & 0 \end{bmatrix}$  and

$C_{1,4} = \tilde{R}_{1,4}^* A \tilde{R}_{1,4}$ . By (14) we take:  $t_{1,4} = \frac{57600i}{14401} = 3.9997i \in \langle \lambda_1, \lambda_5 \rangle$  and  $|t_{1,4}| = 3.9997222 \approx \lambda_{\max}(H(e^{-i\theta_1} C_{1,4})) = 3.9997225$ . Since  $q_3 = \frac{100}{\sqrt{15}} < \frac{120}{\sqrt{15}} = q_4$ , then  $NR[C_{1,3}] \subseteq NR[C_{1,4}]$  and we expect  $t_{1,4}$  to approximate  $\partial NR[C_{1,4}]$ . In fact,  $f_{1,4} = [0.8945 \ 0.1826 \ 0.3162 \ 0.2581]^T$  is eigenvector of  $H(e^{-i\theta_1} C_{1,4})$  and  $t_{1,4} \approx f_{1,4}^* C_{1,4} f_{1,4} = 0.00000017808543 + 3.99972250302240i$ .

In the next figure the numerical ranges of the compressions  $P^*AP$ ,  $C_{1,1}$  and  $C_{1,2}$  are illustrated.



Let  $Q = \langle \lambda_1, \lambda_2, \dots, \lambda_v \rangle$  be subpolygon of  $\mathcal{P}$ , with  $3 \leq v < k$  and the sequence of vectors of  $\mathbb{C}^n$

$$\eta_\mu = \sum_{i=1}^v v_i e_i + \varphi_\mu e_j; \quad j \in \{k+1, \dots, n\}$$

and suppose furthermore that the eigenvalue  $\lambda_j$  may not belong to  $Q$ . Denoting by  $G_{1,\mu} = T_{1,\mu}^* D T_{1,\mu}$  the  $v$ -compression of  $D$ , then  $NR[G_{1,\mu}]$  is tangent to the polygon  $\langle \lambda_1, \lambda_2, \dots, \lambda_v, \lambda_j \rangle$ . Thus, when  $\lim_{\mu \rightarrow \infty} \|\eta_\mu\|_2 = \infty$ , by Theorem 4 we conclude the equality

$$\lim_{\mu \rightarrow \infty} NR[G_{1,\mu}] = Q.$$

Therefore, the separation of polygon  $\mathcal{P} = \bigcup_{\delta=1}^p Q_\delta$  into  $p$ -subpolygons leads to

$$\bigcup_{\delta=1}^p \left( \lim_{\mu \rightarrow \infty} NR[G_{1,\mu}^\delta] \right) = NR[A],$$

where  $G_{1,\mu}^\delta$  is a compression of associated  $Q_\delta$ , according to Theorem 4.

#### REFERENCES

- [1] M. ADAM AND J. MAROULAS, *On compressions of normal matrices*, Linear Algebra Appl., **341** (2002), 403–418.
- [2] M. ADAM AND P. PSARRAKOS, *On a compression of normal matrix polynomials*, Linear and Multilinear Algebra, **52**, 3-4 (2004), 251–263.
- [3] H.-L. GAU AND P.Y. WU, *Numerical range of a normal compression*, Linear and Multilinear Algebra, **52**, 3-4 (2004), 195–201.
- [4] H.-L. GAU AND P.Y. WU, *Numerical range of a normal compression II*, Linear Algebra Appl., **390** (2004), 121–136.
- [5] R.A. HORN AND C.R. JOHNSON, *Topics in Matrix Analysis*, Cambridge University Press, Cambridge, 1991.

(Received December 15, 2008)

Maria Adam  
Department of Computer Science and Biomedical Informatics  
University of Central Greece  
Lamia 35100, Greece  
e-mail: madam@ucg.gr

John Maroulas  
Department of Mathematics  
National Technical University  
Zografou Campus  
Athens 15780, Greece  
e-mail: maroulas@math.ntua.gr