

SELF-INVERSE MATRIX POLYNOMIALS WITH SEMISIMPLE SPECTRUM ON THE UNIT CIRCLE

NAOHARU ITO, REINHOLD KÜSTNER AND HARALD K. WIMMER

(Communicated by F. Kittaneh)

Abstract. The spectrum of a class of self-inverse matrix polynomials is studied. It is shown that the characteristic values are semisimple and lie on the unit circle if the inner radius of an associated matrix polynomial is greater than 1.

1. Introduction

The starting point of our paper is the following Eneström-Kakeya type theorem on self-inverse polynomials. It has an application in the theory of extremal self-dual codes [7].

THEOREM 1.1. *Let*

$$f(z) = a_0 + a_1z + \cdots + a_kz^k + a_kz^{m-k} + \cdots + a_1z^{m-1} + a_0z^m$$

be a real polynomial with $m > 2k$ and

$$a_0 > a_1 > \cdots > a_k > 0.$$

Then the zeros of $f(z)$ lie on the unit circle.

In this note we deal with matrix polynomials. Let

$$P(z) = A_0 + A_1z + \cdots + A_kz^k \in \mathbb{C}^{n \times n}[z] \tag{1.1}$$

be nonsingular (i.e. $\det P(z) \neq \text{zero polynomial}$). We say that λ is a *characteristic value* of $P(z)$ if $\det P(\lambda) = 0$, and we call

$$\sigma(P) = \{\lambda \in \mathbb{C}; \det P(\lambda) = 0\}$$

Mathematics subject classification (2010): 15A33, 15A24, 15A57, 26C10, 30C15, 39A11.

Keywords and phrases: Self-inverse polynomials, zeros of polynomials, matrix polynomials, numerical range, inner radius, semisimple characteristic values, normal characteristic values, Eneström-Kakeya theorem, system of linear difference equations.

The first author was supported by KAKENHI (19540128).

the spectrum of $P(z)$. A characteristic value λ is said to be *normal* if

$$P(\lambda)y = 0 \iff y^*P(\lambda) = 0,$$

$y \in \mathbb{C}^n$. Moreover, λ is called *semisimple* if the corresponding elementary divisors are linear. Let

$$W(P) = \{\lambda; v^*P(\lambda)v = 0 \text{ for some } v \in \mathbb{C}^n, v \neq 0\}$$

be the *numerical range* of $P(z)$. Then $W(P)$ is closed [17] and $\sigma(P) \subseteq W(P)$. The number

$$r_i(P) = \min\{|\lambda|; \lambda \in W(P)\}$$

is the *inner radius* of $P(z)$. Let $\mathbb{D} = \{\lambda; |\lambda| < 1\}$ and $\partial\mathbb{D} = \{\lambda; |\lambda| = 1\}$ be the open unit disc and the unit circle.

A generalization of Theorem 1.1 to matrix polynomials is part (i) of the following proposition.

PROPOSITION 1.2. *Let the coefficients A_0, \dots, A_k of the matrix polynomial*

$$F(z) = A_0 + A_1z + \dots + A_kz^k + A_kz^{m-k} + \dots + A_1z^{m-1} + A_0z^m, \tag{1.2}$$

$m > 2k$, be hermitian $n \times n$ matrices satisfying

$$A_0 > A_1 > \dots > A_k > 0. \tag{1.3}$$

(i) *Then the spectrum of $F(z)$ lies on the unit circle.* (ii) *All characteristic values of $F(z)$ are normal and semisimple.*

The preceding proposition follows from a general result on self-inversive matrix polynomials, which we state as Theorem 1.3 below and which is our main result. We adapt a definition of Marden [21], Sheil–Small [26] and Rahman and Schmeisser [23] from complex polynomials to matrix polynomials and say that

$$F(z) = F_0 + F_1z + \dots + F_{m-1}z^{m-1} + F_mz^m \in \mathbb{C}^{n \times n}[z] \tag{1.4}$$

with $F_0 \neq 0, F_m \neq 0$, is γ -self-inversive if

$$F(z) = \gamma(F_m^* + F_{m-1}^*z + \dots + F_1^*z^{m-1} + F_0^*z^m) \quad \text{and} \quad |\gamma| = 1.$$

To $F(z)$ in (1.4) we associate the conjugate-reverse matrix polynomial

$$\hat{F}(z) = F_m^* + \dots + F_1^*z^{m-1} + F_0^*z^m. \tag{1.5}$$

Thus, $F(z)$ is γ -self-inversive if $F(z) = \gamma\hat{F}(z)$.

THEOREM 1.3. *Let $P(z) = \sum_{j=0}^k A_jz^j \in \mathbb{C}^{n \times n}[z]$ be given with $A_k \neq 0, A_0 \neq 0$. If $|\gamma| = 1$ and $r \geq 0$ then*

$$F(z) = P(z) + \gamma z^r \hat{P}(z)$$

is γ -self-inversive.

(i) If $r_i(P) \geq 1$, i.e.

$$W(P) \subseteq \{\lambda; |\lambda| \geq 1\}, \tag{1.6}$$

then

$$\sigma(F) \subseteq W(F) \subseteq \partial\mathbb{D}, \tag{1.7}$$

and the characteristic values of $F(z)$ are normal.

(ii) If $r_i(P) > 1$ then the characteristic values of $F(z)$ lie on the unit circle, and they are normal and semisimple.

The proofs of Proposition 1.2 and Theorem 1.3 will be given in Section 4. They require a lower bound for the inner radius $r_i(P)$ in Section 2 and an auxiliary result on polynomials in Section 3.

Self-inversive polynomials and matrix polynomials can be found in the literature under various names including reciprocal [15], self-reciprocal [7], palindromic [30] and conjugate symmetric [3]. To place our results in a wider context we indicate some of the recent work in that area. We note that self-inversive polynomials have applications in numerous areas of engineering. They appear in optimal design of problems governed by hyperbolic field equations [30], in the study of line spectral pairs in speech coding [28, Chapter 9.11], in kernel representations of time-reversible systems [22]. Moreover, such polynomials are used in applied mathematics to deal with stability of periodic orbits of autonomous Hamiltonian systems [27, p. 159], and to investigate Lie algebras for semisimple hypersurface singularities [13]. Self-inversive polynomials with random coefficients were studied in the context of quantum chaotic dynamics in [4]. For many purposes (see e.g. [29, p. 108], [7], [13], [15]), [16]) a subclass of self-inversive polynomials is important, namely those polynomials which have all zeros on the unit circle. The study of self-inversive matrix polynomials started with [18]. An interesting example pointed out in [18] is the vibration analysis of railway tracks [11]. Matrix polynomials with self-inversive structure also arise in the solution of discrete-time linear-quadratic optimal control problems [6]. The papers [20], [1], [14], [19], [12] are concerned with Smith forms, and computational and algorithmic aspects of self-inversive matrix polynomials. Self-inversive systems of linear difference equations of the form

$$\begin{pmatrix} c & 0 \\ 0 & \bar{d} \end{pmatrix} x(t+2) - b \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} x(t+1) + \begin{pmatrix} \bar{c} & 0 \\ 0 & d \end{pmatrix} x(t) = 0, \tag{1.8}$$

with $c, d \in \mathbb{C}$, $b \in \mathbb{R}$, arise in the study of discretization schemes for the cubic Schrödinger equation [8, Section 2.2]. First order systems

$$A^*x(t+1) + Ax(t) = 0 \tag{1.9}$$

with bounded (and stably bounded) solutions were investigated in [24]. It is known [10] that stability properties of (1.8) or (1.9) are determined by the characteristic values of the corresponding matrix polynomials

$$F(z) = \begin{pmatrix} \bar{c} & 0 \\ 0 & d \end{pmatrix} - b \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} z + \begin{pmatrix} c & 0 \\ 0 & \bar{d} \end{pmatrix} z^2,$$

or $F(z) = A + A^*z$, respectively. In particular, all solutions are bounded if and only if the spectrum of $F(z)$ lies on the unit circle and the characteristic values are semisimple. In a subsequent paper we study self-inversive difference equations, in particular generalizations of equations of type (1.8). At this point let us only mention a stability result which is an immediate consequence of Proposition 1.2.

PROPOSITION 1.4. *Let A_0, A_1, \dots, A_k be hermitian $n \times n$ matrices. Suppose $A_0 > A_1 > \dots > A_k > 0$. Then the solution $x(t)$ of the difference equation*

$$A_0x(t+m) + A_1x(t+m-1) + \dots + A_kx(t+m-k) + A_kx(t+k) + \dots + A_1x(t+1) + A_0x(t) = 0, \quad m > 2k, \quad (1.10)$$

with initial conditions $x(0) = x_0, \dots, x(m-1) = x_{m-1}$, is bounded for $t \rightarrow \infty$ and $t \rightarrow -\infty$.

2. A lower bound for the inner radius

The Eneström–Kakeya theorem is frequently stated in the following form (due in fact to Eneström).

LEMMA 2.1. (see e.g. [5, p. 12]) *Let $p(z) = a_0 + a_1z + \dots + a_kz^k$ be a real polynomial with positive coefficients. Set*

$$\alpha(p) = \min_{0 \leq j \leq k-1} (a_j/a_{j+1}), \quad \beta(p) = \max_{0 \leq j \leq k-1} (a_j/a_{j+1}).$$

Then the zeros λ of $p(z)$ satisfy

$$\alpha(p) \leq |\lambda| \leq \beta(p).$$

If all eigenvalues of a matrix $M \in \mathbb{C}^{n \times n}$ are real then $\lambda_{\min}(M)$ shall denote the smallest eigenvalue of M . Let A and \tilde{A} be positive definite $n \times n$ matrices. If $v \in \mathbb{C}^n$, $v \neq 0$, then

$$\begin{aligned} \frac{v^* \tilde{A} v}{v^* A v} &= \frac{v^* A^{1/2} (A^{-1/2} \tilde{A} A^{-1/2}) A^{1/2} v}{v^* A^{1/2} A^{1/2} v} \geq \min_{w \in \mathbb{C}^n, w \neq 0} \frac{w^* (A^{-1/2} \tilde{A} A^{-1/2}) w}{w^* w} \\ &= \lambda_{\min}(A^{-1/2} \tilde{A} A^{-1/2}) = \lambda_{\min}(\tilde{A} A^{-1}). \end{aligned} \quad (2.1)$$

In [9, Theorem 2.6] we have a result which extends Lemma 2.1 to matrix polynomials. It is related to the following.

THEOREM 2.2. *Let the coefficients A_j , $j = 0, \dots, k$, of $P(z) = \sum_{j=0}^k A_j z^j$ be hermitian and positive definite. Set*

$$\mu(P) = \min \{ \lambda_{\min}(A_j A_{j+1}^{-1}); j = 0, \dots, k-1 \}.$$

Then $r_i(P) \geq \mu(P)$.

Proof. Let $v \in \mathbb{C}^n$, $v \neq 0$. Set $p_v(z) = v^*P(z)v$. Then (2.1) implies

$$\alpha(p_v) = \min_{0 \leq j \leq k-1} \frac{v^*A_j v}{v^*A_{j+1}v} \geq \min_{0 \leq j \leq k-1} \lambda_{\min}(A_j A_{j+1}^{-1}) = \mu(P).$$

Hence, if $\lambda \in W(P)$ and $p_v(\lambda) = 0$ then $\mu(P) \leq \alpha(p_v) \leq |\lambda|$, and therefore $\mu(P) \leq r_i(P)$. \square

COROLLARY 2.3. *If $A_0 > A_1 > \dots > A_k > 0$ then $r_i(P) > 1$.*

Proof. If $A_j > A_{j+1}$ then $A_{j+1}^{-1/2}A_j A_{j+1}^{-1/2} > I$. Hence $\lambda_{\min}(A_j A_{j+1}^{-1}) > 1$, and therefore $\mu(P) > 1$. \square

3. A theorem of Schur

Let $p(z) = a_0 + a_1z + \dots + a_kz^k$ be a complex polynomial with $a_k \neq 0$, $a_0 \neq 0$. In accordance with definition (1.5) we set

$$\hat{p}(z) = \bar{a}_k + \dots + \bar{a}_1z^{k-1} + \bar{a}_0z^k.$$

If $p(z) = c(z - \omega_1) \dots (z - \omega_k)$ then

$$\hat{p}(z) = \bar{c}(-1)^k \bar{\omega}_1 \dots \bar{\omega}_k \left(z - \frac{1}{\bar{\omega}_1}\right) \dots \left(z - \frac{1}{\bar{\omega}_k}\right).$$

Let $r \geq 0$ and $|\gamma| = 1$. Set

$$q(z) = \gamma z^r \hat{p}(z) \quad \text{and} \quad f(z) = p(z) + q(z).$$

Then $f(z) = \gamma \hat{f}(z)$. Thus we have $f(\varepsilon) = 0$ if and only if $f(\bar{\varepsilon}^{-1}) = 0$. To obtain information on zeros of $f(z)$ we consider the Blaschke product

$$g(z) = \frac{q(z)}{p(z)} = \gamma \bar{c} c^{-1} z^r \prod_{j=1}^k \frac{1 - z\bar{\omega}_j}{z - \omega_j}.$$

Set

$$b(\omega) = \frac{1 - z\bar{\omega}}{z - \omega}.$$

LEMMA 3.1. *If $|z| < 1$ then*

$$|\omega| > 1 \iff |b(\omega)| < 1, \tag{3.1}$$

and $|\omega| = 1 \iff |b(\omega)| = 1$.

Proof. The identity $|1 - z\bar{\omega}|^2 - |z - \omega|^2 = (1 - |z|^2)(1 - |\omega|^2)$ yields

$$|b(\omega)|^2 - 1 = \frac{1 - |z|^2}{|z - \omega|^2} (1 - |\omega|^2).$$

Hence $\operatorname{sgn}(|b(\omega)|^2 - 1) = \operatorname{sgn}(1 - |\omega|^2)$, if $|z| < 1$. \square

We extend a result of Schur [25, XII].

THEOREM 3.2. (i) *If the zeros ω_j of $p(z)$ satisfy $|\omega_j| \geq 1$, $j = 1, \dots, k$, then the zeros of $f(z) = p(z) + \gamma z^r \hat{p}(z)$ lie on the unit circle.*

(ii) *If*

$$|\omega_j| > 1, j = 1, \dots, k, \tag{3.2}$$

then all zeros of $f(z)$ are distinct.

Proof. (i) Suppose $|\omega_j| = 1$, $j = 1, \dots, k$. Then

$$\hat{p}(z) = \bar{c}c^{-1}(-1)^k \bar{\omega}_1 \cdots \bar{\omega}_k p(z),$$

and $f(z) = (1 + \tau z^r)p(z)$ for some $|\tau| = 1$. Hence all zeros of $f(z)$ have absolute value 1. Now, suppose $|\omega_j| > 1$ for some j . Let us show that $f(z) \neq 0$ if $z \in \mathbb{D}$. If $|z| < 1$ then (3.1) implies $|g(z)| < 1$. Hence $|q(z)| < |p(z)|$, and therefore

$$|f(z)| = |p(z) + q(z)| \geq \left| |p(z)| - |q(z)| \right| > 0.$$

Let $f(\varepsilon) = 0$. It is impossible that $|\varepsilon| > 1$. Otherwise we would have $f(\bar{\varepsilon}^{-1}) = 0$ and $|\bar{\varepsilon}^{-1}| < 1$. Thus we obtain $|\varepsilon| = 1$ if $f(\varepsilon) = 0$.

(ii) Let $f(\varepsilon) = 0$. Then $|\varepsilon| = 1$. The assumption (3.2) implies $p(\varepsilon) \neq 0$. Then $p(\varepsilon) = -q(\varepsilon)$ yields

$$\begin{aligned} \frac{f'(\varepsilon)}{p(\varepsilon)} &= \frac{p'(\varepsilon)}{p(\varepsilon)} - \frac{q'(\varepsilon)}{q(\varepsilon)} = \sum_{j=1}^k \frac{1}{\varepsilon - \omega_j} - \sum_{j=1}^k \frac{1}{\varepsilon - \bar{\omega}_j^{-1}} - \frac{r}{\varepsilon} \\ &= \sum_{j=1}^k \frac{|\omega_j|^2 - 1}{|\varepsilon - \omega_j|^2(-\varepsilon)} - \frac{r}{\varepsilon} = -\varepsilon^{-1}(s + r), \end{aligned}$$

where $s = \sum_{j=1}^k (|\omega_j|^2 - 1)|\varepsilon - \omega_j|^{-2}$. Condition (3.2) implies $s > 0$. Hence $f'(\varepsilon) \neq 0$, and therefore the zeros of $f(z)$ are simple. \square

4. Proofs

We show first that the unimodular characteristic values of self-inversive matrix polynomials are normal, but not necessarily semisimple.

LEMMA 4.1. *Let $F(z) \in \mathbb{C}^{n \times n}[z]$ be self-inversive. Then the characteristic values of $F(z)$ on the unit circle are normal.*

Proof. Suppose $F(z) = \sum_{i=0}^m F_i z^i$, $F_m \neq 0$. Let $\lambda \in \sigma(F)$, $|\lambda| = 1$, and $F(\lambda)v = 0$, $v \neq 0$. Then

$$\begin{aligned} 0 &= v^*F(\lambda)^* = \gamma\lambda^{-m}v^*[\bar{\gamma}(F_m^* + F_{m-1}^*\bar{\lambda} + \dots + F_0^*\bar{\lambda}^{-m})] = \gamma\lambda^{-m}v^*\bar{\gamma}\hat{F}(\lambda) \\ &= \gamma\lambda^{-m}v^*F(\lambda). \end{aligned}$$

Therefore $F(\lambda)v = 0$ is equivalent to $v^*F(\lambda) = 0$, and λ is normal. \square

EXAMPLE 4.2. The linear pencil

$$F(z) = \begin{pmatrix} 0 & 0 & 1+z & 1 \\ 0 & 0 & 0 & 1+z \\ 1+z & 0 & 0 & 0 \\ z & 1+z & 0 & 0 \end{pmatrix}$$

is self-inversive. We have $\sigma(F) = \{-1\}$ and

$$\text{Ker } F(-1) = \text{span}\{(0, 1, 0, 0)^T, (0, 0, 1, 0)^T\} = \text{Ker } F(-1)^*.$$

The Smith form of $F(z)$ is $\text{diag}((1+z)^2, (1+z)^2, 1, 1)$. Hence $-1 \in \sigma(F) \cap \partial\mathbb{D}$ is not semisimple.

Proof of Theorem 1.3. (i) Let $\lambda \in W(F)$ and $v^*F(\lambda)v = 0$, $v \neq 0$. Set

$$p_v(z) = v^*P(z)v \quad \text{and} \quad f_v(z) = v^*F(z)v.$$

Then $f_v(z) = p_v(z) + \gamma z^r \hat{p}_v(z)$. The assumption (1.6) implies that $|\omega| \geq 1$ if $p_v(\omega) = 0$. Hence Theorem 3.2(i) yields $\lambda \in \partial\mathbb{D}$, and we obtain (1.7). Let $\lambda \in \sigma(F)$. Then (1.7) implies $|\lambda| = 1$. Hence λ is normal (by Lemma 4.1).

(ii) Suppose $\lambda \in \sigma(F)$ is not semisimple. Then (see e.g. [2], [10]) there exists a corresponding Jordan chain of length 2. That is, $F(\lambda)v = 0$, $v \neq 0$, and

$$F'(\lambda)v + F(\lambda)w = 0, \tag{4.1}$$

for some $v, w \in \mathbb{C}^n$. Consider the polynomial $p_v(z) = v^*F(z)v$. Now the stronger assumption $r_i > 1$ implies $|\omega| > 1$ if $p_v(\omega) = 0$. Therefore (by Theorem 3.2(ii)) all zeros of $f_v(z) = v^*F(z)v$ are simple. We know that λ is a normal characteristic value. Hence $v^*F(\lambda) = 0$. Then (4.1) yields $f'_v(\lambda) = v^*F'(\lambda)v = 0$, and λ would be a multiple zero of $f_v(z)$, which is a contradiction. \square

Proof of Proposition 1.2. We have $F(z) = P(z) + \hat{P}(z)$. The coefficients of $P(z)$ satisfy (1.3). From Corollary 2.3 we obtain $r_i(P) > 1$. Thus we can apply Theorem 1.3(ii). \square

REFERENCES

- [1] B. ADHIKARI AND R. ALAM, *Structured backward errors and pseudospectra of structured matrix pencils*, SIAM J. Matrix Anal. Appl., **31** (2009), 331–359.
- [2] H. BAUMGÄRTEL, *Analytic Perturbation Theory for Matrices and Operators*, Operator Theory, Advances and Applications, Vol. 15, Birkhäuser, Basel, 1985.
- [3] Y. BISTRITZ, *A circular stability test for general polynomials*, Systems Control Lett., **7** (1986), 89–97.
- [4] E. BOGOMOLNY, O. BOHIGAS, AND P. LEBOEUF, *Quantum chaotic dynamics and random polynomials*, J. Statistical Physics, **85** (1996), 639–679.
- [5] P. BORWEIN AND T. ERDÉLYI, *Polynomials and Polynomial Inequalities*, Springer, New York, 1995.
- [6] R. BYERS, D. S. MACKEY, V. MEHRMANN, AND H. XU, *Symplectic, BVD, and palindromic approaches to discrete-time control problems*, in P. Petkov and N. Christov, editors, A Collection of Papers Dedicated to the 60th Anniversary of Mihail Konstantinov, pp. 81–102, Sofia, 2009, Publ. House Rodina. http://eprints.ma.man.ac.uk/1062/01/ByeMMX08.ppt-final_26-03-2008_00-26-49.pdf
- [7] K. CHINEN, *An abundance of polynomials satisfying the Riemann hypothesis*, Discrete Math., **308** (2008), 6426–6440.
- [8] M. DAHLBY AND B. OWREN, *Plane wave stability of some conservative schemes for the cubic Schrödinger equation*, ESAIM: M2AN, **43** (2009), 677–687.
- [9] G. DIRR AND H. K. WIMMER, *An Eneström-Kakeya theorem for hermitian polynomial matrices*, IEEE Trans. Automat. Control, **52** (2007), 2151–2153.
- [10] I. GOHBERG, P. LANCASTER AND L. RODMAN, *Matrix Polynomials*, Academic Press, New York, 1982.
- [11] A. HILLIGES, *Numerische Lösung von quadratischen Eigenwertproblemen mit Anwendung in der Schienendynamik*, Diplomarbeit, TU Berlin, Institut f. Mathematik, 2004.
- [12] B. IANNAZZO AND B. MEINI, *Palindromic matrix polynomials, matrix functions and integral representations*, Linear Algebra Appl., **434** (2011), 174–184.
- [13] M. JIBLADZE AND D. NOVIKOV, *Unimodularity of Poincaré polynomials of Lie algebras for semisimple singularities*, Moscow Math. J., **7** (2007), 481–487.
- [14] D. KRESSNER, CH. SCHROEDER, AND D. S. WATKINS, *Implicit QR algorithms for palindromic and even eigenvalue problems*, Numer. Algorithms, **51** (2009), 209–238.
- [15] P. LAKATOS, *On polynomials with zeros on the unit circle*, C. R. Math. Acad. Sci. Soc. R. Can., **24** (2002), 91–96.
- [16] P. LAKATOS AND L. LOSONCZI, *Polynomials with all zeros on the unit circle*, Acta Math. Hung., **125** (2009), 341–356.
- [17] CH.-K. LI AND L. RODMAN, *Numerical range of matrix polynomials*, SIAM J. Matrix Anal. Appl., **15** (1994), 1256–1265.
- [18] D. ST. MACKEY, N. MACKEY, CH. MEHL, AND V. MEHRMANN, *Structured polynomial eigenvalue problems: good vibrations from good linearizations*, SIAM J. Matrix Anal. Appl., **28** (2006), 1029–1051.
- [19] D. ST. MACKEY, N. MACKEY, CH. MEHL, AND V. MEHRMANN, *Numerical methods for palindromic eigenvalue problems: Computing the anti-triangular Schur form*, Numer. Linear Algebra Appl., **16** (2009), 63–86.
- [20] D. ST. MACKEY, N. MACKEY, CH. MEHL, AND V. MEHRMANN, *Smith forms of palindromic matrix polynomials*, Electron. J. Linear Algebra, **22** (2011), 53–91.
- [21] M. MARDEN, *Geometry of Polynomials*, 3rd ed., Mathematical Surveys No. 3, Amer. Math. Soc., Providence, RI, 1966.
- [22] I. MARKOVSKY AND S. RAO, *Palindromic polynomials, time-reversible systems, and conserved quantities*, in: 16th Mediterranean Conference on Control and Automation, 2008, Ajaccio. <http://eprints.ecs.soton.ac.uk/16592/1/Med08.pdf>
- [23] Q. I. RAHMAN AND G. SCHMEISSER, *Analytic Theory of Polynomials*, Oxford University Press, Oxford, 2002.
- [24] L. RODMAN, *Bounded and stably bounded palindromic difference equations of first order*, Electron. J. Linear Algebra, **15** (2006), 22–49.
- [25] J. SCHUR, *Über Potenzreihen, die im Innern des Einheitskreises beschränkt sind*, J. reine angew. Math., **147** (1917), 205–232.

- [26] T. SHEIL–SMALL, *Complex Polynomials*, Cambridge University Press, Cambridge, 2002.
- [27] CH. SKOKOS, *On the stability of periodic orbits of high dimensional autonomous Hamiltonian systems*, *Physica D*, **159** (2001), 155–179.
- [28] J. Y. STEIN, *Digital Signal Processing: A Computer Science Perspective*, Wiley Series in Telecommunications and Signal Processing, Wiley, 2000.
- [29] P. STOICA AND R. L. MOSES, *On the unit circle problem: The Schur–Cohn procedure revisited*, *Signal Processing*, **26** (1992), 95–118.
- [30] A. P. VELO, G. A. GAZONAS, AND T. AMEYA, *z-Transform methods for the optimal design of one-dimensional layered elastic media*, *SIAM J. Appl. Math.* **70**, 3 (2009), 762–788.

(Received November 3, 2010)

Naoharu Ito
Department of Mathematical Education
Nara University of Education
Takabatake-cho, Nara-shi
Nara 630-8528
Japan
e-mail: naoharu@nara-edu.ac.jp

Reinhold Küstner
Mathematisches Institut
Universität Würzburg
97074 Würzburg
Germany
e-mail: kuestner@mathematik.uni-kassel.de

Harald K. Wimmer
Mathematisches Institut
Universität Würzburg
97074 Würzburg
Germany
e-mail: wimmer@mathematik.uni-wuerzburg.de