

## TRACIAL MOMENT PROBLEMS ON HYPERCUBES

CONG TRINH LE

*Dedicated to Prof. Ha Huy Vui on the occasion of his 70th birthday*

*(Communicated by D. R. Farenick)*

*Abstract.* In this paper we introduce the *tracial  $K$ -moment problem* and the *sequential matrix-valued  $K$ -moment problem* and show the equivalence of the solvability of these problems. Using a Haviland's theorem for matrix polynomials, we solve these  $K$ -moment problems for the case where  $K$  is the hypercube  $[-1, 1]^n$ .

### 1. Introduction

Let  $\mathbb{R}[X] := \mathbb{R}[X_1, \dots, X_n]$  denote the algebra of polynomials in  $n$  variables  $X_1, \dots, X_n$  with real coefficients.

For a real-valued linear functional  $L$  on  $\mathbb{R}[X]$  and a closed subset  $K \subseteq \mathbb{R}^n$ , the  *$K$ -moment problem* asks *when does there exist a positive Borel measure  $\sigma$  on  $\mathbb{R}^n$  supported on  $K$  such that*

$$L(f) = \int f(x) d\sigma(x), \quad \forall f \in \mathbb{R}[X]?$$

Each functional of this form is called a  *$K$ -moment functional*.

Another kind of  $K$ -moment problems for sequences of numbers is stated as follows. For a multi-index sequence  $s = (s_\alpha)_{\alpha \in \mathbb{N}_0^n}$ , the  *$K$ -moment problem for sequences* asks *when does there exist a positive Borel measure  $\sigma$  on  $\mathbb{R}^n$  supported on  $K$  such that*

$$s_\alpha = \int x^\alpha d\sigma(x), \quad \forall \alpha \in \mathbb{N}_0^n,$$

where  $x^\alpha = x_1^{\alpha_1} \dots x_n^{\alpha_n}$ . Each sequence of this form is called a  *$K$ -moment sequence*.

For each sequence  $s = (s_\alpha)_{\alpha \in \mathbb{N}_0^n}$ , the *Riesz functional*  $L_s$  on  $\mathbb{R}[X]$  associated to  $s$  is defined by

$$L_s(x^\alpha) = s_\alpha, \quad \forall \alpha \in \mathbb{N}_0^n.$$

---

*Mathematics subject classification* (2010): 44A60, 30E05, 13J30, 47A57, 11E25.

*Keywords and phrases:* Moment problem, matrix polynomial, moment functional, moment sequence, hypercube.

This research is funded by the Vietnam Ministry of Education and Training under the decision number 3813/QĐ-BGDĐT dated November 20, 2020.

It is easy to check that  $s$  is a  $K$ -moment sequence if and only if  $L_s$  is a  $K$ -moment functional.

In the case  $K = \mathbb{R}$  (resp.  $[0, +\infty)$ ,  $[0, 1]$ ), we go back to the *Hamburger* (resp. *Stieltjes*, *Hausdorff*) *moment problem* which was solved completely by many authors (see, for example, a very excellent book of Schmüdgen [10] on the moment problem). If  $K$  is a *compact basic closed semi-algebraic set* in  $\mathbb{R}^n$ , the  $K$ -moment problem was solved by Schmüdgen (1991, [9]).

One of the power tools to solve  $K$ -moment problems is the *Haviland theorem*, which was proved by E.K. Haviland (1935) and stated as follows.

*Given a linear functional  $L$  on  $\mathbb{R}[X]$  and a closed subset  $K$  in  $\mathbb{R}^n$ . Then  $L$  is a  $K$ -moment functional if and only if  $L(f) \geq 0$  for all  $f \geq 0$  on  $K$ .*

Here the notation  $f \geq 0$  on  $K$  means that  $f(x) \geq 0, \forall x \in K$ . A similar meaning is defined for the notation  $f > 0$  on  $K$ .

For a fixed integer  $t > 0$ , we denote by  $M_t(\mathbb{R}[X])$  the algebra of  $t \times t$  matrices with entries in  $\mathbb{R}[X]$ , and by  $S_t(\mathbb{R}[X])$  the subalgebra of symmetric matrices. Each element  $\mathbf{A} \in M_t(\mathbb{R}[X])$  is a matrix whose entries are polynomials in  $\mathbb{R}[X]$ , which is called a *polynomial matrix*.  $\mathbf{A}$  is also called a *matrix polynomial*, because it can be viewed as a polynomial in  $X_1, \dots, X_n$  whose coefficients come from  $M_t(\mathbb{R})$ . Namely, we can write  $\mathbf{A}$  as

$$\mathbf{A}(X) = \sum_{|\alpha|=0}^d \mathbf{A}_\alpha X^\alpha,$$

where  $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}_0^n$ ,  $|\alpha| := \alpha_1 + \dots + \alpha_n$ ,  $X^\alpha := X_1^{\alpha_1} \dots X_n^{\alpha_n}$ ,  $\mathbf{A}_\alpha \in M_t(\mathbb{R})$ . To unify notation, throughout the paper each element of  $M_t(\mathbb{R}[X])$  is called a *matrix polynomial*.

Let  $\mathbf{A}(X) = \sum_{|\alpha|=0}^d \mathbf{A}_\alpha X^\alpha$  be a matrix polynomial. Denote

$$\mathbf{A}_d(X) := \sum_{|\alpha|=d} \mathbf{A}_\alpha X^\alpha.$$

If  $\mathbf{A}_d \neq 0$ , the matrix polynomial  $\mathbf{A}$  is called of *degree  $d$* . In particular, when  $n = 1$  and  $\mathbf{A}(x) = \sum_{i=0}^d \mathbf{A}_i x^i$  is a univariate matrix polynomial of degree  $d$ , the non-zero matrix  $\mathbf{A}_d$  is called the *leading coefficient* of  $\mathbf{A}$ .

Recently, there are some research on (truncated) *non-commutative  $K$ -moment problems*, e.g. Dette and Studden (2002, [4]), Burgdorf and Klep (2011, [1]), Burgdorf, Klep and Povh (2013, [2]), Cimpric and Zalar (2013, [3]), Kimsey and Woerdeman (2013, [6]), Kimsey (2011, [5]) and the references therein. In particular, in [3], Cimpric and Zalar proved a version of the Haviland theorem for matrix polynomials, and they applied this development to solve the matrix-valued moment problem on compact basic closed semi-algebraic sets.

In the case where  $K \subseteq \mathbb{R}^n$  is a basic closed semi-algebraic set defined by a subset  $G = \{g_1, \dots, g_m\}$  of  $\mathbb{R}[X]$ , almost all solutions of the matrix-valued  $K$ -moment problems are obtained using the positivity of the given linear functional on the quadratic

module  $M(G)^t$  in  $M_t(\mathbb{R}[X])$ , where  $M(G)$  is the quadratic module in  $\mathbb{R}[X]$  generated by  $G$  and

$$M(G)^t = \left\{ \sum_{i=1}^r m_i \mathbf{A}_i^T \mathbf{A}_i \mid r \in \mathbb{N}, m_i \in M(G), \mathbf{A}_i \in M_t(\mathbb{R}[X]) \right\}.$$

The main aim of this paper is to introduce some kinds of  $K$ -moment problems for matrix polynomials and solve these matrix-valued  $K$ -moment problems in the case where  $K$  is the hypercube  $[-1, 1]^n$ . The main advantage working on the hypercube  $[-1, 1]^n$  (and in general, on convex, compact polyhedra with non-empty interior) is that any matrix polynomial which is positive definite on  $[-1, 1]^n$  can be represented in terms of positive definite scalar matrices (see Theorem 3.1 for the case  $n = 1$  and Corollary 4.1.1 for  $n \geq 1$ ). Then the solution of the matrix-valued  $[-1, 1]^n$ -moment problem is obtained based on the positivity of the given linear functional on the cone of positive definite scalar matrices, which is well described.

The paper is organized as follows. In Section 2 we introduce the definition of the *tracial  $K$ -moment problem* and the *sequential matrix-valued  $K$ -moment problem* and establish an equivalence of the solvability of these problems (Proposition 1). Moreover, we also prove in this section a version of Haviland’s theorem for matrix polynomials which was proved mainly by Cimprič and Zalar in [3]. Next, in Section 3 we solve the tracial (resp. sequential matrix-valued)  $[-1, 1]$ -moment problem, which is established based on a representation of univariate matrix polynomials positive definite on  $[-1, 1]$ . A solution of the tracial (resp. sequential matrix-valued)  $[0, 1]$ -moment problem is also given in this section. Finally, in Section 4 we solve the tracial (resp. sequential matrix-valued) moment problem on the general hypercube  $[-1, 1]^n$ .

NOTATION. For any matrix  $\mathbf{A} \in M_t(\mathbb{R}[X])$ , the notation  $\mathbf{A} \geq 0$  means  $\mathbf{A}$  is *positive semidefinite*, i.e. for each  $x \in \mathbb{R}^n$ ,  $v^T \mathbf{A}(x)v \geq 0$  for all  $v \in \mathbb{R}^t$ ;  $\mathbf{A} > 0$  means  $\mathbf{A}$  is *positive definite*, i.e. for each  $x \in \mathbb{R}^n$ ,  $v^T \mathbf{A}(x)v > 0$  for all  $v \in \mathbb{R}^t \setminus \{0\}$ .

## 2. Haviland’s theorem for matrix polynomials

### 2.1. Matrix-valued measures and integrals

In this section we recall some basic notions of the matrix-valued measures and integrals on a closed subset of  $\mathbb{R}^n$ , which can be seen from the thesis of Kimsey [5].

Throughout this subsection, let  $X \subseteq \mathbb{R}^n$  be a non-empty closed set.

DEFINITION 1. Denote by  $\mathcal{B}(X)$  the smallest  $\sigma$ -algebra generated from the open (or equivalently closed) subsets of  $X$ . A measure  $\sigma$  defined on  $\mathcal{B}(X)$  is called a *Borel measure*. A Borel measure  $\sigma$  on  $\mathcal{B}(X)$  is called *finite* if  $\sigma(X) < +\infty$ . Denote by  $m(X)$  the set of all finite measures on  $X$ .

A measure  $\sigma \in m(X)$  is called *positive* if  $\sigma(E) \geq 0$  for all  $E \in \mathcal{B}(X)$ . The set of all finite positive Borel measures on  $X$  is denoted by  $m^+(X)$ .

For each  $\sigma \in m(X)$ , the *support* of  $\sigma$  is defined by

$$\text{supp}(\sigma) := \overline{\{E \in \mathcal{B}(X) : |\sigma|(E) > 0\}},$$

where  $|\sigma|(E) := |\sigma(E)|$  for all  $E \in \mathcal{B}(X)$ .

DEFINITION 2. Let  $t$  be a positive integer. Let  $\sigma_{ij} \in m(X)$ ,  $i, j = 1, \dots, t$ . Define the matrix-valued function  $\mathbf{E} : \mathcal{B}(X) \rightarrow M_t(\mathbb{R})$  by

$$\mathbf{E}(A) := (\sigma_{ij}(A))_{i,j=1,\dots,t} \in M_t(\mathbb{R}), \forall A \in \mathcal{B}(X).$$

The matrix-valued function  $\mathbf{E}$  defined by this way is called a *matrix-valued measure* on  $X$ . The set of all matrix-valued measure on  $X$  is denoted by  $M(X)$ . The set

$$\text{supp}(\mathbf{E}) := \bigcup_{i,j=1}^t \text{supp}(\sigma_{ij})$$

is called the *support of the matrix-valued measure*  $\mathbf{E}$ .

If  $\sigma_{ij} = \sigma_{ji}$  for all  $i, j = 1, \dots, t$ , we say that  $\mathbf{E}$  is a *symmetric measure*. In addition, if for all  $A \in \mathcal{B}(X)$  and for all  $v \in \mathbb{R}^t$  we have  $v^T \mathbf{E}(A)v \geq 0$ , then  $\mathbf{E}$  is called a *positive semidefinite matrix-valued measure*. The set of all positive semidefinite matrix-valued measures on  $X$  is denoted by  $M^+(X)$ .

DEFINITION 3. Let  $\mathbf{E} = (\sigma_{ij}) \in M(X)$ . A function  $f : X \rightarrow \mathbb{R}$  is called  *$\mathbf{E}$ -measurable* if  $f$  is  $\sigma_{ij}$ -measurable for every  $i, j = 1, \dots, t$ .

Let  $\mathbf{E} = (\sigma_{ij}) \in M(X)$  and  $f : X \rightarrow \mathbb{R}$  be  $\mathbf{E}$ -measurable. The *matrix-valued integral* of  $f$  with respect to the matrix measure  $\mathbf{E}$  is defined by

$$\int_X f(x)d\mathbf{E}(x) := \left( \int_X f(x)d\sigma_{ij}(x) \right)_{i,j=1,\dots,t} \in M_t(\mathbb{R}).$$

### 2.2. Tracial $K$ -moment problems

Let  $\mathcal{L}$  be a real-valued linear functional on  $M_t(\mathbb{R}[X])$ , where  $\mathbb{R}[X] := \mathbb{R}[X_1, \dots, X_n]$  and  $K \subseteq \mathbb{R}^n$  a closed subset.

DEFINITION 4. (cf. [3])  $\mathcal{L}$  is called a *tracial  $K$ -moment functional* if there exists a positive semidefinite matrix-valued  $\mathbf{E} \in M^+(\mathbb{R}^n)$  such that

$$\text{supp}(\mathbf{E}) \subseteq K \text{ and } \mathcal{L}(\mathbf{F}) = \int \text{tr}(\mathbf{F}(x)d\mathbf{E}(x)), \forall \mathbf{F} \in M_t(\mathbb{R}[X]). \tag{2.1}$$

The matrix-valued measure  $\mathbf{E} \in M^+(\mathbb{R}^n)$  satisfying (2.1) is called a *representing measure* of the tracial  $K$ -moment functional  $\mathcal{L}$ .

PROBLEM 1. Let  $\mathcal{L}$  be a linear functional on  $M_t(\mathbb{R}[X])$  and  $K \subseteq \mathbb{R}^n$  a closed set. The *tracial  $K$ -moment problem* asks when does there exist a matrix-valued measure  $\mathbf{E} \in M^+(\mathbb{R}^n)$  satisfying the conditions (2.1)?

The following definition of matrix-valued  $K$ -moment sequences is learned from Kimsey in [5].

DEFINITION 5. Let  $\mathbf{S} = (\mathbf{S}_\alpha)_{\alpha \in \mathbb{N}_0^n}$  be a multi-indexed sequence of symmetric matrices in  $M_t(\mathbb{R})$  and  $K \subseteq \mathbb{R}^n$  a closed subset. The sequence  $\mathbf{S}$  is called a *matrix-valued  $K$ -moment sequence* if there exists a positive semidefinite matrix-valued  $\mathbf{E} \in M^+(\mathbb{R}^n)$  such that

$$\text{supp}(\mathbf{E}) \subseteq K \text{ and } \mathbf{S}_\alpha = \int x^\alpha d\mathbf{E}(x), \forall \alpha \in \mathbb{N}_0^n, \tag{2.2}$$

where  $x^\alpha = x_1^{\alpha_1} \dots x_n^{\alpha_n}$  with  $\alpha = (\alpha_1, \dots, \alpha_n)$  and  $x = (x_1, \dots, x_n) \in \mathbb{R}^n$ .

The matrix-valued measure  $\mathbf{E} \in M^+(\mathbb{R}^n)$  satisfying (2.2) is called a *representing measure* of the matrix-valued  $K$ -moment sequence  $\mathbf{S}$ .

PROBLEM 2. Let  $\mathbf{S} = (\mathbf{S}_\alpha)_{\alpha \in \mathbb{N}_0^n}$  be a multi-indexed sequence of symmetric matrices in  $M_t(\mathbb{R})$  and  $K \subseteq \mathbb{R}^n$  a closed set. *The sequential matrix-valued  $K$ -moment problem* asks when does there exist a matrix-valued measure  $\mathbf{E} \in M^+(\mathbb{R}^n)$  satisfying the conditions (2.2)?

For each matrix-valued sequence  $\mathbf{S} = (\mathbf{S}_\alpha)_{\alpha \in \mathbb{N}_0^n}$ , let us consider the real-valued function  $\mathcal{L}_\mathbf{S}$  on  $M_t(\mathbb{R}[X])$  defined as follows. For each  $\mathbf{F}(X) = \sum_{|\alpha|=0}^d \mathbf{F}_\alpha X^\alpha \in M_t(\mathbb{R}[X])$ , define

$$\mathcal{L}_\mathbf{S}(\mathbf{F}) := \sum_{|\alpha|=0}^d \text{tr}(\mathbf{F}_\alpha \mathbf{S}_\alpha).$$

It is easy to check that  $\mathcal{L}_\mathbf{S}$  is *linear*, which is called the *Riesz functional associated to the sequence  $\mathbf{S}$* .

PROPOSITION 1. Let  $\mathbf{S} = (\mathbf{S}_\alpha)_{\alpha \in \mathbb{N}_0^n}$  be a sequence of symmetric matrices in  $M_t(\mathbb{R})$  and  $K \subseteq \mathbb{R}^n$  a closed set. Then  $\mathbf{S}$  is a matrix-valued  $K$ -moment sequence if and only if  $\mathcal{L}_\mathbf{S}$  is a *tracial  $K$ -moment functional*.

*Proof.* Assume that  $\mathbf{S}$  is a matrix-valued  $K$ -moment sequence. Then there exists a matrix-valued measure  $\mathbf{E}$  in  $M^+(\mathbb{R}^n)$  supported in  $K$  and

$$\mathbf{S}_\alpha = \int x^\alpha d\mathbf{E}(x), \forall \alpha \in \mathbb{N}_0^n.$$

Then, for each  $\mathbf{F}(X) = \sum_{|\alpha|=0}^d \mathbf{F}_\alpha X^\alpha \in M_t(\mathbb{R}[X])$ , by definition of  $\mathcal{L}_\mathbf{S}$ , we have

$$\begin{aligned} \mathcal{L}_\mathbf{S}(\mathbf{F}) &= \sum_\alpha \text{tr}(\mathbf{F}_\alpha \mathbf{S}_\alpha) = \sum_\alpha \text{tr} \left( \int x^\alpha \mathbf{F}_\alpha d\mathbf{E}(x) \right) \\ &= \int \text{tr} \left( \sum_\alpha \mathbf{F}_\alpha x^\alpha d\mathbf{E}(x) \right) = \int \text{tr}(\mathbf{F}(x) d\mathbf{E}(x)). \end{aligned}$$

This implies that  $\mathcal{L}_\mathbf{S}$  is a *tracial  $K$ -moment functional*.

Conversely, assume that  $\mathbf{E}$  is a matrix-valued measure in  $M^+(\mathbb{R}^n)$  satisfying

$$\text{supp}(\mathbf{E}) \subseteq K \text{ and } \mathcal{L}_{\mathbf{S}}(\mathbf{F}) = \int \text{tr}(\mathbf{F}(x)d\mathbf{E}(x)), \forall \mathbf{F} \in M_t(\mathbb{R}[X]).$$

In particular, for each coordinate matrix  $\mathbf{W}_{k,l}$  of the algebra  $M_t(\mathbb{R})$  and each  $\alpha \in \mathbb{N}_0^n$ , we have

$$\mathcal{L}_{\mathbf{S}}(X^\alpha \mathbf{W}_{k,l}) = \int x^\alpha \text{tr}(\mathbf{W}_{k,l}d\mathbf{E}(x)) = \int x^\alpha d\mathbf{E}_{l,k}.$$

Here we use the fact that for each  $k, l = 1, \dots, t$  and for each matrix  $\mathbf{A} = (\mathbf{A}_{i,j})_{i,j} \in M_t(\mathbb{R})$ , we have

$$\text{tr}(\mathbf{W}_{k,l}\mathbf{A}) = \mathbf{A}_{l,k}.$$

On the other hand, by definition of the associated Riesz functional, we have

$$\mathcal{L}_{\mathbf{S}}(X^\alpha \mathbf{W}_{k,l}) = \text{tr}(\mathbf{W}_{k,l}\mathbf{S}_\alpha) = (\mathbf{S}_\alpha)_{l,k}.$$

It follows that

$$\mathbf{S}_\alpha^T = ((\mathbf{S}_\alpha)_{l,k})_{k,l} = \left( \int x^\alpha d\mathbf{E}_{l,k}(x) \right)_{k,l} = \left( \int x^\alpha d\mathbf{E}(x) \right)^T.$$

Equivalently,  $\mathbf{S}_\alpha = \int x^\alpha d\mathbf{E}(x)$ , i.e.  $\mathbf{S}$  is a matrix-valued  $K$ -moment sequence.  $\square$

### 2.3. Haviland’s theorem for matrix polynomials

**THEOREM 2.1.** *Let  $\mathcal{L}$  be a real-valued linear functional on  $M_t(\mathbb{R}[X])$  and  $K \subseteq \mathbb{R}^n$  a closed set. Then the following are equivalent.*

- (1)  $\mathcal{L}$  is a tracial  $K$ -moment functional, i.e. there exists a positive semi-definite matrix-valued measure  $\mathbf{E}$  on  $\mathbb{R}^n$  whose support is contained in  $K$  such that

$$\mathcal{L}(\mathbf{F}) = \int \text{tr}(\mathbf{F}d\mathbf{E}).$$

- (2)  $\mathcal{L}(\mathbf{F}) \geq 0$  for all  $\mathbf{F} \geq 0$  on  $K$ .
- (3)  $\mathcal{L}(\mathbf{F} + \varepsilon \mathbf{I}) \geq 0$  for all  $\mathbf{F} \geq 0$  on  $K$  and for all  $\varepsilon > 0$ .
- (4)  $\mathcal{L}(\mathbf{F}) \geq 0$  for all  $\mathbf{F} > 0$  on  $K$ .

*Proof.* The equivalence of (1) and (2) was proved by Cimpric and Zalar [3, Theorem 3 and Remark 5].

Assume that (2) is satisfied. Then  $\mathcal{L}(\mathbf{I}) \geq 0$ , since  $\mathbf{I} \geq 0$  on  $K$ . It follows (3) by the linearity of  $\mathcal{L}$  on  $M_t(\mathbb{R}[X])$ .

Now we check (3)  $\Rightarrow$  (2). Take  $\mathbf{F} \geq 0$  on  $K$ . Consider  $\varepsilon = \frac{1}{n}$  for  $n \in \mathbb{N}$ . Then by the assumption of (3),

$$0 \leq \mathcal{L}\left(\mathbf{F} + \frac{1}{n}\mathbf{I}\right) = \mathcal{L}(\mathbf{F}) + \frac{1}{n}\mathcal{L}(\mathbf{I}),$$

where  $\mathcal{L}(\mathbf{I}) \geq 0$  by hypothesis. Letting  $n \rightarrow +\infty$ , we get  $\mathcal{L}(\mathbf{F}) \geq 0$ , i.e. we have (2).

It is obvious that (2)  $\Rightarrow$  (4). Now we verify (4)  $\Rightarrow$  (3). Take  $\mathbf{F} \geq 0$  on  $K$  and  $\varepsilon > 0$ . Then  $\mathbf{F} + \varepsilon \mathbf{I} > 0$  on  $K$ . Hence by the assumption we have  $\mathcal{L}(\mathbf{F} + \varepsilon \mathbf{I}) \geq 0$ , i.e. we have (3). The proof is complete.  $\square$

### 3. Tracial moment problems on the intervals $[-1, 1]$ and $[0, 1]$

Firstly we propose a version of the Bernstein theorem (cf. [10, Prop. 3.4]) for matrix polynomials, representing a matrix polynomial positive definite on the interval  $[-1, 1]$ . This is a special case of the Handelman’s Positivstellensatz for matrix polynomials established in [7], but the proof given here is special for the case of one-variable matrix polynomials.

**THEOREM 3.1.** *Let  $\mathbf{F} \in M_r(\mathbb{R}[x])$  be a matrix polynomial of degree  $d > 0$ . Assume  $\mathbf{F}(x) > 0$  for all  $x \in [-1, 1]$ . Then there exists a positive number  $N \in \mathbb{N}$  and positive definite matrices  $\mathbf{G}_i, i = 1, \dots, N + d$  such that*

$$\mathbf{F}(x) = \sum_{i=0}^{N+d} \mathbf{G}_i (1+x)^i (1-x)^{N+d-i}.$$

The main idea of the proof is learned from the proof [10, Prop. 3.4] for polynomials, using the *Goursat transform* and the Pólya theorem for homogeneous matrix polynomials established in [8].

Let  $\mathbf{F}$  be as in Theorem 3.1. The *Goursat transform* of  $\mathbf{F}$  is defined by

$$\tilde{\mathbf{F}}(x) := (1+x)^d \mathbf{F}\left(\frac{1-x}{1+x}\right).$$

**LEMMA 3.1.1.**  *$\deg(\tilde{\mathbf{F}}) = d$ ,  $\tilde{\mathbf{F}}_d > 0$ , and  $\tilde{\mathbf{F}}(t) > 0$  for all  $t \in [0, +\infty)$ , where  $\tilde{\mathbf{F}}_d$  denotes the leading coefficient of the matrix polynomial  $\tilde{\mathbf{F}}$ .*

*Proof.* Let  $\mathbf{F}(x) = \sum_{i=0}^d \mathbf{F}_i x^i$ , with  $\mathbf{F}_d \neq 0$ . Then we can write  $\tilde{\mathbf{F}}$  as

$$\tilde{\mathbf{F}}(x) = \sum_{i=0}^d \mathbf{F}_i (1-x)^i (1+x)^{d-i}.$$

The leading coefficient of  $\tilde{\mathbf{F}}$  is

$$\tilde{\mathbf{F}}_d := \sum_{i=0}^d (-1)^i \mathbf{F}_i = \mathbf{F}(-1)$$

which is positive definite by the assumption of  $\mathbf{F}$ .

For every  $t \in [0, +\infty)$ ,  $x := \frac{1-t}{1+t} \in (-1, 1]$ . Observe that  $t = \frac{1-x}{1+x}$ , and hence

$$\tilde{\mathbf{F}}(t) = (1+t)^d \mathbf{F}(x).$$

It follows from the assumption of  $\mathbf{F}$  that  $\tilde{\mathbf{F}}(t) > 0$ .  $\square$

LEMMA 3.1.2. Let  $\mathbf{G} \in M_t(\mathbb{R}[x])$  be a matrix polynomial of degree  $d$  with a positive definite leading coefficient  $\mathbf{G}_d$ . If  $\mathbf{G}(x) > 0$  for all  $x \in [0, +\infty)$  then there exists a positive integer  $N$  and positive definite matrices  $\mathbf{H}_i$ ,  $i = 0, \dots, N + d$  such that

$$(1 + x)^N \mathbf{G}(x) = \sum_{i=0}^{N+d} \mathbf{H}_i x^i.$$

*Proof.* Assume  $\mathbf{G}(x) = \sum_{i=0}^d \mathbf{G}_i x^i$ . Denote by  $\Delta$  the standard simplex in  $\mathbb{R}^2$

$$\Delta = \{(x, y) \in \mathbb{R}^2 \mid x \geq 0, y \geq 0, x + y \leq 1\}.$$

Let  $\mathbf{G}^h(x, y) \in M_t(\mathbb{R}[x, y])$  be the homogenization of  $\mathbf{G}$  with respect to the new variable  $y$ , defined by

$$\mathbf{G}^h(x, y) := \sum_{i=0}^d \mathbf{G}_i x^i y^{d-i}.$$

For any  $(x, y) \in \Delta$ , if  $y = 0$ , we have  $x = 1$ , then

$$\mathbf{G}^h(1, 0) = \mathbf{G}_d > 0.$$

If  $y > 0$ , we have

$$\mathbf{G}^h(x, y) = y^d \mathbf{G}\left(\frac{x}{y}\right) > 0,$$

since  $\frac{x}{y} \in [0, +\infty)$ . In summary, we have  $\mathbf{G}^h(x, y) > 0$  for all  $(x, y) \in \Delta$ .

It follows from the Pólya theorem for homogeneous matrix polynomials proved by Scherer and Hol [8, Theorem 3] that there exist a positive integer  $N$  and positive definite matrices  $\mathbf{G}_{k,l}$ , with  $k, l \in \mathbb{N}_0$ ,  $k + l = 0, \dots, N + d$  such that

$$(x + y)^N \mathbf{G}^h(x, y) = \sum_{k+l=N+d} \mathbf{G}_{k,l} x^k y^l. \tag{3.1}$$

Substituting  $y = 1$  in both sides of the expression (3.1), observing that  $\mathbf{G}^h(x, 1) = \mathbf{G}(x)$ , we get

$$(1 + x)^N \mathbf{G}(x) = \sum_{k+l=N+d} \mathbf{G}_{k,l} x^k = \sum_{k=0}^{N+d} \mathbf{G}_{k,N+d-k} x^k.$$

For each  $k = 0, \dots, N + d$ , the matrices  $\mathbf{H}_k := \mathbf{G}_{k,N+d-k}$  satisfy conclusion of the lemma.  $\square$

Now we are ready to prove Theorem 3.1.

*Proof of Theorem 3.1.* Let  $\mathbf{F}$  satisfy the hypothesis of the theorem. It follows from Lemma 3.1.1 that the Goursat transform  $\tilde{\mathbf{F}}$  satisfy the following properties

- $\deg(\tilde{\mathbf{F}}) = d$ ;
- $\tilde{\mathbf{F}}_d > 0$ ;



- $\tilde{\mathbf{F}}(t) > 0$  for all  $t \in [0, +\infty)$ .

It follows from Lemma 3.1.2 that there exist a positive integer  $N$  and positive definite matrices  $\mathbf{H}_i, i = 0, \dots, N+d$  such that

$$(1+t)^N \tilde{\mathbf{F}}(t) = \sum_{i=0}^{N+d} \mathbf{H}_i t^i, \quad \forall t \in [0, +\infty). \tag{3.2}$$

For any  $x \in (-1, 1]$ , there exists a unique  $t \in [0, +\infty)$  such that  $x = \frac{1-t}{1+t}$ . Then

$$t = \frac{1-x}{1+x}, \quad (1+t)^{-1} = \frac{1+x}{2}, \quad \text{and } \tilde{\mathbf{F}}(t) = (1+t)^d \mathbf{F}(x).$$

Substituting into (3.2), we get

$$(1+t)^N (1+t)^d \mathbf{F}(x) = \sum_{i=0}^{N+d} \mathbf{H}_i \left(\frac{1-x}{1+x}\right)^i.$$

It follows that

$$\begin{aligned} \mathbf{F}(x) &= (1+t)^{-N+d} \sum_{i=0}^{N+d} \mathbf{H}_i \left(\frac{1-x}{1+x}\right)^i \\ &= \frac{1}{2^{N+d}} (1+x)^{N+d} \sum_{i=0}^{N+d} \mathbf{H}_i \left(\frac{1-x}{1+x}\right)^i \\ &= \sum_{i=0}^{N+d} \left(\frac{1}{2^{N+d}} \mathbf{H}_i\right) (1-x)^i (1+x)^{N+d-i}. \end{aligned}$$

Hence the matrices  $\mathbf{G}_i := \frac{1}{2^{N+d}} \mathbf{H}_i, i = 0, \dots, N+d$ , satisfy the conclusion of the theorem.  $\square$

It follows from the Haviland theorem for matrix polynomials (Theorem 2.1) and Theorem 3.1 the following solution of the tracial  $[-1, 1]$ -moment problem.

**COROLLARY 3.1.1.** *Let  $\mathcal{L}$  be a real-valued linear functional on  $M_t(\mathbb{R}[x])$ . Then the following are equivalent.*

- (1)  $\mathcal{L}$  is a tracial  $[-1, 1]$ -moment functional.
- (2)  $\mathcal{L}((1+x)^k (1-x)^l \mathbf{G}) \geq 0$  for all positive definite matrices  $\mathbf{G} \in M_t(\mathbb{R})$  and  $k, l \in \mathbb{N}_0$ .

*Proof.* If (1) holds, for each  $\mathbf{G}$  positive definite and  $k, l \in \mathbb{N}_0$ , since  $(1+x)^k (1-x)^l \mathbf{G} \geq 0$  on  $[-1, 1]$ , it follows from Theorem 2.1, (1)  $\Rightarrow$  (2), that  $\mathcal{L}((1+x)^k (1-x)^l \mathbf{G}) \geq 0$ . Hence we have (1)  $\Rightarrow$  (2).

Conversely, assume (2) holds. Then for each  $\mathbf{F} \in S_t(\mathbb{R}[x])$  of degree  $d$ ,  $\mathbf{F} > 0$  on  $[-1, 1]$ , it follows from Theorem 3.1 that

$$\mathbf{F}(x) = \sum_{i=0}^{N+d} \mathbf{G}_i(1+x)^i(1-x)^{N+d-i}$$

for some  $N > 0$  and  $\mathbf{G}_i > 0$  for all  $i = 0, \dots, N+d$ . Then

$$\mathcal{L}(\mathbf{F}) = \sum_{i=0}^{N+d} \mathcal{L}(\mathbf{G}_i(1+x)^i(1-x)^{N+d-i}) \geq 0.$$

It follows from Theorem 2.1, (4)  $\Rightarrow$  (1), that  $\mathcal{L}$  is a tracial  $[-1, 1]$ -moment functional, i.e. (2)  $\Rightarrow$  (1). The proof is complete.  $\square$

As a consequence of this result, we obtain the following solution of the tracial  $[0, 1]$ -moment problem, which was also solved by Cimpric and Zalar [3, Coro. 1], using sums of Hermitian squares.

**COROLLARY 3.1.2.** *Let  $\mathcal{L}$  be a real-valued linear functional on  $M_t(\mathbb{R}[x])$ . Then the following are equivalent.*

- (1)  $\mathcal{L}$  is a tracial  $[0, 1]$ -moment functional.
- (2)  $\mathcal{L}(x^k(1-x)^l \mathbf{G}) \geq 0$  for all positive definite matrices  $\mathbf{G} \in M_t(\mathbb{R})$  and  $k, l \in \mathbb{N}_0$ .

*Proof.* Observe that there is a bijection from the interval  $[-1, 1]$  onto the interval  $[0, 1]$  given by

$$x \mapsto \frac{x+1}{2}, \forall x \in [-1, 1].$$

Using this bijection, it follows from Theorem 3.1 that for  $\mathbf{F} \in S_t(\mathbb{R}[x])$  of degree  $d$ , if  $\mathbf{F} > 0$  on  $[0, 1]$  then

$$\mathbf{F}(x) = \sum_{i=0}^{N+d} \mathbf{G}_i x^i (1-x)^{N+d-i}$$

for some  $N > 0$  and  $\mathbf{G}_i > 0$  for all  $i = 0, \dots, N+d$ . Then the result follows from this representation and Theorem 2.1.  $\square$

For the sequential matrix-valued  $[-1, 1]$ -problem, we have the following solution.

**COROLLARY 3.1.3.** *Let  $\mathbf{S} = (\mathbf{S}_i)_{i \in \mathbb{N}_0}$  be a sequence of symmetric matrices in  $M_t(\mathbb{R})$ . Then  $\mathbf{S}$  is a matrix-valued  $[-1, 1]$ -moment sequence if and only if*

$$\sum_{i=0}^k \sum_{j=0}^l (-1)^j \binom{k}{i} \binom{l}{j} \text{tr}(\mathbf{G} \mathbf{S}_{i+j}) \geq 0$$

for all positive definite matrices  $\mathbf{G} \in M_t(\mathbb{R})$  and  $k, l \in \mathbb{N}_0$ .

*Proof.* Let  $\mathcal{L}_{\mathbf{S}}$  be the Riesz functional on  $M_t(\mathbb{R}[x])$  associated to the sequence  $\mathbf{S}$ . For each  $\mathbf{G} \in S_t(\mathbb{R}[x])$ ,  $\mathbf{G} > 0$ , and  $k, l \in \mathbb{N}_0$ , we have

$$\begin{aligned} \mathcal{L}_{\mathbf{S}}((1+x)^k(1-x)^l\mathbf{G}) &= \mathcal{L}_{\mathbf{S}}\left(\sum_{i=0}^k \sum_{j=0}^l (-1)^j \binom{k}{i} \binom{l}{j} x^{i+j}\mathbf{G}\right) \\ &= \sum_{i=0}^k \sum_{j=0}^l (-1)^j \binom{k}{i} \binom{l}{j} \mathcal{L}_{\mathbf{S}}(x^{i+j}\mathbf{G}) \\ &= \sum_{i=0}^k \sum_{j=0}^l (-1)^j \binom{k}{i} \binom{l}{j} \text{tr}(\mathbf{G}\mathbf{S}_{i+j}). \end{aligned}$$

By Proposition 1,  $\mathbf{S}$  is a matrix-valued  $[-1, 1]$ -moment sequence if and only if  $\mathcal{L}_{\mathbf{S}}$  is a tracial  $[-1, 1]$ -moment functional. It follows from Corollary 3.1.1 that this is equivalent to  $\mathcal{L}_{\mathbf{S}}((1+x)^k(1-x)^l\mathbf{G}) \geq 0$  for all positive definite matrices  $\mathbf{G} \in M_t(\mathbb{R})$  and  $k, l \in \mathbb{N}_0$ . This implies the result.  $\square$

Similarly, using Corollary 3.1.2, we have the following solution of the sequential matrix-valued  $[0, 1]$ -moment problem.

**COROLLARY 3.1.4.** *Let  $\mathbf{S} = (\mathbf{S}_i)_{i \in \mathbb{N}_0}$  be a sequence of symmetric matrices in  $M_t(\mathbb{R})$ . Then  $\mathbf{S}$  is a matrix-valued  $[0, 1]$ -moment sequence if and only if*

$$\sum_{i=0}^k (-1)^i \binom{k}{i} \text{tr}(\mathbf{G}\mathbf{S}_{i+l}) \geq 0$$

for all positive definite matrices  $\mathbf{G} \in M_t(\mathbb{R})$  and  $k, l \in \mathbb{N}_0$ .

#### 4. Tracial moment problems on the hypercube $[-1, 1]^n$

In this section we consider a convex, compact polyhedron  $K \subseteq \mathbb{R}^n$  with non-empty interior. Assume that  $K$  is the basic closed semi-algebraic set defined by linear polynomials  $L_1, \dots, L_m \in \mathbb{R}[X] := \mathbb{R}[X_1, \dots, X_n]$ , i.e.

$$K = \{x \in \mathbb{R}^n \mid L_1(x) \geq 0, \dots, L_m(x) \geq 0\}.$$

The following version of Handelman’s Positivstellensatz for matrix polynomials was proved by the author in [7].

**THEOREM 4.1.** *Let  $\mathbf{F} \in S_t(\mathbb{R}[X])$  be a matrix polynomial of degree  $d > 0$ . If  $\mathbf{F}(x) > 0$  for all  $x \in K$ , then there exist a positive integer  $N$  and positive definite matrices  $\mathbf{G}_{\alpha} \in M_t(\mathbb{R})$ ,  $\alpha \in \mathbb{N}_0^m$ ,  $|\alpha| = 0, \dots, N + d$ , such that*

$$\mathbf{F}(X) = \sum_{|\alpha|=0}^{N+d} \mathbf{G}_{\alpha} L_1^{\alpha_1} \dots L_m^{\alpha_m}.$$

Applying Theorem 4.1 for the linear polynomials

$$l_1 = 1 + X_1, l_2 = 1 - X_1, \dots, l_{2n-1} = 1 + X_n, l_{2n} = 1 - X_n$$

we obtain the following representation for matrix polynomials positive definite on the hypercube  $[-1, 1]^n$ .

**COROLLARY 4.1.1.** *Let  $\mathbf{F} \in S_t(\mathbb{R}[X])$  be a matrix polynomial of degree  $d > 0$ . If  $\mathbf{F}(x) > 0$  for all  $x \in [-1, 1]^n$ , then there exist a positive integer  $N$  and positive definite matrices  $\mathbf{G}_\alpha \in M_t(\mathbb{R})$ ,  $\alpha \in \mathbb{N}_0^{2n}$ ,  $|\alpha| = 0, \dots, N + d$ , such that*

$$\mathbf{F}(X) = \sum_{|\alpha|=0}^{N+d} \mathbf{G}_\alpha l_1^{\alpha_1} l_2^{\alpha_2} \dots l_{2n-1}^{\alpha_{2n-1}} l_{2n}^{\alpha_{2n}}.$$

Now, applying the Haviland theorem for matrix polynomials (Theorem 2.1) and Corollary 4.1.1 we obtain the following solution of the tracial  $[-1, 1]^n$ -moment problem.

**COROLLARY 4.1.2.** *Let  $\mathcal{L}$  be a real-valued linear functional on  $M_t(\mathbb{R}[X])$ . Then the following are equivalent.*

- (1)  $\mathcal{L}$  is a tracial  $[-1, 1]^n$ -moment functional.
- (2)  $\mathcal{L}(l_1^{\alpha_1} l_2^{\alpha_2} \dots l_{2n-1}^{\alpha_{2n-1}} l_{2n}^{\alpha_{2n}} \mathbf{G}) \geq 0$  for all positive definite matrices  $\mathbf{G} \in M_t(\mathbb{R})$  and  $\alpha = (\alpha_1, \dots, \alpha_{2n}) \in \mathbb{N}_0^{2n}$ .

Observe that the solution of the tracial  $K$ -moment problem, where  $K$  is a convex, compact polyhedron in  $R^n$  with non-empty interior, can be solved by the same way, using the linear polynomial  $L_i$  instead of  $l_j$  in Corollary 4.1.2.

Finally, applying Proposition 1 and Corollary 4.1.2 we obtain the following solution of the sequential matrix-valued  $[-1, 1]^n$ -moment problem.

**COROLLARY 4.1.3.** *Let  $\mathbf{S} = (\mathbf{S}_\alpha)_{\alpha \in \mathbb{N}_0^n}$  be a sequence of symmetric matrices in  $M_t(\mathbb{R})$ . Then  $\mathbf{S}$  is a matrix-valued  $[-1, 1]^n$ -moment sequence if and only if*

$$\sum_{\beta \in \mathbb{N}_0^{2n}, \beta \leq \alpha} (-1)^{e(\alpha)} \binom{\alpha}{\beta} \text{tr}(\mathbf{G}\mathbf{S}_\beta) \geq 0$$

for all positive definite matrices  $\mathbf{G} \in M_t(\mathbb{R})$  and  $\alpha \in \mathbb{N}_0^{2n}$ .

Here, for  $\alpha = (\alpha_1, \dots, \alpha_{2n})$  and  $\beta = (\beta_1, \dots, \beta_{2n})$ ,  $\beta \leq \alpha$  means  $\beta_i \leq \alpha_i$  for all  $i = 1, \dots, 2n$ ;  $\binom{\alpha}{\beta} := \binom{\alpha_1}{\beta_1} \dots \binom{\alpha_{2n}}{\beta_{2n}}$ ; and  $e(\alpha) := \sum_{i=1}^n \alpha_{2i}$ .

*Proof.* The proof is similar to that of Corollary 3.1.3, using Proposition 1, Corollary 4.1.2 and the identity

$$(1 + X_1)^{\alpha_1} (1 - X_1)^{\alpha_2} \dots (1 + X_n)^{\alpha_{2n-1}} (1 - X_n)^{\alpha_{2n}} = \sum_{\beta \in \mathbb{N}_0^{2n}, \beta \leq \alpha} (-1)^{e(\alpha)} \binom{\alpha}{\beta} X^\beta,$$

where  $\alpha = (\alpha_1, \dots, \alpha_{2n})$ .  $\square$

*Acknowledgements.* The author would like to express his sincere gratitude to Prof. Konrad Schmüdgen for his series of lectures on moment problems given at the Vietnam Institute for Advanced Study in Mathematics (VIASM), which motivated him to study matrix-valued moment problems. This paper was finished during a visit of the author to VIASM. He thanks VIASM for financial support and hospitality.

#### REFERENCES

- [1] S. BURGDORF AND I. KLEP, *The truncated tracial moment problem*, J. Oper. Theory **68** (1) (2012), 141–163.
- [2] S. BURGDORF, I. KLEP AND J. POVH, *The tracial moment problem and trace-optimization of polynomials*, Math. Prog. **137**, Issue 1–2 (2013), 557–578.
- [3] J. CIMPRIČ, A. ZALAR, *Moment problems for operator polynomials*, J. Math. Anal. Appl. **401** (1) (2013), 307–316.
- [4] H. DETTE AND W.J. STUDDEN, *Matrix measures, moment spaces and Favard's theorem for the interval  $[0, 1]$  and  $[0, +\infty)$* , Linear Algebra Appl. **345** (2002), 169–193.
- [5] D. P. KIMSEY, *Matrix-valued moment problems*, Ph. D. dissertation, Drexel University, Philadelphia, PA, 2011.
- [6] D. P. KIMSEY AND H. J. WOERDEMAN, *Truncated matrix-valued moment problems on  $\mathbb{R}^d, \mathbb{C}^d$ , and  $\mathbb{T}^d$* , Trans. Amer. Math. Soc. **365** (10) (2013), 5393–5430.
- [7] C.-T. LÊ, T. H. B. DU, *Handelman's Positivstellensatz for polynomial matrices positive definite on polyhedra*, Positivity **22** (3) (2018), 449–460.
- [8] C. W. SCHERER, C. W. J. HOL, *Matrix sum-of-squares relaxations for robust semi-definite programs*, Math. Program. **107** no. 1–2, Ser. B (2006), 189–211.
- [9] K. SCHMÜDGEN, *The  $K$ -moment problem for compact semi-algebraic sets*, Math. Ann. **289** (1) (1991), 203–206.
- [10] K. SCHMÜDGEN, *The moment problem*, Springer, 2017.

(Received August 1, 2019)

Cong Trinh Le  
 Department of Mathematics and Statistics  
 Quy Nhon University  
 170 An Duong Vuong, Quy Nhon, Vietnam  
 e-mail: lecongtrinh@qnu.edu.vn